



The Apples Of New York (1905)



**S. A. Beach
N. O. Booth**



Digitized by the Internet Archive
in 2022 with funding from
Kahle/Austin Foundation

The Apples Of New York

S. A. Beach

Printing Statement:

Due to the very old age and scarcity of this book, many of the pages may be hard to read due to the blurring of the original text, possible missing pages, missing text, dark backgrounds and other issues beyond our control.

Because this is such an important and rare work, we believe it is best to reproduce this book regardless of its original condition.

Thank you for your understanding.

SB
363
B35a
v. 2

INDEX TO ILLUSTRATIONS.

HALF-TONE PLATES.

	FACING PAGE.
Admirable	2
Arctic	8
Blenheim	22
Cabashea	28
Cox Orange	42
Cranberry Pippin	44
Early Harvest	50
Early Joe	52
Early Ripe	54
Fall Orange	60
Gladstone	76
Golden Sweet	82
Haas	90
Hawley	94
Hoadley	100
Hook	102
Jefferis	108
Jersey Sweet	110
Judson	112
Landsberg	118
Late Strawberry	120
Longfield	122
Lowell	128
McMahon	136
Magog	138
Ohio Pippin	148
Parry White	156
Patten	158
Pomona	164
Porter	166
Pumpkin Russet	170
Sharp	198
Sops of Wine	202
Sour Bough	204
Tetofsky	220
Workaroe	246
Montreal Beauty (Crab)	262

COLOR PLATES.

Alexander	4
Benoni	16
Bismarck	20

COLOR PLATES — Continued.

	FACING PAGE.
Champlain	30
Chenango	34
Collamer	36
Constantine	40
Cranberry Pippin	44
Detroit Red	46
Dudley	48
Early Harvest	50
Early Strawberry	56
Fall Pippin	62
Fameuse	66
Fanny	68
Fishkill	70
Gravenstein	84
Keswick	116
Lee Sweet (section).	136
Lee Sweet (whole fruit), see Volume I.	230
McIntosh	132
McLellan (Section).	136
McLellan (whole fruit).	134
Maiden Blush	140
Mother	144
Munson	146
Oldenburg	150
Pease	160
Primate	168
Pumpkin Sweet	172
Red Astrachan	178
Red June	180
Ribston	184
Shiawassee	200
Stump	208
Sweet Bough	216
Twenty Ounce	228
Victoria	232
Wealthy	236
Williams	242
Wolf River	244
Yellow Transparent	248
Excelsior (Crab).	254
Hyslop (Crab).	256
Large Red Siberian (Crab).	258
Martha (Crab).	260
Red Siberian (Crab).	264
Transcendent (Crab).	266

THE APPLES OF NEW YORK.

ADIRONDACK.

REFERENCE. 1. Taylor, *U. S. Pom. Rpt.*, 1893:285.

SYNONYMS. None.

This variety is said to have originated in Clinton county. We have received no reports of its being grown outside of the locality of its origin. Taylor gives the following description of it. (1). "Roundish, conical; regular, of medium size, with smooth surface, becoming glossy when rubbed; color rich yellow, washed and striped with red; dots small, straw color, slightly elevated; cavity large, round, deep, flaring; stem of medium length and thickness, slightly knobbed; basin small, nearly round, very shallow, with convex sides, slightly and regularly ribbed and downy; calyx segments rather small, meeting; eye small, closed. Skin thin, tough; core large, broad, heart-shaped, moderately open, clasping; seeds numerous, oval, plump, grayish brown; flesh yellowish white, granular, rather dry, tender; flavor mild subacid; quality good. Season, October to January in Clinton county, New York. Tree a good grower; resembles Baldwin in its bearing habit; hardy at its place of origin. This variety is said to have originated from seed of Westfield (Seek-No-Further) crossed with Hubbardston. It is a promising early winter apple for Northern New York and New England."

ADMIRABLE.

REFERENCES. 1. Downing, 1859:354. 2. Leroy, 1873:813. fig. 3. Hogg, 1884:214. 4. *N. Y. Sta. An. Rpt.*, 8:339, 348. 1889. 5. Beach, *ib.*, 11:589, 595. 1892. 6. Beach and Clark, *N. Y. Sta. Bul.*, 248:110. 1904.

SYNONYMS. ADMIRABLE (6). SMALL'S ADMIRABLE (1, 2, 3, 4, 5). *Small Admirable* (6).

Fruit green or yellow, not particularly attractive. At its best it is very good for dessert use, but as grown at this Station it usually ranks but fair to good in quality and the fruit is very apt to show imperfect spots in the flesh which are evidently due to some physiological defect. In England it is regarded as an excellent kitchen and dessert apple and the tree is said to be an immense bearer and well adapted for dwarf culture (3). So far as tested at this Station the tree has been an annual cropper and very productive, often yielding full crops. A portion of the fruit may

sometimes be kept through the winter, but ordinarily the season for this variety is November and December (6). Not recommended for cultivation in New York.

Historical. This variety originated in England (2, 3).

TREE.

Tree dwarfish with short, stout, slow-growing branches. *Form* rather flat, spreading and somewhat drooping. *Twigs* short, somewhat curved, stout to rather slender; internodes short to above medium. *Bark* dark brown tinged with olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* numerous, small to medium, oval, slightly raised. *Buds* deeply set in bark, medium to small, broad, plump, obtuse to somewhat acute, appressed or free, pubescent.

FRUIT.

Fruit quite uniform in size and shape for any particular crop but varies under different conditions from below medium to above. *Form* oblate conic to roundish conic, pretty regular but sometimes indistinctly ribbed. *Stem* medium to long, slender, pubescent. *Cavity* acute, moderately deep to deep, broad, smooth or thinly russeted. *Calyx* closed or open; lobes reflexed. *Basin* moderately shallow, rather narrow, sometimes abrupt, somewhat furrowed and wrinkled.

Skin varies from pale green to attractive lemon-yellow, sometimes with brownish blush. *Dots* numerous, small, light or russet.

Calyx tube rather narrow, elongated funnel-form. *Stamens* median.

Core rather large, somewhat abaxile; cells open; core lines clasping. *Carbels* long, obovate. *Seeds* rather large, long, plump, pointed, medium brown.

Flesh whitish, firm, moderately coarse, crisp, quite juicy, mild subacid, very aromatic, good to very good in quality in well grown fruit.

Season October to January.

ALBION.


Fruit intermediate in type between Fall Pippin and Lowell. In color it resembles Fall Pippin but in form it is more like Lowell. *Season* October to late fall or early winter. We have been unable to learn the origin of this variety. It is but little cultivated in this State.

FRUIT.

Fruit large to very large. *Form* roundish oblong, irregular. *Stem* short to medium, slender. *Cavity* acuminate, deep, medium to wide, russeted. *Calyx* small to medium, tightly closed; lobes short, narrow, acute. *Basin* shallow, medium to narrow in width, obtuse, furrowed and wrinkled.

Skin smooth, waxy, pale yellow mingled with green. *Dots* green or russet or submerged and whitish.

Calyx tube narrow, cone-shape to funnel-form. *Stamens* marginal.



ADMIRABLE

Core large, abaxile; cells open or partly closed; core lines clasping. Car-pels elongated, ovate, pointed, mucronate. Seeds few, medium to large, rather narrow, long, acute.

Flesh moderately coarse, very tender, juicy, mild subacid, good.

ALEXANDER.

REFERENCES. 1. Forsyth, 1824:131. 2. *London Hort. Soc. Cat.*, 1831: No. 10. 3. Kenrick, 1832:65. 4. Floy-Lindley, 1833:11. 5. *Mag. Hort.*, 1:395. 1835. 6. Downing, 1845:79. fig. 7. Thomas, 1849:146. 8. Cole, 1849:118. 9. Emmons, *Nat. Hist. N. Y.*, 3:32. 1851. 10. Barry, 1851:282. 11. Elliott, 1854:120. 12. Gregg, 1857:41. 13. Hooper, 1857:11. 14. *Am. Pom. Soc. Cat.*, 1862. 15. Warder, 1867:510. 16. Fitz, 1872:152. 17. Leroy, 1873:333. fig. 18. *Ia. Hort. Soc. Rpt.*, 1879:538. 19. Downing, 1881:11 index, app. 20. Hogg, 1884:72. 21. Roach, *Montreal Hort. Soc. Rpt.*, 1886-87:27. 22. Wickson, 1889:244. 23. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 24. Bailey, *An. Hort.*, 1892:234. 25. Hoskins, *Rural N. Y.*, 53:278. 1894. 26. Gaucher, 1894: No. 14. col. pl. 27. Dempsey, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. 28. Burrill and McCluer, *Ill. Sta. Bul.*, 45:312. 1896. 29. Stinson, *Ark. Sta. Bul.*, 43:102. 1896. 30. Waugh, *Ut. Sta. Bul.*, 61:29. 1897. 31. *Can. Hort.*, 20:283. 1897. 32. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 5:5. 1898. figs. 33. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 34. Craig, *Cyclopedia of Amer. Hort.*, 3:1404. 1901. 35. Hansen, *S. D. Sta. Bul.*, 76:22. 1902. 36. Munson, *Me. Sta. Rpt.*, 1902:83, 85, 86. 37. Budd-Hansen, 1903:36. fig. 38. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:36. 1903. 39. Beach and Clark, *N. Y. Sta. Bul.*, 248:111. 1904. 40. Scriber, *Can. Hort.*, 28:248. 1905.

SYNONYMS. *Albertin* (17, 26). *ALEXANDER* (1, 2, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 19, 21, 22, 23, 24, 27, 28, 29, 30, 31, 32, 34, 35, 36, 37, 38, 39, 40). *Alexander* (34, 26). *Alexander the First* (31). *Alexandre* (17, 26). *Aporta* (3, 4, 6, 9, 13, 17, 20, 26, 31). *Aporta Nalivia* (26). *Aubertin* (17, 26). *Beauty of Queen* (17, 26). *Belle d'Orleans* (17, 26). *Comte Woronzoff* (17, 26). *Corail* (17, 26). *English King* (19, 26). *Empereur Alexandre I* (17, 26). *Empereur Alexandre de Russie* (17, 26). *Empereur de Russie* (17, 26). *EMPEROR ALEXANDER* (3, 4, 20, 33). *Emperor Alexander* (2, 6, 9, 10, 11, 13, 26, 32). *Fini d'Automne* (17, 26). *Grand Alexander* (26). *GRAND-ALEXANDRE* (17). *Grand Alexandre* (26). *Gros-Alexandre* (17, 26). *Jolly Gentleman* (26). *KAISER ALEXANDER* (26). *Korallen Apfel* (26). *Phanix* (26). *Phönix* (17, 26). *Pomona Britannica* (17, 26). *Président Napoléon* (17, 26). *Russian Emperor* (2, 6, 9, 13, 20, 26). *Stoke Tulip* (26). *Wolf River* incorrectly (39). *Wunderapfel* (26).

Alexander is a typical representative of the class of Russian apples commonly known as the Aport group. Fruit very large, attractive red or striped, coarse in texture, medium to good in quality, suitable for culinary rather than for dessert use. The fruit is apt to crack and decay about the stem and calyx and often becomes discolored where it is chafed by constantly rubbing

against some twig or branch; there is also a considerable loss from premature dropping of the fruit. Notwithstanding these faults many fruit growers now regard Alexander favorably as a commercial variety as in some markets there is a strong demand for the fruit at good prices. It is being used to some extent for export trade (40). Its season begins in September and extends through October or into November. It may be held in cold storage till November. It goes down quickly and as it does not stand heat well before going into storage it should be shipped the day it is picked and under ice (39). As it ripens continuously during a period of from four to six weeks it should have more than one picking. The tree is hardy, vigorous and moderately productive. In some localities it is subject to blight. It can be recommended for planting in commercial orchards to a limited extent. In the West it is now largely supplanted by its Wisconsin seedling Wolf River (34, 36).

Historical. Introduced into England from Russia in 1817 (20). The exact date of the introduction of this variety into America is not known. The Massachusetts Horticultural Society made several importations of European varieties which were distributed among the members of the society. Mr. Manning exhibited what was supposed to be Alexander before the Massachusetts Horticultural Society at its meeting on September 18, 1830. Whether this was Alexander or not, the shipment of varieties of which Alexander was one had evidently been made prior to that date.¹

It has been widely disseminated and is now pretty well known in the apple growing districts from the Atlantic to the Pacific. Thus far it has not been grown to any considerable extent in New York state but at the present time its cultivation is on the increase.



TREE.

Tree large to medium, vigorous to moderately vigorous with long, stout branches. *Form* upright spreading to roundish, open and somewhat inclined to droop after bearing heavy crops. *Twigs* short, curved, stout with large terminal buds; internodes medium. *Bark* brown mingled with olive-green, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* scattering, medium in size, oval, raised. *Buds* medium in size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit large, uniform in size and shape. *Form* roundish conic to slightly oblate conic, regular or approaching broadly angular, symmetrical. *Stem* medium to rather short, moderately thick. *Cavity* acute to acuminate, deep, broad, symmetrical, occasionally lipped, russeted, often with broad, con-

¹ N. E. Farmer, Sept. 24, 1880:78.



ALEXANDER (Reduced Size)

spicuous, outspreading russet rays. *Calyx* medium to large, usually open; lobes medium to short, rather narrow, acute. *Basin* rather small, deep, narrow to nearly medium in width, abrupt, nearly smooth, symmetrical.

Skin moderately thick, tough, smooth, glossy, somewhat waxy, greenish or pale yellow deepening to orange-yellow in the sun, often entirely over-spread with lively red or handsomely striped and splashed with bright carmine. *Dots* inconspicuous, small, scattering. *Prevailing effect* red or striped.

Calyx tube variable, long to short, wide to medium, conical to funnel-shape. *Stamens* median to basal.

Core small, usually axile; cells often not uniformly developed, closed or very slightly open; core lines slightly clasping. *Carpels* elliptical to slightly ovate, emarginate. *Seeds* medium in size, wide, short, rather plump, obtuse to acute.

Flesh nearly white with faint yellow tinge, firm, coarse, moderately crisp, tender, juicy, mild subacid, fair to good.

Season September and October or early November.

AMSTERDAM.

REFERENCES. 1. Downing, 1876:43 app. 2. Ragan, *U. S. B. P. I. Bul.*, 56:26. 1905.

SYNONYMS. AMSTERDAM (2). AMSTERDAM SWEET (1). *Amsterdam Sweet* (2). *Hightop Sweet* incorrectly (1).

Downing describes this fruit as medium in size, pale greenish-yellow, striped and splashed with light and dark bright red; flesh white, half fine, tender, juicy, rather rich, sweet, slightly aromatic; season October and November. Origin Amsterdam, N. Y., on the farm of Joseph Britten (1). We find no account of Amsterdam except the one above mentioned, and have not seen the variety.

ANISIM.

REFERENCES. 1. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:71. 2. Hoskins, *Rural N. Y.*, 52:299. 1893. figs. 3. Greene, *Minn. Sta. Bul.*, 32:240. 1893. 4. Budd, *Ia. Sta. Bul.*, 31:334. 1895. 5. Thomas, 1897:626. 6. Budd, *Ia. Sta. Bul.*, 41:66, 69, 71. 1899. 7. *Am. Pom. Soc. Cat.*, 1899:15. 8. Munson, *Me. Sta. Rpt.*, 1902:83, 86, 95. 9. Hansen, *S. D. Sta. Bul.*, 76:23. 1902. fig. 10. Budd-Hansen, 1903:38. fig.

SYNONYMS. ANISIM (1, 2, 3, 4, 5, 6, 7, 8, 9, 10). *Anisim* of Peterson (9). *Borsdorfer* of Wragg (9). 13 *M* (3, 8. of Budd 9). 14 *M* of Budd (9). *Good Peasant* of Patten (9). *Jonathan* of the North (4). *Swedish Borsdorf* of Patten (9). *Zuzoff* of Tuttle (9).

Hansen (9) reports that this Russian apple is proving very valuable in Minnesota and other parts of the Northwest. It may be of some value in those portions of New York state where superior hardiness is a prime requisite. Hansen's description is given below. "Tree a strong grower in the nursery and orchard and a prodigious bearer; young trees upright, spreading with age; limbs long, slender with a very strong shoulder; leaves

narrow, pointed, dark green. The beautiful color of the fruit attracts favorable attention—Fruit below medium, roundish conical, slightly angular; surface greenish-yellow, covered almost wholly with a beautiful dark crimson, with heavy blue bloom; dots white, minute; cavity regular, acute, usually slightly russeted; stem medium; basin narrow, very shallow, corrugated, sometimes flat; calyx closed. Core closed, clasping; tube short, broad; stamens median, flesh greenish white, with green veins, good. Early winter."

ANTONOVKA.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 8:28, 79, 130. 1881-82. fig. 2. Budd, *Ja. Hort. Soc. Rpt.*, 1882:75. 3. Gibb, *Ib.*, 1883:432. 4. Budd, *Am. Pom. Soc. Rpt.*, 1883:71. 5. *Ib.*, *Ja. Agr. Coll. Bul.*, 1883:22. 6. *Ib.*, 1885:8. 7. Van Deman, *U. S. Pom. Rpt.*, 1886:272. fig. 8. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886 87:72. 9. Budd, *Ja. Agr. Coll. Bul.*, 1890:24. 10. *Can. Hort.*, 13:175, 216. 1890. 11. Budd, *Ja. Sta. Bul.*, 19:537. 1892. 12. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57, 58. 13. Bailey, *Am. Hort.*, 1892:234. 14. *Can. Hort.*, 16:359. 1893. 15. *Ib.*, 17:289, 290, 291. 1894. 16. Troop, *Ind. Sta. Bul.*, 53:124. 1894. 17. *Nat. Nurseryman*, 3:32. 1895. 18. Stinson, *Ark. Sta. Bul.*, 43:104. 1896. 19. Hamilton, *Can. Hort.*, 20:412. 1897. 20. *Am. Pom. Soc. Cat.*, 1897:11. 21. Thomas, 1897:286, 626. fig. 22. Budd, *Ja. Sta. Bul.*, 41:66, 69, 70, 71, 73, 74. 1899. 23. Troop, *Ind. Sta. Rpt.*, 1899:79. 24. Macoun, *Can. Dept. Agr. Rpt.*, 1899:78. 25. Hansen, *S. D. Sta. Bul.*, 76:25. 1902. fig. 26. Farrand, *Mich. Sta. Bul.*, 205:43. 1903. 27. Budd-Hansen, 1903:38. fig.

SYNONYMS. ANTENOVKA (23). ANTONOVKA (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 18, 19, 20, 21, 22, 24, 25, 26, 27). ANTONOWKA (15, 17). Antony (3, 14). Bergamot, 421 (25). Cinnamon (24). German Calville (24). German Calville, 324 (spurious) (25). No. 224 (25). No. 236 (1, 2, 4, 5, 6, 9, 25). Possaris Nalivia (2, 4, 5). Russian Graevenstein, 105 (25). 26 M (1, 2, 4, 5, 9, 25). Vargul, 277 (25).

A Russian fruit of no practical value for this state. Hansen (25) describes it as "large, roundish, irregular, obscurely angular; surface yellow; dots minute, raised, white, suffused; cavity deep, regular, with radiating, often large patch of russet, stem medium; basin abrupt, corrugated or wavy; calyx closed. Core closed; cells ovate, slit; tube funnel-shaped; stamens median; seeds ten to sixteen, small, pointed, plump, a few imperfect; flesh yellow, juicy, sprightly spicy subacid, good. October."

APORT.

REFERENCE. 1. Gibb, *Montreal Hort. Soc. Rpt.*, 8:32. 1881-82.

SYNONYMS. None.

This name is applied to a pretty well defined group of Russian apples. Alexander is the typical variety of this group.

The name Aport has also been applied to a particular Russian variety which resembles Alexander closely.¹

¹ Hansen, *S. D. Sta. Bul.*, 76:26. 1902.

APOINT ORIENT.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 8:73. 1881-82. 2. *Ia. Hort. Soc. Rpt.*, 1882:78. 3. Budd, *Ia. Agr. Coll. Bul.*, 1883:30. 4. *Montreal Hort. Soc. Rpt.*, 9:82. 1883. fig. 5. Budd, *Ia. Agr. Coll. Bul.*, 1885:9. 6. Schroeder, *Montreal Hort. Soc. Rpt.*, 12:72. 1886-87. 7. Budd, *Ia. Agr. Coll. Bul.*, 1890:24. 8. *ib.*, *Ia. Sta. Bul.*, 19:538. 1892. 9. (?) Beach, *N. Y. Sta. An. Rpt.*, 13:579. 1894. 10. Budd, *Ia. Sta. Bul.*, 41:71. 1899. 11. Munson, *Me. Sta. Rpt.*, 1902:83. 12. Hansen, *S. D. Sta. Bul.*, 76:26. 1902. 13. Budd-Hansen, 1903:39.

SYNONYMS. APOINT (1, 2, 3, 4, 5, 6, 7, 11, 12, 13). APOINT ORIENT (8, 10). (APOINT ORIENTAL, 9)? *No. 12 Orel* (7). *No. 252* (1, 2, 3, 4, 7, 11). *23 M* (1, 2, 3). *Oporto* (4).

A Russian apple, large, yellow, mostly covered with mixed red, striped and splashed with dark crimson, very attractive but coarse-grained and inferior in quality. It begins to ripen about the middle of August. The tree comes into bearing rather young and gives full crops in alternate years. Not recommended for New York state.

ARCTIC.

REFERENCES. 1. *N. Y. Sta. An. Rpt.*, 8:347. 1889. 2. Munson, *Me. Sta. Rpt.*, 1896:70. 3. Waugh, *Vt. Sta. Bul.*, 61:29. 1897. 4. *Am. Pom. Soc. Cat.*, 1899:15. 5. Beach, *W. N. Y. Hort. Soc. Rpt.*, 1900:34. 6. Waugh, *Vt. Sta. An. Rpt.*, 14:286. 1901. 7. *Me. Sta. Rpt.*, 1902:83, 89, 95, 96. 8. Budd-Hansen, 1903:39. 9. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:36. 1903.

SYNONYMS. None.

One of the most valuable characteristics of the Arctic is its ability to endure cold climates. It has probably been planted more extensively in Northern New York, New England and Canada than in any other regions. It is reported as being pretty hardy in Central Iowa where the climate is too severe for Baldwin, Rhode Island *Greening* and other varieties of a similar grade of hardiness. Munson (7) states that it is worthy of trial where Baldwin will not succeed. The tree is vigorous. In some districts it has the reputation of being productive, in others it is called a shy bearer. The fruit is very attractive, mild subacid, good but not high in quality. It somewhat resembles Baldwin in size and color, but is more oblate, and the skin is of a somewhat lighter and brighter red than that of Baldwin. The cavity is often marked with outspreading rays of reddish or green russet as in the Baldwin. The dots are round, scattering, whitish, often

areolar and not elongated in the region of the cavity as they often are on the Baldwin. It does not keep so well as the Hubbardston. In Western New York its season may extend from October to February, but when grown farther north is later (6, 7). In some parts of Northern New York it is being grafted over to other sorts. Waugh (6) states that this is also being done in the Isle La Motte region of Vermont, nevertheless he believes it will be grown there in moderate quantities for years to come.

Historical. The Arctic was introduced by Mr. O. K. Gerrish, now of Lakeville, Mass. He states that it originated as a chance seedling in a garden near Cape Vincent, N. Y., about 1862. About 1887 he bought the tree from Mr. John H. Esselyne on whose farm it was growing. After taking propagating wood from it, he destroyed the tree to prevent theft of scions.

TREE.

Tree a moderate grower with long, moderately stout branches. *Form* spreading and open somewhat like Tompkins King. *Twigs* short, stocky; internodes medium length; slightly pubescent near tips. *Bark* dark reddish-brown, streaked and mottled with thin scarf-skin. *Lenticels* scattering, conspicuous, medium to large, roundish to somewhat elongated, raised. *Buds* medium to large, broad, obtuse, flat, free, pubescent. *Leaves* large, dark green, broad and rather thick.

FRUIT.

Fruit above medium to large. *Form* oblate, sometimes roundish conic, often faintly ribbed; pretty uniform in size and shape. *Stem* short and rather thick to medium. *Cavity* moderately shallow to rather deep, broad, usually symmetrical or slightly furrowed, and having outspreading rays of red or green russet. *Calyx* medium to rather large; segments broad, obtuse, closed or partly open. *Basin* abrupt, medium to wide and deep, often compressed or slightly furrowed and corrugated.

Skin nearly smooth, slightly roughened by the light russet or whitish dots; deep yellow or greenish-yellow, often almost wholly covered with a bright red obscuring the stripes of deeper red.

Calyx tube short, conical varying to funnel-form. *Stamens* median.

Core medium, axile or nearly so; cells closed or partly open; core lines clasping the cylinder of the tube. *Carpels* broadly roundish to nearly obcordate, emarginate, slightly tufted. *Seeds* often abortive; when normally developed they are medium to rather large, flat, obtuse, sometimes slightly tufted, dark.

Flesh somewhat tinged with yellow, firm, moderately coarse, crisp, juicy, mild subacid, good in quality.

Uses. Adapted rather for market and culinary uses than for dessert.

Season October to February or later.



ARCTIC

AUGUST.

REFERENCES. 1. *N. Y. Sta. An. Rpt.*, 7:89. 1888. 2. *Me. Sta. Rpt.*, 1893:132. 3. *N. Y. Sta. An. Rpt.*, 13:586. 1894. 4. *Mich. Sta. Bul.*, 118:59. 1895. 5. *Ib.*, 129:39. 1896. 6. *N. Y. Sta. An. Rpt.*, 15:270. 1896. 7. *Mich. Sta. Bul.*, 143:200. 1897. 8. *Ib.*, 152:219. 1898. 9. *Ib.*, 205:47. 1903.

SYNONYMS. None.

This hybrid is classed by some as an apple and by others as a crabapple. The tree is hardy, comes into bearing early and is reliably productive. The fruit is medium to small for an apple but very large for a crabapple. It has a slight crabapple flavor and is of fairly good quality for culinary use. Not recommended for planting in New York.

Historical. Originated from seed of Wealthy by Peter M. Gideon, Excelsior, Minn., from whom it was received in 1888 for testing at this Station. It has been tested at experiment stations in different States but it appears to be practically unknown to fruit growers.

TREE.

Tree moderately vigorous. *Form* upright spreading and somewhat drooping, open. *Twigs* short, curved, slender; internodes short. *Bark* clear brown, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, medium in size, oblong, slightly raised. *Buds* medium in size, plump, acute, free, not pubescent.

FRUIT.

Fruit medium to small, occasionally above medium, uniform in size and shape. *Form* roundish oblate to roundish conic, nearly symmetrical, regular or somewhat ribbed. *Stem* rather short to medium in length, moderately slender. *Cavity* acute approaching acuminate, medium to deep, moderately broad, symmetrical, usually not russeted. *Calyx* rather large, closed; lobes long, acute, reflexed. *Basin* moderately shallow to rather deep, moderately wide, somewhat abrupt, slightly furrowed and wrinkled.

Skin pale yellow or greenish, sometimes almost entirely overspread and mottled with rather bright red, striped and splashed with carmine, covered with bloom. *Dots* whitish, small, scattering, inconspicuous. *Prevailing color* red in well colored specimens.

Calyx tube rather small, short, conical. *Stamens* median to nearly basal.

Core medium in size to above, usually axile; cells often unsymmetrical, usually closed, sometimes wide open; core lines clasping. *Carpels* ovate. *Seeds* light brown, medium to above, moderately wide, plump, acute.

Flesh slightly tinged with yellow, half-fine, moderately juicy, breaking, mild subacid, with a slight crabapple flavor; quality fairly good for culinary use.

Season August and early September.

AUGUSTINE.

REFERENCES. 1. *Horticulturist*, 1848 (cited by 5). 2. Downing, 1857:207. 3. Warder, 1867:711. 4. Thomas, 1885:502. 5. Ragan, *U. S. B. P. I. Bul.*, 56:31. 1905.

SYNONYMS. None.

A pleasant flavored dessert apple formerly grown to a very limited extent in some portions of the state but now practically unknown. Fruit medium to rather large, roundish conic or slightly inclined to oblong, yellow splashed and striped with red; flesh moderately juicy to rather dry, not crisp, tender, sweet; season August.

AUTUMN BOUGH.

REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 36. 2. Parsons, *Horticulturist*, 1:209. 1846. 3. *Am. Pom. Soc. Cat.*, 1852. 4. *Mag. Hort.*, 19:68. 1853. 5. Elliott, 1854:66. fig. 6. Downing, 1857:71. 7. *Mag. Hort.*, 25:154. 1859. 8. Warder, 1867:712. 9. Thomas, 1875:198. 10. Downing, 1881:11 index, app. 11. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 12. Bailey, *Am. Hort.*, 1892:234.

SYNONYMS. AUTUMN BOUGH (1, 2, 3, 4, 5, 7, 11, 12). *Autumn Bough* (9). *AUTUMNAL BOUGH* (8). *AUTUMN SWEET BOUGH* (6, 9, 10). *Autumn Sweet Bough* (5). *Fall Bough* (6, 9). *Late Bough* (6, 9). *Montgomery Sweet* (10). *Philadelphia Sweet* (6, 9). *Summer Bellflower* (6).

This is regarded by many as one of the best sweet apples of its season for dessert use and is esteemed also for culinary purposes. The tree is medium in size, upright or roundish, moderately vigorous to vigorous, healthy, long-lived, comes into bearing fairly young and is reliably productive. The fruit hangs well to the tree. It is suitable for local market but it does not ship well. So far as we can learn it is not grown commercially but it is occasionally cultivated for home use and is still listed by some nurserymen.

In 1846 Robert B. Parsons, of Flushing, N. Y., described it as "a very superior fruit, ranking indeed among our best sweet apples, and worthy of extensive cultivation. It is rather large, somewhat of a calville-shape, though with the ribs not quite so prominent as is usual with apples of that class; oblong, diminishing very much to the eye. Skin smooth, pale yellow, with a few scattered dots. Eye of medium size, and very deeply sunken. Stalk rather slender, set in a deep narrow cavity. Flesh white, very tender, and with a rich and sweet, yet sprightly flavor. Ripens from 25th of Eighth month to the 20th of Ninth month. The tree is exceedingly productive, and of very vigorous growth" (2).

AUTUMN STREAKED.

REFERENCES. 1. Gihb, *Am. Pom. Soc. Rpt.*, 1887:50. 2. Lyon, *U. S. Pom. Bul.*, 2:39. 1888. 3. *N. Y. Sta. An. Rpt.*, 8:349. 1889. 4. *Ib.*, 13:579. 1894.

SYNONYMS. AUTUMN STREAKED (1, 2, 3, 4). *Herbst Streifling* (1). *Herbst Streifling* (2). *No. 964* (1).

This fruit approaches the Oldenburg type in some respects. It is of good size and usually attractive in color, sprightly subacid, very good for culinary purposes; season September. The tree is hardy, comes into bearing young and is a good biennial bearer.

Historical. A Russian apple received from T. H. Hoskins, Newport, Vt., in 1888 for testing at this Station (3, 4).

TREE.

Tree moderately vigorous with short, stout branches. *Form* spreading, flat, rather dense. *Twigs* short, curved, stout with large terminal buds; internodes short. *Bark* dull brown, mingled with olive-green, heavily coated with gray scarf-skin; pubescent. *Lenticels* scattering, medium to large, oval, slightly raised. *Buds* prominent, large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit large. *Form* roundish to roundish oblate, somewhat inclined to conic, regular or obscurely ribbed; sides often unequal. *Stem* short to medium, rather slender. *Cavity* medium to rather large, acute to acuminate, moderately deep, rather wide, slightly furrowed, greenish-russet. *Calyx* large, closed or partly open. *Basin* uneven, wide, abrupt, wrinkled.

Skin yellow or pale yellow, shaded, striped and splashed with red and overspread with pinkish bloom. *Prevailing effect* striped red.

Core large, open.

Flesh yellowish, firm, a little coarse, rather crisp, moderately juicy, sprightly subacid, good.

AUTUMN SWAAR.

REFERENCES. 1. *Genesee Farmer*, 1838 (cited by 10). 2. Downing, 1857:115. 3. Hooper, 1857:14. 4. *Am. Pom. Soc. Cat.*, 1862. 5. Warder, 1867:572. *fig.* 6. Downing, 1869:82. 7. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 8. Bailey, *An. Hort.*, 1892:234. 9. Budd-Hansen, 1903:41. 10. Ragan, *U. S. B. P. I. Bul.*, 56:107. 1905.

SYNONYMS. AUTUMN SWAAR (4, 7, 8, 9). AUTUMNAL SWAAR (1, 2, 3, 6). *Autumnal Swaar* (10). FALL SWAAR (10, of the West 5). *Fall Swaar* of West (6, 9, 10).

This belongs to the Fall Orange group and the fruit resembles Fall Orange very closely. It is very good in quality for either dessert or culinary uses. The tree is hardy, vigorous and spreading; not satisfactorily productive (7). It is occasionally found in cultivation in this state but is now seldom or never planted. Its origin is unknown.

FRUIT.

Fruit above medium to medium, sometimes large. *Form* oblate to roundish conic. *Stem* often short, thick and irregularly knobbed. *Cavity* acute, deep, broad, often lipped or irregular, with concentric russet marks and with outspreading russet rays. *Calyx* medium to small, closed or slightly open. *Basin* medium in depth, medium to narrow, abrupt, slightly ridged.

Skin orange-yellow or greenish, in some cases with a decided blush but

not striped, roughened by almost invisible, capillary netted russet lines which become more distinct, larger and concentric about the base and apex. *Dots* conspicuous, irregular, russet or red areolar with russet center. *Prevailing effect* yellow.

Calyx tube funnel-form. *Stamens* median.

Core medium to rather small, nearly axile; cells partly open or closed; core lines clasp the funnel-cylinder. *Carpels* emarginate, somewhat elliptical, tufted. *Seeds* numerous, large to medium, plump, tufted, brown.

Flesh yellow, tender, breaking, juicy, agreeable, mild subacid, decidedly aromatic, sprightly, very good.

Season September.

AUTUMN SWAAR AND FALL ORANGE COMPARED.

The fruit of Fall Orange as compared with that of Autumn Swaar is larger, more inclined to conic, has smaller seeds and a fleshy pistil point projecting into the base of the calyx tube. The flesh is whiter, more acid, less aromatic and less tender. The flesh of well grown and well ripened Autumn Swaar is decidedly yellower, more tender and milder with a very pleasant, peculiarly aromatic flavor.

AUTUMN SWEET SWAAR.

REFERENCES. 1. *Albany Cultivator*, 5:247. 1848. 2. Thomas, 1849:145. 3. Barry, 1851:282. 4. Elliott, 1854:121. 5. Downing, 1857:115. 6. Gregg, 1857:41. 7. Warder, 1867:47t. fig. 8. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 9. Ragan, *U. S. B. P. J. Bul.*, 56:33. 1905.

SYNONYMS. AUTUMN SWAAR (3, 9). *Autumn Sweet* (9). AUTUMN SWEET SWAAR (8). AUTUMNAL SWAAR (1, 2, 4, 6). *Autumnal Sweet* (9). AUTUMNAL SWEET SWAAR (5, 7). *Autumnal Sweet Swaar* (9). *Sweet Golden Pippin* (5, 9). *Sweet Swaar* (2, 3, 5, 6, 7, 9).

In 1848 Thomas described this as one of the finest autumnal sweet apples (1). It is now seldom found in cultivation in this state. Its origin is unknown.

TREE.

Tree moderately vigorous to vigorous, productive. *Form* upright spreading.

FRUIT (1, 2, 4, 6).

Fruit large. *Form* roundish oblate, sometimes slightly ribbed. *Stem* varying from long and slender to thick and fleshy, yellow and red. *Cavity* acute, deep, wavy, green. *Calyx* medium in size, closed. *Basin* shallow, wide, slightly furrowed.

Skin smooth, waxen yellow, sometimes blushed. *Dots* rare, minute.

Core medium in size; cells somewhat open; core lines clasping. *Seeds* numerous, plump, pale.

Flesh whitish tinged with yellow, fine, moderately juicy, very sweet, spicy, agreeable, very good to best.

Season September and October.

BAILEY SPICE.

REFERENCES. 1. *N. Y. Agr. Soc. Trans.*, 1849:350. 2. Bailey, *Horticulturist*, 5:286. 1850. fig. 3. Hovey, *Mag. Hort.*, 16:542. 1850. fig. 4. Barry, 1851:283. 5. Elliott, 1854:121. 6. Downing, 1857:116. 7. Hooper, 1857:14. 8. Gregg, 1857:41. 9. Warder, 1867:712. 10. Thomas, 1875:205.

SYNONYMS. BAILEY SPICE (2, 3, 4, 5, 7, 8). BAILEY'S SPICE (1, 6, 9, 10).

A dessert apple of medium size, light yellow color and subacid, spicy flavor, in season in September and October.

Historical. In 1850 J. W. Bailey, of Plattsburgh, published the following account of the origin of this variety (2, 3). "The original tree is now growing in my grounds, and was planted there fifty years ago by my grandfather, Captain Nathaniel Platt. It is a great bearer, and I think I never knew an apple so invariably fair and perfect as this."

So far as we have learned this variety is no longer planted and is nearly obsolete in New York.

BAKER SWEET.

REFERENCES. 1. Downing, 1857:117. 2. Warder, 1867:712. 3. Thomas, 1875:492.

SYNONYMS. BAKER SWEET (2, 3). BAKER'S SWEET (1). *Late Golden Sweet* (1). *Long Stem Sweet* (1). *Winter Golden Sweet* (1, 3).

A golden yellow apple of good size and attractive appearance. Because it is sweet, not a late keeper and drops readily from the tree it is of little commercial value except where it can be disposed of in local market, notwithstanding that the tree is very productive. It is a good variety for the home orchard where a sweet apple, ripening in late autumn, is desired.

Historical. This is an old variety, formerly much grown in parts of New England (1). It is but little grown in New York state.

TREE.

Tree medium size, only moderately vigorous or a slow grower; branches dark, rather slender, somewhat resembling Jonathan (1). *Form* spreading. *Twigs* rather stout.

FRUIT.

Fruit large to medium, pretty uniform in size and shape. *Form* roundish to oblate, usually regular. *Stem* short to rather long, rather slender. *Cavity* large, acute to acuminate, deep, rather broad, sometimes partly russeted and with outspreading rays, symmetrical. *Calyx* pubescent, medium, closed; lobes broad at base, acute. *Basin* shallow to moderately deep, narrow to above medium in width, somewhat abrupt, a little furrowed.

Skin moderately thin, tough, nearly smooth except for some patches of russet and conspicuous russet dots, good yellow with shade of brownish-red blush on exposed cheek. *Prevailing effect* good yellow.

Calyx tube medium, somewhat funnel-shape. *Stamens* median.

Core above medium to large, abaxile; cells open, sometimes unsymmetrical; core lines meeting. Carpels very broadly ovate to roundish, tufted. Seeds dark, medium to rather small, plump, acute, tufted.

Flesh yellowish, firm, moderately fine, rather tender, rather juicy, very sweet, pleasant, good to very good.

Season October to December.

BANKS.

REFERENCES. 1. Craig and Allen, *Can. Hort.*, 16:420. 1893. fig. 2. *Nova Scotia Fr. Gr. Assn. Rpt.*, 1894:81, 129. 3. Sears, *Can. Hort.*, 22:476. 1899. 4. Caston, *Ont. Fr. Stas. An. Rpt.*, 9:55. 1902. 5. *Am. Pom. Soc. Rpt.*, 1903:166.

SYNONYMS. BANKS (2, 3, 4, 5). BANKS GRAVENSTEIN (2). BANKS RED GRAVENSTEIN (1). *Banks Red Gravenstein* (4). *Red Gravenstein* (3).

In 1903 R. W. Starr, Wolfville, Nova Scotia, presented to the American Pomological Society the following report concerning this variety (5). "A bud sport from Gravenstein, much the same in season and flavor, but bright red, less ribbed, more regular in shape, and generally a little smaller in size. First noticed and propagated by C. E. Banks, of Berwick, Kings Co., N. S. It is well liked and is being quite largely planted." It appears that this sport first came into bearing about 1880 (1). In 1899 Sears (3) remarked, "The Banks or Red Gravenstein is gaining in popularity because it combines with the superior quality of the ordinary Gravenstein the bright red color which people demand who judge the apple by its appearance alone."

Except in the points of difference above noted Banks appears to be identical with Gravenstein and the reader is referred to the description of that variety for a technical account of the tree and fruit. So far as we can learn this variety is not yet planted to any considerable extent in New York.

BEAUTIFUL ARCAD.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 9:104. 1883. 2. *Id. Hort. Soc. Rpt.*, 1883:443. 3. *Id. Agr. Coll. Bul.*, 1885:17. 4. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:15. 5. *Id.*, *Am. Pom. Soc. Rpt.*, 1887:48. 6. Budd, *Rural N. Y.*, 47:692. 1888. 7. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57. 8. Harris, *U. S. Pom. Rpt.*, 1892:274. 9. Thomas, 1897:248. fig. 10. Hansen, *S. D. Sta. Bul.*, 76:28. 1902. 11. Budd-Hansen, 1903:44. 12. Ragan, *U. S. B. P. I. Bul.*, 56:29, 39, 353. 1905.

SYNONYMS. *Arcad Krasivui* (3). *Arkad Krasivui* (5, 12). ARKAD KRASIVUI (1). *Arkad Krasivui* (2, 5, 12). BEAUTIFUL ARCAD (2, 3, 4, 5, 6, 7, 8, 9, 11). BEAUTIFUL ARCADE (10, 11). *Beautiful Arcade* (1, 5, 12). *No. 453* (5, 6, 10, 11, 12).

This is a Russian apple of good medium size, yellow, partly shaded and splashed with red, sweet, in season in August and September. It is considered a desirable variety in portions of the Upper Mississippi valley and in other districts where superior hardiness is a prime requisite.

BEAUTY OF KENT.

REFERENCES. 1. Forsyth, 1803:55. 2. *Ib.*, 1824:93. 3. *London Hort. Soc. Cat.*, 1831:No. 59. 4. Kenrick, 1832:92. 5. Floy-Lindley, 1833:20. 6. Downing, 1845:81. 7. Hovey, *Mag. Hort.*, 14:250. 1848. fig. 8. Kirtland, *Horticulturist*, 2:544. 1848. 9. Thomas, 1849:146. 10. Cole, 1849:114. 11. Emmons, *Nat. Hist. N. Y.*, 3:34. 1851. 12. Barry, 1851:283. 13. Elliott, 1854:167. 14. Hooper, 1857:15. 15. Gregg, 1857:41. 16. Downing, *Horticulturist*, 19:364. 1864. figs. 17. Warder, 1867:584. 18. Fitz, 1872:152. 19. Leroy, 1873:97. fig. 20. Hogg, 1884:17. 21. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 22. Wickson, 1889:245. 23. Bailey, *Am. Hort.*, 1892:234. 24. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 25. Budd-Hansen, 1903:109. 26. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:46. 1903. 27. Garden, 64:239. 1903. fig. bearing nursery trees.

SYNONYMS. BEAUTE DE KENT (19). BEAUTY OF KENT (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 27). *Beauty of Kent* (19, 26). KENT BEAUTY (25, 26). *Kentish Pipkin* (19, of some 3). *Pepin de Kent* (19). *Pippin Kent* (19).

Fruit large, beautiful, showy, suitable for culinary use, in season from late September to November. The tree is large, vigorous, upright, comes into bearing rather young, is a reliable cropper and moderately productive. In England where it originated it is said to do best under garden culture in warm soil and on Paradise stock; grown in clay and other uncongenial soils it loses quality (27). It is but little known in New York.

BELBORODOOSKOE.

REFERENCES. 1. Gibb, *Am. Pom. Soc. Rpt.*, 1887:55. No. 37. 2. *N. Y. Sta. An. Rpt.*, 8:349. 1889. 3. Beach, *Ib.*, 12:599. 1893. 4. Thomas, 1897:265. fig. 5. Hansen, *S. D. Sta. Bul.*, 76:29. 1902.

SYNONYMS. BELBORODOOSKOE (2, 3). BELLERIOVSKOE (4, 5). *Bielborodovska* (1). WHITE BOROMOVKA (1).

A Russian apple, medium to large, pale greenish-yellow, sometimes blushed, coarse, rather juicy, mild subacid to nearly sweet, good; season August. It does not appear to be worthy the attention of New York fruit growers.

BENNINGER.

REFERENCES. 1. Churchill, *N. Y. Sta. An. Rpt.*, 8:355. 1889. 2. Bailey, *Am. Hort.*, 1892:235. 3. Heiges, *U. S. Pom. Rpt.*, 1894:17. 4. Beach, *N. Y. Sta. An. Rpt.*, 15:270. 1896. 5. Taylor, *Am. Pom. Soc. Rpt.*, 1897:36.

SYNONYMS. BENNINGER (2). BENNINGER (1, 3, 4, 5).

A pleasant-flavored dessert apple of good medium size and attractive appearance; in season during late August and September. It is too mild in flavor to be very desirable for culinary purposes. The tree is a pretty good grower, comes into bearing young and is productive.

Historical. Originated about 1830 on the farm of Uhlie Benninger near Slatington, Lehigh county, Pa. In that region it is said to be a good grower and a reliable and abundant cropper (4).

TREE.

Tree moderately vigorous with short stout branches. *Form* spreading, open. *Twigs* short, curved, stout with large terminal buds; internodes medium. *Bark* clear brownish tinged with olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* conspicuous, quite numerous, medium in size, oblong, not raised. *Buds* deeply set in bark, medium in size, broad, obtuse, appressed, pubescent.

FRUIT.

Fruit medium or above. *Form* roundish oblate to roundish ovate, somewhat irregular; sides unequal. *Stem* short to medium, rather slender. *Cavity* acute or approaching acuminate, medium in width, moderately deep to deep, often somewhat russeted. *Calyx* medium in size, usually closed; lobes narrow, acuminate. *Basin* wide, moderately deep to shallow, smooth or slightly furrowed.

Skin rather thin, nearly smooth, yellow, blushed and streaked with red. *Dots* rather small, greenish.

Calyx tube usually short, wide, conical. *Stamens* marginal.

Core medium, abaxile; cells open; core lines slightly clasping or sometimes meeting.

Flesh whitish tinged with yellow, firm, moderately fine, crisp, rather juicy, mild subacid, good.

Season late August and September.

BENONI.

REFERENCES. 1. *N. E. Farmer*, 9:46. 1830. 2. Kenrick, 1832:25. 3. *Mag. Hort.*, 1:149, 363. 1835. 4. Manning, 1838:49. 5. *Mag. Hort.*, 6:172. 1840. 6. *Ib.*, 7:43. 1841. 7. Downing, 1845:70. 8. Hovey, *Mag. Hort.*, 14:17. 1848. fig. 9. Thomas, 1849:136. 10. Cole, 1849:101. 11. Emmons, *Nat. Hist. N. Y.*, 3:12. 1851. 12. Hovey, 1:83. 1851. col. pl. and fig. 13. Barry, 1851:279. 14. *Am. Pom. Soc. Cat.*, 1854. 15. Elliott, 1854:122. 16. *Mag. Hort.*, 21:63. 1855. 17. Gregg, 1857:35. 18. Hooper, 1857:17, 106, 108. 19. *Horticulturist*, 14:425. 1859. 20. Warder, 1867:650. fig. 21. Fitz, 1872:121, 148, 177. 22. Hogg, 1884:20. 23. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 24. Bailey, *Am. Hort.*, 1892:235. 25. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. 26. Burrill and McCluer, *Ill. Sta. Bul.*, 45:314. 1896. 27. Dickens and Greene, *Kan. Sta. Bul.*, 106:51. 1902. 28. Budd-Hansen, 1903:47. fig.

SYNONYMS. None.

Benoni is a fine dessert apple, very attractive in appearance and excellent in quality but not large enough to be a good market variety. The tree comes into bearing moderately young and



GENONI

yields fair to good crops biennially. It begins to ripen early in August and its season extends into September.

Historical. Originated in Dedham, Massachusetts, where the original tree was still standing in 1848. It was introduced to notice by Mr. E. M. Richards shortly before 1832 (2). It is highly esteemed throughout the country and is generally listed by nurserymen throughout the middle and northern portions of the apple-growing regions of this continent (24).

TREE.

Tree rather large, vigorous. *Form* erect to somewhat roundish, dense. *Twigs* moderately long, straight, slender; internodes medium. *Bark* olive-green, shaded with light reddish-brown, lightly coated with scarf-skin, pubescent. *Lenticels* scattering, medium, oblong, slightly raised. *Buds* deeply set in bark, medium size, plump, obtuse, appressed, slightly pubescent.

FRUIT.

Fruit medium to rather small. *Form* roundish inclined to conic, faintly ribbed toward the apex; sides unequal. *Stem* short to very short, slender. *Cavity* acute, rather narrow, moderately deep, wavy, greenish-russet. *Calyx* rather small to above medium, partly open, slightly pubescent. *Basin* medium in width and depth, abrupt, somewhat wrinkled.

Skin smooth, orange-yellow partly covered with lively red striped with deep carmine. *Dots* scattering, minute, whitish.

Stamens basal.

Core small to medium, axile; cells closed; core lines meeting. *Carpels* roundish, slightly elongated, emarginate. *Seeds* few, dark brown, medium in size, plump, obtuse.

Flesh yellow, firm, crisp, fine-grained, tender, juicy, pleasant subacid, good to very good.

Season August and early September.

BIETIGHEIMER.

REFERENCES. 1. Downing, 1881:101 app. 2. Thomas, 1885:521. 3. Hoskins, *Rural N. Y.*, 47:646. 1888. 4. Wickson, 1889:243. 5. *Can. Hort.*, 13:239, 301. 1890. fig. 6. Bailey, *An. Hort.*, 1892:234. 7. *Can. Hort.*, 16:14. 1893. 8. *Ib.*, 17:413. 1894. col., pl. 9. *Gard. and For.*, 8:390, 428. 1895. 10. *Am. Pom. Soc. Cat.*, 1897:12. 11. Dickens and Greene, *Kan. Sta. Bul.*, 106:54. 1902. 12. Budd-Hansen, 1903:48. fig. 13. Farrand, *Mich. Sta. Bul.*, 205:43. 1903.

SYNONYMS. BEITIGHEIMER (6). BIETIGHEIMER (7, 10, 12, 13). RED BEITIGHEIMER (9). *Red Beitigheimer* (6). RED BIETIGHEIMER (1, 2, 3, 4, 5, 8, 11).

Fruit remarkable only for its great size and beauty. It is a good cooking apple but coarse, subacid and not desirable for dessert use. The fruit being extremely large, drops badly before

the crop is ready to pick. In the nursery it is a rough grower forming many badly shaped trees and for this reason it is best to topwork it on some good straight stock. The tree comes into bearing rather early and under favorable conditions is an annual cropper but only moderately productive. It is a fine fruit for exhibition but is not worthy of cultivation for either home use or market.

Historical. Origin, Germany.

TREE.

Tree large, moderately vigorous to vigorous. *Form* upright spreading or roundish, dense, with laterals inclined to droop. *Twigs* short, curved, stout, with large terminal buds; internodes long. *Bark* dull brown tinged with green, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, conspicuous, medium in size, oval, raised. *Buds* prominent, large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit very large, pretty uniform in size and shape. *Form* roundish oblate or inclined to conic, with broad, flat base, somewhat irregular. *Stem* medium to short, thick. *Cavity* large, acute, or approaching acuminate, wide, moderately shallow to rather deep, sometimes furrowed, occasionally lipped, often much russeted and with outspreading russet rays. *Calyx* medium to small, closed; lobes rather narrow, acute. *Basin* varies from shallow and obtuse to deep and abrupt, medium in width, somewhat wrinkled, often marked with mammiiform protuberances.

Skin thick, tough, smooth, bright pale yellow to greenish or whitish washed with pinkish-red and sparingly and obscurely splashed with deeper red. *Dots* numerous, small, inconspicuous, yellowish or russet.

Calyx tube broadly conical. *Stamens* usually basal or nearly so.

Core medium to large, axile to somewhat abaxile; cells partly open; core lines clasping. *Carpels* cordate or broadly roundish, a little tufted. *Seeds* numerous, large to medium, rather wide, broadly acute, rather light brown.

Flesh almost white, firm, very coarse, crisp, somewhat tough, juicy, sub-acid, fair to nearly good.

Season September and October.

BIRTH.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 8:42, 73. 1881-82. 2. Budd, *Ia. Hort. Soc. Rpt.*, 1882:79. 3. *Ib.*, 1883:444. 685. fig. 4. *Montreal Hort. Soc. Rpt.*, 1883:107. fig. 5. *N. Y. Sta. An. Rpt.*, 2:35. 1883. 6. *Ia. Agr. Coll. Bul.*, 1885:18. 7. *Montreal Hort. Soc. Rpt.*, 1886-87:80. 8. *N. Y. Sta. An. Rpt.*, 11:588. 1892. 9. *Ia. Sta. Bul.*, 41:70. 1899.

SYNONYMS. BIRTH (8, 9). CHRIST BIRTH (6). CHRIST BIRTH APPLE (2, 3). Christ Birth Apple (4). CHRISTMAS (7). No. 161 (7). No. 477



PICTURE

(2, 6, 9). 161 *M* (2). RESCHESTWENSKOE (5). *Roschdestwenskoe* (6). ROSCHDESTWENSKOE (4). *Roschdestwenskoe* (3). ROSHDESTRENSKOE (1).

A Russian apple received in 1888 from Dr. T. H. Hoskins, Newport, Vt., for testing at this Station. Fruit above medium, roundish conic, slightly ribbed; skin greenish-yellow with a shade of brownish-red; flesh mild sub-acid, fair quality; ripens here in September. Not valuable.

BISMARCK.

REFERENCES. 1. Hogg, 1884:181. 2. *Rural N. Y.*, 55:275, 288, 321, 690, 1896. 3. Van Deman, *ib.*, 56:241, 503, 534, 598, 662. 1897. *figs.* 4. Green, *Am. Pom. Soc. Rpt.*, 1897:32. 5. *Rural N. Y.*, 57:786. 1898. 6. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:356. 7. *Can. Hort.*, 22:240. 1899. 8. *Amer. Gard.*, 20:124, 782. 1899. *figs.* 9. Van Deman, *Rural N. Y.*, 58:688. 1899. 10. Beach, *W. N. Y. Hort. Soc. Rpt.*, 1900:35. 11. Waugh, *Vt. Sta. An. Rpt.*, 14:288. 1901. 12. *Can. Hort.*, 25:47. 1902. 13. *Rural N. Y.*, 61:626. 1902. 14. Budd-Hansen, 1903:49. 15. *Rural N. Y.*, 62:809. 1903.

SYNONYMS. BISMARCK (2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15). BISMARCK (10). PRINCE BISMARCK (1). *Prince Bismark* (10).

Bismarck is evidently related to the Apport group of apples. In size and general appearance it somewhat resembles Alexander. Fruit large, attractive in color, suitable for kitchen and market purposes but inferior in dessert qualities. It ranks about with Alexander and Wolf River in quality. It begins to ripen in October and its season extends from October to early winter. It has not been tested enough in this country to demonstrate its market value. The tree is dwarfish, healthy, hardy, comes into bearing very young, is a reliable cropper and very productive. Even when grown as standards the trees may be planted much more closely together than ordinary commercial varieties.

Historical. Originated in the Province of Canterbury, New Zealand. Introduced into this country from England about ten years ago.

TREE.

Tree dwarfish with very short, stout, drooping branches. *Form* spreading, open. *Twigs* short, curved, moderately stout; internodes medium. *Bark* dull brownish, tinged with green, lightly coated with scarf-skin, pubescent. *Lenticels* scattering, medium to large, oval, slightly raised. *Buds* medium in size, plump, obtuse, free, pubescent.

FRUIT.

Fruit very large or large, rather uniform in size and shape. *Form* roundish oblate to roundish conic, flattened at the base, pretty regular; sides often

unequal. *Stem* short to medium, thick. *Cavity* usually rather large, acuminate, moderately wide to wide, deep, often compressed, greenish or russet with outspreading russet rays. *Calyx* large, open; lobes short, rather broad, nearly obtuse. *Basin* large to very large, usually symmetrical, deep, moderately wide to wide, very abrupt, sometimes broadly and irregularly furrowed and wrinkled.

Skin rather thick, tough, smooth, greenish or yellow washed, mottled and striped with two shades of red becoming solid dark red on the exposed cheek, overspread with thin bloom and often marked with thin scarf-skin about the base. *Dots* minute and russet or large and pale gray. *Prevailing effect* attractive red with less of a striped appearance than Alexander.

Calyx tube wide, broadly conical to somewhat funnel-form. *Stamens* basal.

Core medium to rather small, axile to somewhat abaxile; cells closed or sometimes open; core lines meeting or slightly clasping. *Carpels* flat, broadly ovate to nearly cordate, tufted. *Seeds* few, often abortive, medium size, rather wide, short, plump, obtuse to acute, medium brown.

Flesh nearly white, moderately firm, coarse, rather tender, juicy, subacid, sprightly, fair to good or nearly good.

Season October to early winter.

BLACK ANNETTE.


REFERENCES. 1. Elliott, 1854:167. 2. ? Warder, 1867:713. 3. Downing, 1869:99.

SYNONYMS. None.

A rather small dark red apple formerly grown to a limited extent in some sections of New York and other Eastern states. Season November and December. It is now practically obsolete here. The Black Annette mentioned by Hansen when grown in Central Iowa keeps through the winter which indicates that it is distinct from the variety here described. See Vol. I.

BLENHEIM.

REFERENCES. 1. Turner, *London Hort. Soc. Trans.*, 3:322. 1819. 2. Forsyth, 1824:134. 3. *Pom. Mag.*, 1:No. 28. 1828. col. pl. 4. Ronalds, 1831:61. fig. 5. *London Hort. Soc. Cat.*, 1831:No. 104. 6. Kenrick, 1832:72. 7. Floy-Lindley, 1833:29. 8. Downing, 1845:81. 9. *Horticulturist*, 1:389. 1847. 10. Thomas, 1849:144. 11. Emmons, *Nat. Hist. N. Y.*, 3:29. 1851. 12. Elliott, 1854:167. 13. Hooper, 1857:19. 14. Lucas, Ed., *Ill. Handb. Obstk.*, 1:515. 1859. 15. Warder, 1867:713. 16. Regel, 1:461. 1868. 17. Berghuis, 1868:No. 61. col. pl. 18. *Tilt. Jour. Hort.*, 7:166. 1870. 19. Downing, 1872:3 app. 20. Leroy, 1873:139. fig. 21. *Am. Pom. Soc. Cat.*, 1875:6. 22. Lauche, 1882:No. 36. col. pl. 23. Bensen, *Rural N. Y.*, 42:65. 1883. 24. Hogg, 1884:24. 25. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:288. 26. *Can. Hort.*, 15:40, 72, 111, 124, 156, 188, 206. 1892. 27. Bailey, *Am. Hort.*, 1892:235. 28. *Can. Hort.*, 16:113, 299. 1893. 29. Bredsted, 2:157. 1893. 30. *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. 31. Gaucher, 1894:No. 8. col. pl. 32. *Gard. and For.*, 9:15. 1896. 33. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 34.



BISMARCK (Reduced Size)

Amer. Gard., 19:344. 1898. 35. *Can. Hort.*, 24:353. 1901. figs. 36. *Ont. Fr. Stas. An. Rpt.*, 8:6. 1901. figs. 37. Eneroth-Smirnoff, 1901:197. 38. Macoun, *Can. Dept. Agr. Bul.*, 37:40. 1901. 39. Budd-Hansen, 1903:51. 40. Beach and Clark, *N. Y. Sta. Bul.*, 248:113. 1904.

SYNONYMS. *Belle d'Angers* (31). *BLenheim* (20, 23, 34, 35, 36, 39, 40). *Blenheim* (5, 18, 19). *BLenheim ORANGE* (1, 2, 4, 6, 13, 15, 26, 28, 32, 33). *Blenheim Orange* (3, 5, 7, 8, 10, 11, 12, 18, 19, 20, 24, 31, 34, 35, 36, 39, 40). *BLenheim PIPPIN* (3, 5, 7, 8, 9, 10, 11, 12, 16, 17, 18, 19, 21, 24, 25, 27, 29, 30, 38). *Blenheim Pippin* (6, 20, 26, 28, 31, 34, 35, 36, 39, 40). *BLenheim REINETTE* (37). *Blooming Orange* (18, 19, 31). *Dredge's Fane* (31). *Dutch Mignonette*, etc. (18, 19). *Gloucester Pippin* (31). *GOLDREINETTE VON BLenheim* (14, 22, 31). *Goldreinette von Blenheim* (20). *Impératrice Eugénie*. (31). *Kempster's Pippin* (18, 19, 20, 24, 31, 32). *LORD NELSON* (incorrectly in some nursery catalogues). *Lucius Apfel* (31). *Northampton* (31). *Northwick Pippin* (5, 18, 19, 20, 24, 31). *Orange Blenheim* (31). *Orange Pippin* (31). *Perle d'Angleterre* (31). *Prince of Wales* (31). *Pomme de Blenheim* (31). *Reinette de Blenheim* (31). *Ward's Pippin* (31). *Woodstock* (2, 35, 36). *Woodstock Pippin* (3, 5, 6, 7, 8, 10, 11, 12, 18, 19, 20, 24, 31, 34).

Fruit large to very large, yellow, more or less washed and striped with red, attractive in appearance and of excellent quality. The commercial season in the southeastern portions of the State is October. In Western New York it comes into season with the Twenty Ounce and keeps into early winter (31). Often specimens of it may be kept much later. Macoun gives its season in Ontario as November and December (28). The fruit is desirable both for home and market uses but the variety is usually unsatisfactory for commercial planting because it is not a good keeper, is variable in season and commonly suffers considerable loss in drops and culls. In some locations, however, it is considered a good profitable variety.

Origin Woodstock, Oxfordshire, England. It found its way into the London nurseries about the year 1818 (24). Although it has long been known in portions of New York and adjacent states and in Canada in no part of this region has it assumed very great commercial importance.

TREE.

Tree large, vigorous, productive, bearing its fruit singly and evenly distributed. *Form* upright spreading. *Twigs* very stout. *Bark* clear, light reddish-brown becoming dark.

FRUIT.

Fruit usually large or above medium. *Form* roundish oblate to roundish inclined to conic, usually pretty regular and symmetrical, sometimes a little

furrowed at the apex. *Stem* short to medium. *Cavity* below medium to rather large, rather narrow to wide, moderately deep to deep, acute to acuminate, usually symmetrical, sometimes compressed or lipped, covered with russet which often extends beyond the cavity. *Calyx* large or very large; segments flat, separated at base plainly exposing the yellowish tube beneath; lobes obtuse. *Basin* large, broad, shallow and obtuse to deep and abrupt, somewhat furrowed and slightly wrinkled.

Skin moderately thin and tough, deep yellow overspread with a rather dull pinkish-red, in highly colored specimens developing a deep and rather bright red somewhat roughened in places with netted capillary russet lines. *Dots* numerous, small or conspicuously large and russet. *Prevailing effect* rather attractive red and yellow.

Calyx tube short, very wide, cone-shape. *Stamens* median to somewhat basal.

Core medium or below, axile or somewhat abaxile; cells often unequally developed, closed or partly open; core lines meeting. *Carpels* flat, tufted, emarginate. *Seeds* few and frequently abortive, irregular, often not plump, long, acute to acuminate, tufted.

Flesh tinged with yellow, rather firm, moderately juicy, crisp, moderately fine grained or a little coarse, somewhat aromatic, agreeable sprightly subacid, becoming rather mild subacid, good to very good; excellent either for dessert or culinary use.

Season. It is at its best from October to December but often may be kept until midwinter or later.

BLUSHED CALVILLE.

REFERENCES. 1. Budd, *Ia. Hort. Soc. Rpt.*, 1882:80. 2. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:72. 3. Budd, *Can. Hort.*, 11:223, 246. 1888. 4. *Ib.*, *Rural N. Y.*, 47:692. 1888. 5. *Ib.*, *Ia. Agr. Coll. Bull.*, 1890:17. 6. *Ib.*, *Can. Hort.*, 13:216. 1890. 7. *Ib.*, *Ia. Sta. Bul.*, 19:535. 1892. 8. Green, *Minn. Sta. Bul.*, 32:241. 1893. 9. Munson, *Me. Sta. Rpt.* 1896:73. 10. Thomas, 1897:265. fig. 11. Hansen, *S. D. Sta. Bul.*, 76:32. 1902. fig. 12. Budd-Hansen, 1903:52.

SYNONYMS. BLUSHED CALVILLE (2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12). CALVILLE KRASNCI (1). 22 (2). 22 M (1, 4, 5, 7, 8, 11, 12).

Blushed Calville is said to be hardy and desirable in northern apple-growing regions (11). As fruited at this Station the tree does not come into bearing very young and is not very productive. It is not recommended for planting in this state.

Historical. Origin Russia.

TREE.

Tree rather small, moderately vigorous with short, stout branches. *Form* upright spreading, open. *Twigs* medium in length, curved and stout with large terminal buds; internodes long. *Bark* brownish mingled with olive-green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, medium in size, round, slightly raised. *Buds* prominent, large, broad, plump, acute, free, slightly pubescent.



BLENHEIM

FRUIT.

Fruit medium to large, fairly uniform in shape and size. *Form* roundish conical, ribbed; sides unequal. *Stem* usually long and slender. *Cavity* acute to acuminate, rather narrow to moderately wide, moderately deep, sometimes russeted. *Calyx* large, closed or open. *Basin* medium in width and depth to wide and deep, a little abrupt, wrinkled.

Skin light green or yellowish, sometimes blushed.

Calyx tube broad, cone-shaped. *Stamens* median.

Core very large, abaxile; cells wide open; core lines clasping. *Seeds* medium in size, acute.

Flesh whitish, firm, rather coarse, crisp, tender, juicy, subacid, fair to good.

Season early summer.

BONUM.

REFERENCES. 1. Robey, *Horticulturist*, 11:89. 1856. 2. Downing, 1857: 122. 3. *Am. Pom. Soc. Cat.*, 1860:240. 4. Warder, 1867:424. fig. 5. Leroy, 1873:147. fig. 6. Thomas, 1875:190. 7. Barry, 1883:343. 8. Bailey, *Am. Hort.*, 1892:235. 9. Dempsey, *Ont. Fr. Stas. An. Rpt.*, 2:32. 1895. 10. Barrill and McCluer, *Ill. Sta. Bul.*, 45:315. 1896. 11. Alwood, *Va. Sta. Bul.*, 130:123. 1901. 12. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:38. 1903. 13. Budd-Hansen, 1903:54. fig.

SYNONYMS. BONUM (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13). *Magnum Bonum* (2, 4, 5, 7).

This is a southern variety of very good quality when grown under favorable conditions. It is in season during late fall and early winter. Probably it is not well adapted to regions as far north as New York, for although it has long been cultivated it is practically unknown among New York fruit growers.

Historical. Origin Davidson county, N. C. It was entered on the catalogue of the American Pomological Society in 1860, dropped from the list in 1862 and reentered in 1869. According to Bailey's Inventory of North American Apples (8) it is now propagated but little by nurserymen.

TREE.

Tree moderately vigorous. *Form* upright spreading or roundish, open. *Twigs* moderately long, curved, moderately stout; internodes medium. *Bark* dull brown, lightly mottled with scarf-skin; slightly pubescent. *Lenticels* quite numerous, small, round, not raised. *Buds* medium in size, flat, acute, free, not pubescent.

FRUIT (2, 4, 6, 7, 13).

Fruit medium to large. *Form* oblate, regular. *Stem* long, slender to moderately thick, green. *Cavity* medium to large, deep, regular, often with a little green russet. *Calyx* large, closed. *Basin* medium in width, shallow, wrinkled.

Skin smooth, yellow, mostly covered with crimson and dark red, striped. *Dots* distinct, large, light with some having a dark center.

Calyx tube funnel-form. *Stamens* marginal.

Core small; cells closed; core lines scarcely meeting. *Carpels* ovate. *Seeds* numerous, large, plump.

Flesh white, often stained next to the skin, firm, fine, tender, juicy, aromatic, mild subacid, very good for dessert.

Season September to November.

BOROVINKA.

REFERENCES. 1. *Montreal Hort. Soc. Rept.*, 8:37. 1881-82. 2. *Ib.*, 8:75. 1881-82. 3. Budd, *Ia. Hort. Soc. Rpt.*, 1882:80. 4. Gibb, *Ib.*, 1883:432. 5. *Montreal Hort. Soc. Rpt.*, 9:80. 1883. fig. 6. Budd, *Ia. Agr. Coll. Bul.*, 1885:9. 7. ? Schroeder, *Montreal Hort. Soc. Rpt.* 1886-87:71. 8. *Ib.*, 1886-87:79. 9. Van Deman, *U. S. Pom. Rpt.*, 1888:571. 10. Budd, *Ia. Agr. Coll. Bul.*, 1890:18. 11. *Can. Hort.*, 13:216. 1890. 12. Budd, *Ia. Sta. Bul.*, 19:536. 1892. 13. Harris, *U. S. Pom. Rpt.*, 1892:273, 278. 14. Taylor, *Me. Pom. Soc. Rpt.*, 1892:58. 15. Niemetz, *Can. Hort.*, 16:113. 1893. 16. Green, *Minn. Sta. Bul.*, 32:240. 1893. 17. Stinson, *Ark. Sta. Bul.*, 43:105. 1896. 18. Thomas, 1897:629. 19. *Am. Pom. Soc. Cat.*, 1897:12. 20. Hansen, *S. D. Sta. Bul.*, 76:33. 1902. 21. Farrand, *Mich. Sta. Bul.*, 205:43. 1903. 22. Budd-Hansen, 1903:54. fig.

SYNONYMS. BOROVINKA (1, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22). *Borovinka* (4). BOROVITSKY (18). BOROVINKA ANGL'SKAIA (2, 3). (ENGLISH BOROVINKA, 7)? MUSHROOM (4). *Mushroom* (5). *g M* (2, 3). *No. 245* (6, 9, 10, 12, 17).

Borovinka resembles Oldenburg so closely that Hansen says the question of their identity has not been settled (20). As fruited at this Station it is distinct from Oldenburg; it is fully as attractive as Oldenburg in color but it lacks uniformity in size and is not equal to that variety in flavor and quality. The stock grown at this Station came from Professor J. L. Budd, Ames, Ia., in 1890, and is doubtless the true Borovinka.

Historical. Origin Russia.

TREE.

Tree below medium size but moderately vigorous. *Form* upright spreading to rather flat, open. *Twigs* short, curved, stout; internodes short. *Bark* dark brown, lightly mottled with scarf-skin; slightly pubescent. *Lenticels* scattering, medium to large, oblong, slightly raised. *Buds* prominent, medium in size, broad, plump, obtuse to acute, free, not pubescent.

FRUIT.

Fruit below medium to large, averaging medium; pretty uniform in shape but not in size. *Form* roundish, slightly flattened at the ends, regular or faintly ribbed. *Stem* medium in length, thick. *Cavity* acute, rather shallow to moderately deep, moderately broad, slightly furrowed, sometimes with faint radiating rays of russet. *Calyx* medium to rather large, closed; lobes broad. *Basin* medium to rather deep, wide, somewhat abrupt, slightly furrowed, occasionally showing mammiform protuberances.

Skin thin, very tender, smooth, pale yellow, often entirely covered with broken stripes and irregular splashes of attractive bright red, overspread with thin bluish bloom. *Dots* numerous, conspicuous, very small, light colored.

Calyx tube large, rather wide, urn-shape to funnel-form widening in the lower part of the funnel cylinder. *Stamens* median to marginal.

Core small to medium, axile; cells closed or nearly so; core lines clasping. *Carpels* roundish, somewhat concave, mucronate, not emarginate. *Seeds* medium to rather large, moderately wide, plump, somewhat obtuse, dark brown.

Flesh tinged with yellow, medium in grain, crisp, tender, moderately juicy to juicy, agreeable subacid, slightly aromatic, good.

Season mid-August to mid-September.

BOSKOOP.

REFERENCES. 1. Berghuis, 1868: col. pl. No. 43. 2. Oberdieck, *Monatshefte*, 1869:193. 3. Oberdieck, *Deutschlands beste Obstsorten*, 212. 4. Lauche, *Ergänzungsband zum Ill. Handb.*, 265. 5. *Montreal Hort. Soc. Rpt.*, 7:56, 155. 1881. 6. Downing, 1881:77 app. fig. 7. Budd, *Ia. Hort. Soc.*, 1882: 8. Barry, 1883:342. 9. Willard, *W. N. Y. Hort. Soc.*, 1889:171 (reprint in *Mich. Hort. Soc. Rpt.*, 1889:329). 10. Palandt, *Gartenflora*, 38:425. 1889. col. pl. 11. Brodie, *Can. Hort.*, 12:238. 1889. 12. Bailey, *Am. Hort.*, 1892:234. 13. Bailey, *Amer. Gard.*, 14:501. 1893. 14. Craig, *Can. Hort.*, 16:138. 1893. fig. 15. Bredsted, 1893:233. 16. Gaucher, *Pomologie*, 1894:No. 20. col. pl. 17. Beach and Clark, *N. Y. Sta. Bul.*, 248:113. 1904.

SYNONYMS. BELLE DE BOSKOOP (11). BELLE OF BOSKOOP (9). BELLE DE BOSKOOP (6, 7, 8, 12, 13, 14, 15). *Belle de Boskoop* (16, 17). BOSKOOP (5, 17). *Reinette Belle de Boskoop* (16). *Reinette Monstrueuse* (16). *Reinette von Montfort* (16). SCHÖNER VON BOSKOOP (2, 3, 4, 10, 16). SCHOONE VON BOSKOOP (1). *Schoone van Boskoop* (16).

In some locations this fruit becomes highly colored with attractive bright red predominating, but more often the color is not good, being predominantly dull green or yellow and more or less russeted. It is more suitable for general market and culinary purposes than for dessert. It is of good size but does not rank high in quality; the texture is somewhat coarse, and the flavor rather too acid for an agreeable dessert apple, but late in the season its acidity becomes modified. It appears to be pretty hardy and a good bearer. When grown on warm soils in Southern New York it may be marketed in September, but in the more northern regions of the state it keeps into the winter. It is perhaps of sufficient merit to be worthy of testing but we are not yet ready to recommend it for general planting.

Historical. This variety is said to have originated from seed in 1856 in the nursery of the Ottolander family at Boskoop (1, 4). Palandt finds that it is identical with the variety described by Lauche and Oberdieck as "*Reinette von Montfort*" (4). It was imported into this country more than twenty-five years ago (5) and has gradually been disseminated to a limited extent in various portions of New York state.

TREE.

Tree rather large, moderately vigorous; branches long, moderately stout, crooked; lateral branches numerous and small. *Form* open, wide-spreading and drooping. *Twigs* rather short to long, straight, rather stout; internodes below medium to very long. *Bark* dark brownish-red, mingled with olive-green; somewhat pubescent. *Lenticels* numerous, conspicuous, small, oblong or roundish. *Buds* rather large, broad, plump, acute, free, slightly pubescent. *Leaves* large, broad.

FRUIT.

Fruit large. *Form* usually oblate, sometimes roundish oblate, obscurely ribbed, sometimes with oblique axis; pretty uniform in size and shape. *Stem* usually short and thick, sometimes rather long. *Cavity* rather large, acute to acuminate, somewhat furrowed, often irregular, deep, russeted. *Calyx* large; segments long or very long, acuminate, closed or somewhat open, sometimes separated at the base. *Basin* abrupt, rather narrow, moderately shallow to rather deep, sometimes slightly furrowed.

Skin dull green or yellowish, sometimes blushed and mottled with rather bright red, and striped with deeper red, roughened with russet flecks, often irregularly overspread with russet. *Dots* small and gray, mingled with others which are large, irregular and russet.

Calyx tube large, cone-shape. *Stamens* median to basal.

Core medium to small, somewhat abaxile; cells often unsymmetrical, closed or open; core lines slightly clasping. *Carpels* roundish or obcordate, a little tufted. *Seeds* apt to be abortive; when well developed they are long, irregular, obtuse to acute, somewhat tufted.

Flesh tinged with yellow, firm, somewhat coarse, tender, juicy, crisp, brisk subacid, good to very good.

Season. Commercial season September to November (17). As grown in Western New York generally some of the fruit may be kept till April.

BOUGH SWEET.

This variety is also known as Bough Apple, Large Yellow Bough, Sweet Bough and Summer Sweet Bough. It is listed in the late catalogues of the American Pomological Society¹ as Bough, *Sweet* but most nurserymen list it as Sweet Bough.² We prefer to recognize the name commonly accepted by nurserymen and accordingly have described the variety under the name Sweet Bough. See page 216.

BRESKOVKA.

REFERENCES. 1. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:80. 2. Budd, *Can. Hort.*, 11:246. 1888. 3. *Ib.*, *Rural N. Y.*, 47:692. 1888. 4. *Ib.*, *la. Agr. Col. Bul.* 1890:17. 5. *Ib.*, *la. Sta. Bul.*, 19:535. 1892. 6. Green, *Minn.*

¹ *Am. Pom. Soc. Cat.*, 1897:12.

² Bailey, *Am. Hort.* 1892:235, 250.

Sta. Bul., 32:240. 1893. 7. Thomas, 1897:629. 8. Hansen, *S. D. Sta. Bul.*, 76:34. 1902. 9. Budd-Hansen, 1903:56.

SYNONYMS. BRESKOVKA (1, 2, 3, 4, 5, 6, 7, 8, 9). 152 M (2, 3, 4, 5, 8, 9).

A hardy Russian variety of Yellow Transparent type, in season during late August and early September. The flesh quickly discolors as the ripening season advances. It is rather attractive in color for a yellow apple but does not average above medium size and it is not equal to Yellow Transparent in either flavor or quality. Not recommended for growing in New York.

BUNKER HILL.

REFERENCE. 1. Downing, 1872:4 app. fig.

SYNONYMS. None.

This variety has been planted to some extent in Central New York and is regarded by some fruit growers in that region as a profitable commercial sort. The tree is large, upright spreading, vigorous to moderately vigorous with long, spreading, moderately stout twigs. It is hardy, healthy, medium to long-lived and a reliable cropper, usually bearing heavy crops biennially. The fruit is subacid and good either for dessert or culinary uses. It is in season from mid-autumn to early winter.

Historical. Originated in the orchard of Dr. Paige, Dryden, Tompkins county, N. Y. (1).

FRUIT (1).

"Fruit medium, roundish conical, regular; skin pale whitish-yellow shaded, mottled, striped and splashed with two shades of red, rather thinly over two-thirds of the surface, and moderately sprinkled with light dots, a few being areole; stalk short, slender; cavity medium or large, a little greenish; calyx closed; basin medium, slightly corrugated; flesh quite white, sometimes a little stained next the skin, fine, tender, juicy, subacid, vinous, slight quince-like flavor; very good; core rather small."

BUTTER.

REFERENCES. 1. Elliott, 1854:125, 159, 174. 2. Downing, 1857:125. 3. Warder, 1867:392. 4. Downing, 1869:112. 5. Fitz, 1872:152. 6. Thomas, 1875:495. 7. Ragan, *U. S. B. P. I. Bul.*, 56:60. 1905.

SYNONYMS. None.

Downing describes a variety under this name which is above medium size, yellow, with whitish flesh, very sweet and rich, valuable for cooking and esteemed for making apple butter; season September and October (2, 4). Other varieties have been known under the name Butter which, as Downing remarks, "appears to be a favorite name with some to apply to any good sweet apple for sauce or cooking."

The references above cited do not all refer to the same variety.

CABASHEA.

REFERENCES. 1. *N. Y. Agr. Soc. Trans.*, 1849:350. 2. Emmons, *Nat. Hist. N. Y.*, 3:103. 1851. 3. Warder, 1867:714. 4. Thomas, 1875:495. 5. Beach, *Apples of New York*, 1:91. 1905.

SYNONYMS. CABASHEA (1, 3, 4, 5). *Cabashea* (2). CABASHIE (2).

The name Cabashea has been applied by many pomologists to the variety commonly known among fruit growers and fruit dealers as Twenty Ounce Pippin (5), an apple which comes in season about with Tompkins King. The variety which is generally known in Western New York as Cabashea comes in season about with the true Twenty Ounce but it is not so good a keeper. In 1851 Emmons published a cut of a section of this Cabashea showing well its characteristically oblate form. Emmons remarked, "This apple is more remarkable for its size than for its valuable qualities. . . . It is not, however, an inferior apple. For cooking it is certainly esteemed, as it has a pleasant and agreeable taste. It is, however, too large." The tree is hardy, healthy, medium to long-lived, and a pretty regular cropper, yielding moderate to rather light crops nearly annually. It is not considered a good commercial variety because it is not sufficiently productive and the fruit does not sell very well.


Historical. This variety appears to be a Western New York seedling (1). It is now seldom or never planted.

TREE.

Tree medium size, moderately vigorous. *Form* erect or somewhat spreading. *Twigs* medium to long, curved, spreading, stout to rather slender; internodes medium. *Bark* reddish-brown tinged with olive-green, streaked with scarf-skin, heavily pubescent near tips. *Lenticels* conspicuous, scattering, large, oval, raised. *Buds* large, broad, obtuse, free, pubescent; tips stout.

FRUIT.

Fruit large to very large, fairly uniform in size and shape. *Form* roundish oblate to decidedly flat, obscurely ribbed; sides somewhat unequal. *Stem* usually short, moderately slender. *Cavity* acute, deep, very broad, often somewhat furrowed, much russeted with greenish russet often extending beyond the cavity. *Calyx* large or sometimes medium, usually closed; lobes long, medium in width, acute. *Basin* large, deep, wide, somewhat furrowed, unsymmetrical.



CABASHEA

Skin moderately tender, smooth, slightly unctuous, yellowish-green mottled and blushed with yellowish-red, with broad stripes and splashes of brighter and deeper red. *Dots* small, inconspicuous, often submerged. *Prevailing color* in many specimens yellowish-green with broad stripes of faint red; in more highly-colored specimens the red striping becomes quite distinct.

Calyx tube large, wide, conical. *Stamens* basal.

Core large, decidedly abaxile; cells wide open; core lines meeting. *Carbels* elongated ovate, distinctly concave, slightly tufted. *Seeds* few, medium size, irregular, plump, obtuse, dark.

Flesh greenish or tinged somewhat with yellow, rather firm, coarse, crisp, juicy, subacid or quite acid, fair for dessert, good for cooking.

Season September and October.

CATHEAD.

REFERENCES. 1. Forsyth, 1803:56. 2. Cox, 1817:133. fig. 3. Thacher, 1822:122. 4. Floy-Lindley, 1833:48. 5. Downing, 1845:103. 6. Thomas, 1849:179. 7. Emmons, *Nat. Hist. N. Y.*, 3:86. 1851. 8. Elliott, 1854:168. 9. Warder, 1867:715. 10. Leroy, 1873:842. fig. 11. Hogg, 1884:41.

SYNONYMS. CATHEAD (2, 3, 8). *Cathead Greening* (5, 6, 8). CATSHEAD (1, 4, 5, 6, 7, 9, 11). *Catshead* (4, 10). *Catshead Greening* (10). *Costard* (4). *Costard Ray* (7). *Coustard* (4). *De Seigncur d'Automne* (10). *Grosse-Schafnasé* (10). *Round Catshead* (5, 8, 10). *Schafnasé* (10). *Tete d'Ange* (10). TETE DE CHAT (10).

Formerly grown in some of the home orchards of the state but now practically obsolete. Fruit very large, pale green, subacid. Used for cooking and evaporating. An old English variety. Ray described it as long ago as 1688 (4).

CELESTIA.

REFERENCES. 1. Warder, 1867:530. fig. 2. Downing, 1869:122. 3. Fitz, 1872:173. 4. Bailey, *An. Hort.*, 1892:236. 5. Dickens and Greene, *Kan. Sta. Bul.*, 106:52. 1902. 6. Beach and Clark, *N. Y. Sta. Bul.*, 248:114. 1904.

SYNONYMS. None.

Fruit not particularly attractive in color and as tested at this Station not superior to ordinary varieties in quality. Warder says (1) that it is essentially an amateur's fruit of very best quality but its texture and color disqualify it for market. The tree is a moderate grower and not very productive. Not recommended for cultivation in New York.

Historical. Originated from seed of Stillwater Sweet by L. S. Mote, Miami county, Ohio (1).

TREE.

Tree moderately vigorous with short, stout, curved branches. *Form* upright spreading, roundish, rather dense. *Twigs* short to moderately long, slightly curved, moderately slender; internodes medium to long. *Bark* clear brownish-red with some olive-green, lightly mottled with scarf-skin; pubescent. *Lenticels* numerous, small to medium, elongated or roundish,

slightly raised. *Buds* medium size, plump, obtuse to somewhat acute, free, slightly pubescent.

FRUIT.

Fruit medium to large, usually above medium, uniform in size and shape. *Form* roundish inclined to conic, somewhat flattened at the base, markedly ribbed, irregular, somewhat angular. *Stem* medium to long, thick. *Cavity* obtuse to somewhat acute, moderately deep to deep, rather broad, somewhat furrowed, usually russeted. *Calyx* medium in size, usually closed; lobes medium in length, rather narrow, acute. *Basin* shallow, narrow, rather abrupt, much furrowed and wrinkled.

Skin thick, smooth, rather tender, yellow marbled with pale green, and occasionally having a thin brownish blush. *Dots* numerous, small, inconspicuous, submerged, light or russet.

Calyx tube very long to medium, deep, funnel-shape. *Stamens* median to somewhat marginal.

Core large, very abaxile to sometimes axile; cells open or closed; core lines clasping the funnel cylinder. *Carpels* elliptic to broadly obcordate, much concave, emarginate, much tufted. *Seeds* large, wide, long, obtuse, dark dull brown.

Flesh very strongly tinged with yellow, rather firm, moderately coarse, crisp, tender, juicy, pleasant, rather mild, subacid, good.


Season October to January.

CHAMPLAIN.

REFERENCES. 1. *N. E. Farmer*, 1853 (cited by 14). 2. Downing, 1857:128. 3. Warder, 1867:637. 4. Downing, 1869:368. fig. 5. *Am. Pom. Soc. Cat.*, 1871:10. 6. Leroy, 1873:828. fig. 7. Thomas, 1875:197, 496. 8. Barry, 1883:334. 9. Bailey, *An. Hort.*, 1892:245. 10. *Ib.*, 1892:250. 11. *Am. Pom. Soc. Cat.*, 1897:12. 12. Waugh, *Vt. Sta. An. Rpt.*, 14:290. 1901. 13. Budd-Hansen, 1903:61. 14. Ragan, *U. S. B. P. I. Bul.*, 56:300. 1905.

SYNONYMS. *Calkin's Pippin* (4, 14). *CHAMPLAIN* (1, 2, 3, 5, 11, 12, 13, 14). *Champlain* (4, 6, 7, 8, 14). *Geneva Pearmain* (4, 6, 14). *Haverstraw Pippin* (4, 14). *Large Golden Pippin* (4, 6, 14). *Nyack* (9, 13). *NYACK PIPPIN* (9). *Nyack Pippin* (4, 14). *Paper* (3, 4, 6, 14). *Paper-Skin* (3, 14). *Sourbough* (13). *Sour Bough* (4, 6, 7, 14). *SUMMER PIPPIN* (4, 6, 7, 8, 10, 14). *Summer Pippin* (13, 14). *Tart Bough* (4, 6, 14). *Underdunk* (4, 6, 14). *Vermont* (14). *Walworth* (4, 6, 14).

Nurserymen sometimes list this variety as Nyack, and sometimes as Summer Pippin, but seldom or never as Champlain (9, 10). Fruit of good size, smooth and attractive for a greenish-yellow apple. It is good for dessert and excellent for culinary use. Since it ripens in succession from late August till October, more than two pickings are required to secure the crop in good condition, neither too green nor too ripe. The tree is a good grower, hardy,



CHAMPLAIN

healthy, and moderately long-lived. It comes into bearing rather young and is a reliable cropper, yielding good crops biennially or almost annually. Some find Champlain a profitable commercial variety, but usually it is grown for home use rather than for market.

Historical. Origin unknown. In 1871 (5) it was included in the list of the American Pomological Society's Catalogue under the name Summer Pippin, but since 1897 it has been listed as Champlain (11). Old trees of it are frequently found in the home orchards throughout the state. It is now seldom planted.

TREE.

Tree medium to large, vigorous with long and moderately stout branches. *Form* upright spreading to roundish, open. *Twigs* long to medium, straight, moderately stout; internodes long. *Bark* dull brown tinged with olive-green, lightly streaked with scarf-skin, heavily pubescent. *Lenticels* scattering, medium size, oblong, slightly raised. *Buds* medium size, plump, obtuse, appressed, pubescent.

FRUIT.

Fruit medium to large, not very uniform in size or shape. *Form* roundish, rather conical to ovate or somewhat oblong, irregularly ribbed; sides somewhat unequal. *Stem* medium to long, medium to rather thick. *Cavity* acuminate to acute, moderately shallow to deep, rather narrow to medium in width, sometimes furrowed and usually lightly russeted. *Calyx* small to medium, closed or slightly open. *Basin* shallow to medium in depth, narrow, a little abrupt, nearly smooth.

Skin tender, greenish-yellow or pale yellow, often with a light crimson blush. *Dots* numerous, small, russet or submerged.

Calyx tube conical to funnel-form, usually rather short but sometimes elongated. *Stamens* median to marginal.

Core large, axile to somewhat abaxile; cells open; core lines clasp the funnel cylinder. *Carpels* smooth, elongated ovate, not emarginate. *Seeds* rather dark brown, medium size, rather narrow and short, plump, sharp pointed, almost acuminate.

Flesh white or with slight tinge of yellow, rather fine, very tender, juicy, sprightly, subacid, good to very good.

Season late August till October.

CHANDLER.

REFERENCES. 1. Kenrick, 1835:65. 2. Floy-Lindley, 1846:410, app. 3. Thomas, 1849:164. 4. Cole, 1849:122. 5. Emmons, *Nat. Hist. N. Y.*, 3:67. 1851. 6. Elliott, 1854:168. 7. Downing, 1857:128. 8. Hooper, 1857:24. 9. Warder, 1867:715. 10. Downing, 1869:122. 11. Fitz, 1872:169.

SYNONYMS. CHANDLER (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11). *Chandler's Red* (11). *General Chandler* (1). *Late Chandler* (10). *Winter Chandler* (6).

In 1854 Elliott included Chandler in a list of varieties unworthy of cultivation (6). It is a late fall apple, yellowish striped with red. Tree moderately vigorous but a great bearer (7, 10). An old variety supposedly of Connecticut origin though Kenrick (1) ascribes it to Chelmsford, Mass. There may be a confusion of two varieties. It is now but little cultivated.

Waugh describes another Chandler of sweet flavor which seems to be unknown in New York. He states that it is an old variety of Connecticut origin.

FRUIT (5, 6, 7, 10).

Fruit large. *Form* roundish, slightly oblate, irregular, unsymmetrical; sides unequal. *Stem* short.

Skin greenish-yellow, shaded and frequently striped with yellowish-red and with a few streaks of bright red. *Dots* light gray.

Core small. *Seeds* small.

Flesh greenish-yellow, tender, juicy, moderately rich, subacid.

Season mid-autumn to early winter.

CHARLAMOFF.

REFERENCES. 1. (?) Manning, *Tilt. Jour. Hort.*, 6:349. 1869. 2. Budd, *Ia. Hort. Soc. Rpt.*, 1880:525. 3. *Montreal Hort. Soc. Rpt.*, 1881:53, 156. 4. *Ib.*, 1881-82:38. 5. *Ib.*, 1883:83. 6. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:434. 7. *Ia. Agr. Coll. Bul.*, 1885:10. 8. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:77. 9. Bailey, *Am. Hort.*, 1892:236. 10. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57, 58. 11. Butz, *Pa. Sta. Rpt.*, 1895:134. *col. pl.* 12. *Am. Pom. Soc. Cat.*, 1899:16. 13. Budd, *Ia. Sta. Bul.*, 41:66. 1899. 14. Macoun, *Can. Dept. Agr. Bul.*, 37:37. 1901. 15. Hansen, *S. D. Sta. Bul.*, 76:38. 1902. *fig.* 16. Budd-Hansen, 1903:61. *fig.*

SYNONYMS. *Arabka* (14). CHARLAMOFF (2, 3, 6, 8, 9, 10, 12, 13, 15). *Charlamoff* (3). (CHARLOMSKI, 1)? CHARLAMOVSKOE (4). CHARLAMOVSKOE (3, 5). *Charlamowskoe* (6, 7). No. 105 (8). No. 262 (3, 5). *Peter-son's Charlamoff* (16). POINTED PIPKA (11). *Pointed Pipka* (14, 16).

A Russian variety of the Oldenburg type imported for the Iowa Agricultural College by J. L. Budd. Macoun states that it has been grown under several different names in this country, the most common being Pointed Pipka and Arabka (14). Hansen declares that it is entirely distinct from the Charlamoff as grown by J. G. Mitchell and A. G. Tuttle which is a flat apple of upright habit of tree and not as valuable as many more of the same season.

It does very well at Ottawa, Canada, and further north. At its best it is a good dessert apple but it has the fault of remaining in prime condition for only a very short time (14). It ripens a little earlier than Oldenburg but as fruited at this Station is inferior to that variety in quality. It comes into bearing young and is a reliable cropper, yielding fair to heavy crops biennially. It is but little known among New York fruit growers. It may be found of some value in those sections of the state where superior hardiness is a prime requisite.

CHEESEBORO.

REFERENCES. 1. *Mag. Hort.*, 17:263. 1851. 2. Emmons, *Nat. Hist. N. Y.*, 3: col. pl. No. 20. 1851. 3. Elliott, 1854:108. 4. Downing, 1857:211. 5. Hooper, 1857:25. 6. *Am. Pom. Soc. Rpt.*, 1860:243. 7. Warder, 1867:522. 8. Thomas, 1875:496. 9. Waugh, *Vt. Sta. An. Rpt.*, 14:291. 1901. 10. Ragan, *U. S. B. P. I. Bul.* 56:70. 1905.

SYNONYMS. *Canada Reinette* (9). *Cathead* (9). *CHEESEBORO* (10). *CHEESEBORO'S RUSSSET* (6). *CHEESBOROUGH* (7). *CHEESEBOROUGH* (2, 9). *CHEESBOROUGH RUSSSET* (1, 3, 4, 5, 8). *Cheesborough Russet* (9, 10). *Forever Pippin* (10, of some West 3, 5). *Howard Russet* (3, 4, 5, 10). *Kingsbury Russet* (3, 4, 5, 10). *Oxheart* (9). *Pumpkin Sweet* of some (10). *Sweet Russet* (10). *York Russet* (10, of some 3, 5). *York Russetting* (10).

This is an old variety of unknown origin which is fast becoming obsolete. Tree large to very large, very vigorous, long-lived, a reliable cropper yielding good to heavy crops biennially or almost annually; form upright spreading or roundish. Fruit large to very large, conical, dull green overspread with thin russet, coarse, rather dry, subacid or becoming almost sweet, inferior in flavor and quality, suitable for kitchen use only; season October to early winter.

CHENANGO.

REFERENCES. 1. *Horticulturist*, 9:475. 1854. 2. *Am. Pom. Soc. Cat.*, 1869. 3. Downing, 1869:124. fig. 4. Thomas, 1875:199. 5. *Ia. Hort. Soc. Rpt.*, 1879:472. 6. *Montreal Hort. Soc. Rpt.*, 1879:24. 7. Barry, 1883:337. 8. Wickson, 1889:244. 9. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 10. Bailey, *An. Hort.*, 1892:236. 11. *Id.*, 1892:249. 12. *Masson, Me. Sta. Rpt.*, 1893:132. 13. Burrill and McCluer, *Ill. Sta. Bul.*, 45:317. 1896. 14. Waugh, *Vt. Sta. An. Rpt.*, 14:291. 1901. 15. Alwood, *Va. Sta. Bul.*, 130:120. 1901. 16. *W. N. Y. Hort. Soc. Rpt.*, 1901:76. 17. Hansen, *S. D. Sta. Bul.*, 76:39. 1902. 18. *Can. Hort.*, 26:345. 1903. figs. 19. Budd-Hansen, 1903:62. 20. Farrand, *Mich. Sta. Bul.*, 205:44. 1903. 21. Bruner, *N. C. Sta. Bul.*, 182:20. 1903. 22. Beach and Clark, *N. Y. Sta. Bul.*, 248:115. 1904.

SYNONYMS. *Buckley* (3, 4). *CHENANGO* (2, 9, 10, 14, 16, 17, 18, 19, 20, 21, 22). *CHENANGO STRAWBERRY* (1, 3, 4, 5, 6, 7, 8, 13, 15). *Chenango Strawberry* (10, 11, 14, 17, 18, 19, 22). *CHENANGO Strawberry* (12). *Frank* (3, 4). *Jackson* (3, 4). *SHERWOOD'S FAVORITE* (11). *Sherwood's Favorite* (3, 4, 6, 7, 12, 14, 17, 18, 19, 22). *Smyrna* (3). *Strawberry* (1, 3, 4).

Fruit beautiful in appearance, yellowish-white striped with red, of excellent dessert quality and good also for culinary uses. The tree is an early and regular bearer, hardy, healthy, and pretty long-lived. Under favorable conditions it is an annual bearer, alternating rather light with heavy crops. The fruit begins to mature in September and ripens continuously during a period of several weeks. For this reason it should have more than one picking in order to secure the crop in the best condition. The latest ripening

fruit may be kept in ordinary storage till November, but after that the color fades and it deteriorates much in quality, even though it may remain apparently sound (22). The fruit does not ship well because its flesh is too tender. Some find it a profitable variety to grow for local or special markets, but other varieties of its season are more desirable than Chenango for general commercial planting. It is recommended as an excellent variety for the home orchard.

Historical. Chenango, according to some accounts, originated in Lebanon, Madison county, N. Y.; others say that it was early brought into Chenango county by settlers from Connecticut. It has certainly been known in cultivation for more than fifty years (3). It is still propagated by nurserymen but the demand for the stock is quite limited.

TREE.

Tree medium size, vigorous with short, stout, curved branches. *Form* upright spreading to roundish, rather dense. *Twigs* long to medium, curved, moderately slender; internodes medium. *Bark* olive-green tinged with dull brown, lightly streaked with scarf-skin; pubescent. *Lenticels* scattering, small, round, not raised. *Buds* deeply set in bark, small, flat, obtuse, appressed, slightly pubescent.

FRUIT.

Fruit above medium to large, but under unfavorable conditions it may be small and poorly colored. *Form* elongated ovate or oblong conic, slightly ribbed. *Stem* short to medium, moderately thick. *Cavity* acute to acuminate, deep, narrow, often somewhat furrowed and compressed, usually not russeted. *Calyx* medium to large, partly open or closed; lobes often separated at the base, long, broad, obtuse. *Basin* usually small, medium to rather shallow, narrow to moderately wide, obtuse to somewhat abrupt, furrowed, sometimes wrinkled.

Skin rather tough, smooth, glossy, yellowish-white, often almost entirely overspread and mottled with attractive pinkish-red, conspicuously striped and splashed with bright carmine. *Dots* few, small, inconspicuous, light colored, often submerged.

Calyx tube long, funnel-shape or nearly so. *Stamens* median.

Core rather large, abaxile; cells often unsymmetrical, wide open or closed; core lines clasping. *Carpels* broadly ovate to oval, smooth. *Seeds* small, moderately wide, plump, obtuse, not tufted.

Flesh white, moderately firm, tender, juicy, mild subacid, very aromatic, good to very good.

Season latter part of August and through September.

CLAPPER FLAT.

REFERENCE. 1. Downing. 1869:127.

SYNONYMS. CLAPPER FLAT (1). Flat (1).



CHENANGO

Downing describes a variety under this name which originated in the town of Bethlehem, Albany county, N. Y. He states (1) that the tree is productive, the fruit above medium size, pale yellow mostly overspread with deep red, pleasant subacid and good in quality for culinary uses; season September and October. We do not know this variety and have found no account of it except that given by Downing.

CLARKE.

REFERENCES. 1. *Rural N. Y.*, 19:375. 1868. 2. Downing, 1869:127. 3. Leroy, 1873:221. fig. 4. Burrill and McCluer, *Ill. Sta. Bul.*, 45:317. 1896. 5. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:39. 1903. 6. Beach and Clark, *N. Y. Sta. Bul.*, 248:115. 1904.

SYNONYMS. CLARKE (1, 2, 3, 4, 5, 6). *Clarke Beauty*.

This variety has been grown to a limited extent locally in some portions of Central New York. It is not a good commercial variety, being too tender and too easily bruised. It is very good for dessert. It is sometimes called Clarke Beauty. The tree is hardy, healthy and long-lived. It does not come into bearing very young but when mature is a reliable biennial cropper.

Historical. Originated with J. N. Clarke, Naples, Ontario county, N. Y. (1, 2). It is now rarely propagated.

TREE.

Tree large to medium, vigorous. *Form* upright spreading to roundish, rather dense. *Twigs* short, curved, stout; internodes medium. *Bark* brownish and olive-green, lightly mottled with scarf-skin; slightly pubescent. *Lenticels* scattering, medium size, oblong, slightly raised. *Buds* medium to large, broad, acute, free, slightly pubescent.

FRUIT.

Fruit medium to large, averaging above medium. *Form* roundish oblate to oblate conic or to oblong conic, usually faintly ribbed, unsymmetrical; not very uniform in shape. *Stem* short to medium in length, slender. *Cavity* acuminate, deep, rather narrow to moderately wide, usually partly russeted and often with narrow, broken, outspreading russet rays. *Calyx* small to rather large, closed or slightly open. *Basin* rather shallow to moderately deep, rather narrow, obtuse to moderately abrupt, slightly furrowed and wrinkled.

Skin thin, smooth, waxy, pale whitish-yellow or greenish, often faintly shaded with orange-red or sometimes blushed with crimson; under some conditions the fruit develops but a slight blush or none. *Dots* numerous, small, pale or russet, often submerged.

Calyx tube cone-shape.

Core medium to rather large, abaxile; cells open; core lines slightly clasping. *Carpels* broadly roundish, mucronate, slightly tufted. *Seeds* medium to rather large, moderately wide, plump, obtuse to acute, slightly tufted, rather light brown.

Flesh whitish, firm, moderately fine, crisp, tender, juicy, rather sprightly subacid, good to very good.

Season October to January; some portion of the fruit may keep till spring but by January it begins to deteriorate in flavor and quality.

CLYDE.

REFERENCES. 1. Barry, 1851:283. 2. Elliott, 1854:127. 3. Downing, 1857:129. 4. Hoffer, *N. A. Pomol.*, 1860. *col. pl.* 5. Warder, 1867:694. *fig.* 6. *Am. Pom. Soc. Cat.*, 1875:6. 7. *1a. Hort. Soc. Rpt.*, 1880:596. 8. *Mo. Hort. Soc. Rpt.*, 1884. 9. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 10. Bailey, *Am. Hort.*, 1892:236. 11. Thomas, 1897:259. 12. Budd-Hansen, 1903:65.

SYNONYMS. CLYDE (3, 8, 9, 10). CLYDE BEAUTY (1, 2, 4, 5, 6, 7, 11). CLYDE BEAUTY (12). *Mackie's Clyde Beauty* (2, 3, 4, 11).

A large, late fall apple. So far as we can learn it is now but little grown in this state. Lyon reports that in Michigan the tree is vigorous, upright, very productive, and the fruit desirable for market (9).

Historical. This is a late autumn variety which originated with Mr. Mackie, of Clyde, Wayne county (3, 4).

TREE.

Tree vigorous, spreading. *Twigs* reddish-brown.

FRUIT.

Fruit large. *Form* roundish to oblong conic, more or less ribbed. *Stem* short, sometimes fleshy. *Cavity* acute, deep, rather wide, furrowed. *Calyx* small, closed. *Basin* medium in depth, somewhat abrupt, furrowed.

Skin waxy, green or yellow, washed and mottled with dull red and striped with carmine becoming bright red on the exposed side.

Core large and open. *Seeds* small, brown. *Flesh* white, often tender, juicy, sprightly, pleasant subacid, good to very good.

Season October to December.

COLLAMER.

The Collamer or Collamer Twenty Ounce is a sport of the Twenty Ounce, from which it differs in being more highly colored. As compared with Twenty Ounce, it is less mottled and striped but more completely covered with red, which often extends in an unbroken blush over a considerable portion of the fruit. In the Twenty Ounce this is seldom or never seen, but the red is mottled or appears in heavy stripes and splashes. So far as we have been able to determine, Collamer is more regular in shape and, if ribbed at all, is less distinctly ribbed than Twenty Ounce. The tree dif-



COLLAME 2

fers from Twenty Ounce in that the bark of the young twigs is more distinctly tinged with red. The fruit being more attractive than Twenty Ounce, Collamer is worthy of consideration for commercial planting where an apple of the Twenty Ounce type is desired.

Except in the points of difference above noted, Collamer appears to be identical with Twenty Ounce, and the reader is referred to the description of that variety for a technical account of the tree and fruit.

Historical. This variety originated as a sport of the Twenty Ounce tree in the orchard of J. B. Collamer, Hilton, N. Y. Mr. Collamer began propagating it about 1900.

COLTON.

REFERENCES. 1. *Amer. Gard.*, 12:573. 1891. *figs.* 2. Bailey, *Am. Hort.*, 1892:237. 3. Munson, *Me. Sta. Rpt.*, 1893:132. 4. Beach, *N. Y. Sta. An. Rpt.*, 15:271. 1896. 5. *Am. Pom. Soc. Cat.*, 1899:16. 6. Bruner, *N. C. Sta. Bul.*, 182:20. 1903. 7. Farrand, *Mich. Sta. Bul.*, 205:44. 1903. 8. Budd-Hansen, 1903:67.

SYNONYMS. COLTON (1, 2, 3, 4, 7). COLTON Early (5, 6, 8). Early Colton (1).

Colton is a green or yellowish apple of fair to good quality, in season from the last of July to early September. The tree is a good grower, hardy, comes into bearing moderately young and yields good crops biennially.

Historical. Colton is said to have originated on the farm of Mr. Colton, Rowe, Franklin county, Mass., where it has been propagated since about 1840 under the name Early Colton (1).

TREE.

Tree large, vigorous with moderately long, stout, crooked branches. *Form* rather upright when young but eventually flat, spreading and open. *Twigs* moderately long, straight, moderately stout; internodes short. *Bark* dark brown, heavily mottled with scarf-skin; much pubescent. *Lenticels* quite numerous, rather conspicuous, medium to large, oblong, raised. *Buds* medium to large, broad, plump, acute, free, pubescent.

FRUIT.

Fruit medium in size. *Form* roundish, narrowing toward either end, slightly ribbed. *Stem* medium in length, stout. *Cavity* small, acute to slightly acuminate, shallow, narrow. *Calyx* medium in size, nearly closed; lobes long, rather recurved. *Basin* small, shallow, obtuse, wrinkled.

Skin pale greenish-yellow, sometimes with a shade of red. *Dots* numerous, large, greenish.

Calyx tube elongated funnel-form. *Stamens* median.

Core medium to rather large, somewhat abaxile; cells open; core lines clasping. *Carpels* broadly roundish.

Flesh whitish, rather coarse, crisp, juicy, mild subacid, fair to good.

Season last of July to early September.

COLVERT.

REFERENCES. 1. Warder, 1867:427. 2. Downing, 1869:131. 3. *Am. Pom. Soc. Cal.*, 1875:6. 4. Thomas, 1885:506. 5. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 6. Bailey, *Am. Hort.*, 1892:237. 7. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:39. 1903.

SYNONYMS. COLVERT (1, 2, 3, 4, 5, 6, 7). *Prussian* (2).

Ripens about with Twenty Ounce. It is inferior to that variety in size, color and quality, and is not as good a seller, but is more productive. The fruit is large, uniform in size, yellowish-green shaded and lightly striped with pinkish-red on the sunny side, smooth, showy and fairly attractive. It needs to be picked early to prevent loss from dropping. It is not a good keeper and is not much in demand among buyers, but sometimes it sells pretty well.

The tree is generally hardy, healthy and an excellent cropper. It generally succeeds well on any good apple land.

Historical. Origin uncertain (2). It has long been known and pretty widely disseminated but it is not much grown in New York. Even in those localities where it is best known the trees of this variety constitute less than one per cent of the orchards.

TREE.

Tree medium size to large, moderately vigorous to vigorous; branches long, medium stout, curved, crooked. *Form* upright spreading or roundish, open. *Twigs* above medium to long, usually nearly straight, moderately stout; internodes medium. *Bark* rather dark brownish-red, shaded with olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* scattering, medium, oblong, raised. *Buds* medium to large, broad, prominent, very plump, obtuse, free, pubescent. *Leaves* medium in size, broad.

FRUIT.

Fruit averages large, fairly uniform in size but rather variable in shape. *Form* oblate to oblate conic, obscurely ribbed, irregular and with sides sometimes unequal. *Stem* short, rather thick. *Cavity* acute to slightly acuminate, medium to nearly deep, medium in width to sometimes broad, usually very heavily russeted, sometimes compressed and frequently lipped. *Calyx* medium, closed or slightly open; lobes short, narrow, acuminate. *Basin* abrupt, medium in depth, narrow, slightly furrowed.

Skin very thick, rather tough, rather dull greenish-yellow, sometimes partly washed with red and striped and splashed with carmine. *Dots* inconspicuous, small, usually submerged; a few scattering ones are large and russet. *Prevailing color* greenish-yellow, not particularly attractive.

Calyx tube broadly conical to funnel-shape. *Stamens* median to basal.

Core axile, small; cells closed or partly open. *Carpels* broad-cordate, emarginate, tufted. *Seeds* large to above medium, wide, rather long, plump, acute; frequently they are abortive.

Flesh tinged with yellow, firm, nearly coarse, crisp, moderately tender, juicy, subacid, good.

Season October to January or February.

CONSTANTINE.

REFERENCES. 1. Leroy, 1873:335. fig. 2. Budd, *Ia. Hort. Soc. Rpt.*, 1880:524. 3. Gibb, *Montreal Hort. Soc. Rpt.*, 1881:155. 4. Hogg, 1884:95. 5. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:14. 6. *Id.*, 14:86. 1888. 7. Hoskins, *Rural N. Y.*, 51:682. 1892. fig.

SYNONYMS. *Berry Apple* (7). CONSTANTINE (2, 3). GRAND DUC CONSTANTIN (1). GRAND DUKE CONSTANTINE (4, 5, 6, 7). *Grand Duke Constantine* (2, 3). No. 457 (7). *Riabinouka* (7).

This fruit is of the Aport type and very closely resembles Alexander. The flesh is rather coarse, subacid and fair to good in quality. Some hold that it is rather better in flavor than Alexander. As grown at this Station, the fruit, as compared with that of Alexander, begins to ripen about a week later and continues longer in season. The trees are not so large and may be planted more closely together than those of Alexander. It is a reliable cropper, yielding good crops biennially or nearly annually. The percentage of marketable fruit is greater than that of Alexander because there are fewer drops, the apples are less apt to show cracks about the calyx and stem and the skin is less often discolored by chafing against the branches. We are not sure that it is as good a variety for commercial planting as Alexander, but it appears to be worthy of testing where a variety of the Alexander type is desired.

TREE.

Tree small to below medium size, at first moderately vigorous but with age it becomes a slow grower with short, stout, curved branches. *Form* spreading, open. *Twigs* moderately long, curved, slender; internodes long. *Bark* brown with some olive-green, lightly streaked with scarf-skin; pubescent near tips. *Lenticels* scattering, medium to small, oblong, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit large or very large. *Form* roundish conic flat at the base, varying to oblate conic, regular or somewhat ribbed, symmetrical. *Stem* below medium to long, rather slender to moderately thick. *Cavity* large, acuminate or acute, very deep, broad, symmetrical, russeted and with outspreading rays of greenish-russet. *Calyx* medium to rather large, usually somewhat open; lobes medium in width and length, acute. *Basin* narrow to medium in width, moderately deep to deep, abrupt, smooth or slightly wrinkled.

Skin thick, tough, smooth, waxy, clear greenish-yellow or whitish, mottled, marbled and blushed with bright red over nearly the whole surface with wide broken stripes of carmine radiating from the cavity, overspread with thin bloom. *Dots* whitish or pale russet. *Prevailing effect* bright red.

Calyx tube long, wide, funnel-shape or conical. *Stamens* median or below.

Core medium size, somewhat abaxile; cells open or partly closed; core lines somewhat clasping. *Carpels* broadly ovate or approaching cordate, emarginate. *Seeds* medium or below, moderately wide, short, thick, plump, obtuse, dark brown.

Flesh whitish, moderately firm, coarse, tender, juicy, sprightly subacid, fair to good; suitable for culinary use and market.

Season late September to November.

COOPER.

REFERENCES. 1. *Horticulturist*, 1:339, 484. 1847. 2. *Mag. Hort.*, 13:105, 200. 1847. 3. Cole, 1849:114. fig. 4. Thomas, 1849:147. 5. Barry, 1851:283. 6. *Horticulturist*, 6:181. 1851. 7. Emmons, *Nat. Hist. N. Y.*, 3:104. 1851. fig. 8. Elliott, 1854:127. 9. Gregg, 1857:41. 10. Hooper, 1857:26, 106, 109. 11. Downing, 1857:130. 12. *Am. Pom. Soc. Cat.*, 1862. 13. Warder, 1867:428. fig. 14. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 15. Bailey, *An. Hort.*, 1892:237. 16. Budd-Hansen, 1903:67.

SYNONYMS. *Beauty Red* (8, 11). COOPER (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16). *Lady Washington* (8, 11). *Seek-No-Further* of some, erroneously (8).

Fruit large, uniform, very attractive, rather light yellow indistinctly streaked with mixed red, mild subacid or nearly sweet, season October to December. The tree is very vigorous, upright spreading. Not recommended for planting in New York.

Historical. This is an old variety of unknown origin. In 1796 it was introduced from Connecticut into Ohio where it has been much esteemed (2). Evidently it has never been cultivated to any considerable extent in this State and is now practically unknown to New York fruit growers.

CORNELL.

REFERENCES. 1. Downing, 1857:131. 2. *Am. Pom. Soc. Cat.*, 1862. 3. Warder, 1867:716. 4. Thomas, 1875:200. 5. Barry, 1883:337. 6. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 7. Bailey, *An. Hort.*, 1892:237. 8. Van



CONSTANTINE

Deman, *Rural N. Y.*, 61:671. 1902. fig. 9. Farrand, *Mich. Sta. Bul.*, 205:44. 1903. 10. Budd-Hansen, 1903:67.

SYNONYMS. CORNELL (6, 8, 9). CORNELL FANCY (1, 2, 3, 4, 5, 7). *Cornell Fancy* (8). *CORNELL Fancy* (10). *Cornell's Favourite* (1).

Fruit usually of good medium size, sometimes large, waxen yellow and red, agreeable for dessert, in season from early September to November. The tree sometimes lacks vigor and productiveness (9).

Historical. Origin Pennsylvania (1). It is but little known in New York.

FRUIT.

Fruit above medium to large, uniform in size, somewhat variable in shape. *Form* roundish conic to oblate conic, often quite strongly ribbed, irregular; sides usually unequal. *Stem* medium to rather long and slender. *Cavity* moderately deep to deep, moderately wide, often compressed, sometimes lipped, sometimes russeted, with the russet extending beyond the cavity. *Calyx* below medium to rather large, closed or slightly open; lobes sometimes separated at the base, often upright, moderately acute. *Basin* deep, wide, rather abrupt, strongly furrowed, slightly wrinkled.

Skin moderately thick, tough, smooth, clear pale waxen yellow, partly overspread with thin attractive pinkish-red, often quite regularly splashed and striped with bright carmine. *Dots* conspicuous, variable, often large, irregular, russet or areolar with russet center, varying to small, light colored and submerged. *Prevailing effect* handsome red striped over clear yellow.

Calyx tube rather large, rather short, conical. *Stamens* median.

Core below medium, variable, nearly axile to decidedly abaxile; cells variable, open or closed; core lines meeting or slightly clasping. *Carpels* broadly ovate, slightly emarginate, sometimes tufted. *Seeds* numerous, rather large, dark brown, rather narrow, long, plump, acute to acuminate, sometimes tufted.

Flesh tinged with yellow, often affected with "Baldwin Spot," firm, moderately coarse, crisp, moderately tender, juicy, agreeable, mild subacid, aromatic, rich, sprightly, very good.

Season early September to November.

CORNER.

REFERENCE. 1. Heiges, *U. S. Pom. Rpt.*, 1894:18.

SYNONYMS. None.

We have neither seen Corner nor received any report concerning it. The following account of it was given in 1894 by S. B. Heiges, then United States Pomologist (1).

"Size above medium; oblate; cavity wide, deep, marked by russet netting; stem one-half inch, medium diameter; basin, medium, regular, marked by russet; calyx segments with mammiform bases, wide, long, converging or slightly reflexed; surface moderately smooth; color yellow, washed with red and striped with crimson; dots numerous, russet, some with dark centers, depressed; flesh yellowish, moderately fine grained, tender, moderately juicy; core large, wide, clasping, closed; flavor mild subacid; quality very good. Season early winter. Well known locally in Orange county, N. Y."

COX ORANGE.

REFERENCES. 1. *Horticulturist*, 13:168. 1858. 2. Downing, 1869:135. 3. Leroy, 1873:517. fig. 4. *Am. Pom. Soc. Cat.*, 1881:8. 5. Hogg, 1884:55. 6. Thomas, 1885:507. 7. Bailey, *Am. Hort.*, 1892:237. 8. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:356. 9. Willard, *Rural N. Y.*, 58:754. 1899. 10. Thomas, *Garden*, 59:34. 1901. figs.

SYNONYMS. *Cos Orange* (3). *COX ORANGE* (8). *Cox's Orange* (3). *COX'S ORANGE PIPPIN* (1, 2, 4, 5, 6, 7, 9, 10). *Cox's Orange Pippin* (3). *ORANGE DE COX* (3). *Reinette Orange de Cox* (3).

One of the best in quality of the English dessert apples; in season from late September to early winter. The fruit is of medium size or above medium, red and yellow. When highly colored it is attractive, with the red predominant. The tree is a moderate grower and productive. It is well adapted for growing on dwarf stock, either Paradise or Doucin. It is not recommended for commercial planting, but it is a desirable variety for the home orchard.

Historical. Cox Orange is said to have originated in 1830 from seed of Ribston, at Colnbrook Lawn near Slough, Bucks, England (5). It is sometimes propagated by American nurserymen but it has never been extensively planted in this country and its cultivation is not increasing.

TREE.

Tree medium size or above, moderately vigorous with rather slender branches. *Form* upright, thickly branched, dense. *Twigs* long to medium, rather slender, irregularly crooked; internodes medium or below. *Bark* olive-green somewhat mottled with reddish-brown, slightly pubescent. *Lenticels* numerous, conspicuous, medium size, oblong, raised. *Buds* medium size to rather small, roundish, obtuse, appressed, pubescent. *Leaves* small to medium size and inclined to be narrow.

FRUIT.

Fruit medium or above, pretty uniform in size and shape. *Form* roundish oblate, sometimes slightly inclined to conic, regular or faintly ribbed, symmetrical, axis sometimes oblique. *Stem* usually obliquely inclined, short, thick, sometimes long. *Cavity* obtuse to somewhat acuminate, rather shallow to moderately deep, rather narrow, often somewhat russeted. *Calyx* rather small, closed or partly open. *Basin* rather shallow and obtuse to moderately deep and abrupt, rather narrow to moderately wide, smooth or slightly furrowed.

Skin rather thin, tough, smooth, attractive, washed with orange-red deepening to bright red and mottled and splashed with carmine, over a deep yellow background. *Dots* conspicuous, large, areolar with pale gray or russet center. *Calyx tube* cone-shape or funnel-form. *Stamens* median to basal.



COX ORANGE

Core medium size, somewhat abaxile; cells usually symmetrical, open or closed; core lines clasping the funnel cylinder. *Carpels* thin, obovate to obcordate, emarginate, usually smooth. *Seeds* reddish-brown, above medium size, wide, obtuse to acute, often abortive.

Flesh yellow, firm, nearly fine, crisp, tender, very juicy, rich, sprightly subacid or becoming mild subacid, decidedly aromatic, very good to best.

Season late September to January.

CRANBERRY PIPPIN.

REFERENCES. 1. Downing, 1845:106. 2. Thomas, 1849:179. 3. Emmons, *Nat. Hist. N. Y.*, 3:88. 1851. 4. Elliott, 1854:169. 5. Warder, 1867:402. 6. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 7. Bailey, *Am. Hort.*, 1892:237. 8. *Amer. Gard.*, 16:425. 1895. 9. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 2:9. 1895. fig. 10. *Ib.*, 3:6. 1896. fig. 11. Macoun, *Can. Dept. Agr. Bul.*, 37:43. 1901. 12. Beach and Clark, *N. Y. Sta. Bul.*, 248:116. 1904.

SYNONYMS. None.

In some sections this has proved a desirable apple, but in others it has not been successful. It is well known in some localities in the Hudson valley, in Northern and Western New York and in Ontario, where it is favorably regarded as a fall or early winter apple because of its good size, bright and attractive color and its uniform size and shape. It is a good storage apple, stands shipping well and brings good prices. It is suitable for market, cooking and evaporating, but not for dessert. It appears to be quite resistant to the attacks of scab. The trees are hardy and often very productive, but in some cases it is reported as undesirable because unproductive. It is said to be a shy bearer when young, but becomes productive with age.

Historical. Originated near Hudson, Columbia county (1).

TREE.

Tree large, very vigorous; branches stout, spreading. *Form* upright becoming somewhat spreading. *Twigs* long, moderately stout, light grayish-brown, quite pubescent; internodes short. *Bark* dull reddish-brown with some olive-green and thickly mottled with scarf-skin. *Lenticels* scattering medium to small, usually roundish. *Buds* medium or sometimes small. Rather broad, deeply set, obtuse or sometimes acute, pubescent, appressed. *Leaves* dark green, broad, medium to large; foliage rather dense.

FRUIT.

Fruit large. *Form* roundish oblate, symmetrical. *Stem* short. *Cavity* broad, wavy. *Calyx* closed or somewhat open. *Basin* moderately deep, russeted. *Skin* smooth, shining, clear light yellow, handsomely blushed, striped

and splashed with scarlet. *Dots* many, large, often red areolar with russet center. General appearance beautiful and attractive. *Flesh* white or with slight yellowish tinge, moderately juicy, mild subacid.

Season October to February. In the vicinity of its origin its season closes from a month to six weeks earlier than either Hubbardston or Tompkins King. In Northern New York and Ontario its season is late fall and early winter and often extends to midwinter.

CREAM.

REFERENCES. 1. *N. E. Farmer*, 1831 (cited by 3). 2. Downing, 1869:137. 3. Ragan, *U. S. B. P. I. Bul.*, 56:82. 1905.
SYNONYMS. None.

This variety originated in Queens county, N. Y. So far as we know it is no longer cultivated. Downing describes the tree as a vigorous grower and an early bearer and the fruit as medium or below, yellowish, fine-grained, pleasant, sweet, in season in September and October. Valued by some for dessert and culinary uses.

CROW EGG.

REFERENCES. 1. Kenrick, 1832:43. 2. Downing, 1857:211. 3. Warder, 1867:716. 4. Burrill and McCluer, *Ill. Sta. Bul.*, 45:318. 1896.
SYNONYMS. CROW EGG (2). CROW'S EGG (1, 3, 4). *Egg Jop?* (2).

A sweet apple which is still occasionally found in very old orchards but is now practically obsolete. Some esteem it highly for dessert. Downing calls it not very good in quality (2). The old trees are productive.

TREE.

Tree moderately vigorous. *Form* upright spreading; top roundish, open; branches long, slender, crooked. *Twigs* medium in size, curved, slender; internodes very short. *Bark* reddish-brown, streaked with scarf-skin, slightly pubescent. *Lenticels* numerous, very small, oblong. *Buds* small, plump, obtuse, deeply set in the bark. *Leaves* medium in size, narrow.

FRUIT.

Fruit about medium in size. *Form* roundish to oblong or ovate. *Stem* long, slender. *Cavity* obtuse to sometimes acute, shallow, medium in width, symmetrical or obscurely furrowed, bright green or sometimes with out-spreading russet. *Calyx* small to medium, closed. *Basin* small, shallow, narrow, somewhat abrupt, furrowed and wrinkled.


Skin tough, nearly smooth, bright pale yellow or greenish sometimes with faint bronze bluish. *Dots* numerous, very small but conspicuous, russet.

Calyx tube rather small, funnel-shape or cone-shape. *Stamens* median.


Core large, abaxile; cells usually symmetrical and open; core lines clasping the funnel cylinder or meeting when the tube is conical. *Carpels* ovate, nearly smooth. *Seeds* numerous, rather light brown, flat, acute to acuminate.

Flesh whitish, firm, crisp, tender, rather juicy, sweet, agreeably flavored, good to very good.

Season October and November.



CRANBERRY PIPPIN



CRANBERRY PIPPIN

CZAR THORN.

REFERENCES. 1. *Montreal Hort. Soc. Rept.*, 1881:53. 2. *Ib.*, 1883:75. 3. *Id.*, *Hort. Soc. Rpt.*, 1883:430. 4. Budd, *Id.*, *Agr. Coll. Bul.*, 1885:7. 5. Gibb, *Am. Pom. Soc. Rpt.*, 1887:44. 6. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57. 7. Beach, *W. N. Y. Hort. Soc. Rept.*, 1896:50. 8. Hansen, *S. D. Sta. Bul.*, 76:43. 1902. fig.

SYNONYMS. CZAR THORN (3, 4, 5, 6, 7, 8). *Czar Thorn* (2). *Czarskui Schip* (4). *No. 140 M* (8). *No. 206* (4, 5, 8). *Tars Thorn* (1). *Tsarskui Schip* (5). *Zarskischip* (7). ZARSKI SCHIP (2). *Zarski Schip* (3). ZARSKI ZARS (1).

A Russian apple of medium size, roundish conic, green and yellow usually shaded and striped with crimson; flesh rather coarse, sweet, hardly fair in quality; season September; not valuable.

DEADERICK.

REFERENCES. 1. *U. S. Pom. Rpt.*, 1895:22. 2. Watts, *Tenn. Sta. Bul.*, 1:11. 1896. fig. 3. Taylor, *Am. Pom. Soc. Rpt.*, 1897:37.

SYNONYMS. *Ben Ford* (2). DEADERICK (1, 2, 3). *Ozark Pippin* (2).

A good-sized green apple, of somewhat better color than *Rhode Island Greening*, but it does not keep as well, and is inferior to that variety in quality. The tree is a strong grower, healthy, and so far as tested here comes into bearing young and gives promise of being very productive. It has not been on trial long enough to indicate whether or not it has sufficient merit to be considered a promising variety for this state. In Tennessee it is considered a very valuable early winter apple (2).

Historical. Originated with Benjamin Ford, Washington county, Tenn. It was first disseminated as *Ozark Pippin* (2).

TREE.

Tree rather vigorous. *Form* spreading and somewhat upright. *Twigs* moderately stout, nearly straight; internodes short. *Bark* bright brownish-red. *Lenticels* roundish, often conspicuous, scattering, small. *Buds* medium size, appressed, obtuse, short, pubescent. *Leaves* medium size, somewhat narrow; often the base of the petioles is conspicuously streaked with red.

FRUIT.

Fruit large. *Form* broadly roundish, often rather conical, sometimes broadly ribbed, pretty regular, uniform. *Stem* medium to rather long, slender. *Cavity* large, acute to acuminate, deep, broad, usually smooth and symmetrical, sometimes slightly furrowed, occasionally prominently lipped. *Calyx* small, partly open or closed; lobes rather narrow, acute. *Basin* small, shallow, obtuse to somewhat abrupt, nearly smooth, a little wrinkled.

Skin moderately thick, tough, smooth, rather bright green becoming yellow, usually partly covered with a thin pinkish-red blush upon which there are often seen red, areolar dots with russet or whitish center; commonly the dots are whitish and often submerged. *Prevailing color* green or yellowish.

Calyx tube long, funnel-form. *Stamens* median to nearly marginal.

Core a little abaxile, medium to small; cells symmetrical, open or nearly so; core lines clasp the base of the cylinder. *Carpels* thin, generally smooth, broadly roundish, narrowing toward the base, slightly emarginate. *Seeds* numerous, medium or above, rather wide, obtuse.

Flesh yellowish, firm, moderately coarse, tender, rather juicy, pleasant sub-acid, good.

Season October to January.

DETROIT RED.

REFERENCES. 1. *Mag. Hort.*, 10:167. 1844. 2. Downing, 1845:106. 3. *Horticulturist*, 1:361. 1846. 4. *N. Y. Agr. Soc. Trans.*, 1846:192. 5. Thomas, 1849:164. 6. Cole, 1849:115. 7. Emmons, *Nat. Hist. N. Y.*, 3:65. 1851. fig., col. pl. 8. Elliott, 1854:130. 9. Downing, 1857:134. 10. Warder, 1867:532. fig. 11. *Am. Pom. Soc. Cat.*, 1873. 12. Wickson, 1889:245. 13. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 14. Bailey, *Am. Hort.*, 1892:237. 15. Waugh, *Vt. Sta. An. Rpt.*, 14:291. 1901.

SYNONYMS. *Black Apple* of some (2, 9). BLACK DETROIT (5, 7). *Black Detroit* (2, 8). *Crimson Pippin* (6, 8, of some 2). DETROIT (1, 2, 4, 6, 8). *Detroit* (9). DETROIT BLACK (10). DETROIT RED (3, 9, 11, 12, 13, 14, 15). *Detroit Red?* (10).

Fruit growers in Western New York have commonly used the names Detroit Red and Detroit Black interchangeably for the remarkably variable variety which we are here describing as Detroit Red. We have been unable to determine whether there are in fact two distinct varieties of this type, or whether the differences which have been observed in the habit of growth and productiveness of the tree and in the form, size, general appearance, season and quality of the fruit, are altogether due to differences in the conditions under which the fruit has been produced. Speaking of these two names, Warder, in 1867, wrote, "I have put these two names together because the fruits presented as Black and as Red Detroit are so very much alike in all respects that it is not worth while to consider them distinct. * * * The Red variety may be distinct, as it keeps later."

Lyon (13) recognized two or more varieties of this type and distinguished them by the names Detroit Black and Detroit Red. Speaking of Detroit Red, he remarks: "There are probably several



DETROIT RED

varieties grown under this name, none of them valuable;" and of The Detroit Black he says: "Unproductive, showy, valueless; it is probably the Detroit Red of Downing."

D. D. Stone, of Oswego, writes:¹ "It seems to me that the two are not the same. Detroit Black seems to be more of a scrubby grower, the shape, size, firmness and color seem to be more constant and it does not crack so badly as the one we know as Detroit Red, but the season appears to be the same."

Detroit Red, or as it is often called, Detroit Black, as usually grown in Western New York, varies from medium to very large, commonly averaging about medium size. It is flattened at the ends, very dark crimson or purplish, becoming almost black, with snow-white flesh occasionally streaked with rose-pink. It is esteemed by many for dessert use because of its mild, pleasant flavor. There is considerable loss from premature dropping of the fruit and from fruit that is too small or too ill-shapen for market. It is quite variable in keeping qualities, being commonly in season about with Maiden Blush. The tree is a moderate grower, comes into bearing rather young, and is not a very reliable cropper. Some report that it is a shy bearer; others that it yields moderate to full crops biennially.

Historical. This is supposed to have been brought into the neighborhood of Detroit by the early French settlers and thence disseminated (1, 2, 4, 8). It was introduced into Ohio and Western New York before the middle of the last century. The variety is still sometimes listed by nurserymen (14). Its cultivation in New York state is declining and it is now seldom planted.

TREE.

Tree medium or eventually large, moderately vigorous to vigorous. *Form* upright spreading or roundish. *Twigs* medium to long, moderately slender, dark brown.

FRUIT.

Fruit very large to medium. *Form* oblate conic varying to roundish oblate, often strongly ribbed, irregular. *Stem* short, usually rather slender. *Cavity* often very large, acute to nearly acuminate, deep, moderately broad to very broad, frequently compressed, usually thinly russeted. *Calyx* variable, usually large, closed or somewhat open; lobes short, broad, rather obtuse. *Basin* medium in width and depth, obtuse to somewhat abrupt, irregularly furrowed and wrinkled and often with mammiform protuberances.

¹ Letter, 1904.

Skin thick, rather tough, dark crimson, largely striped and splashed with purplish-carmine eventually becoming almost black, sometimes having a portion of the greenish-yellow ground color exposed. *Dots* numerous, conspicuous, very small, pale or russet.

Calyx tube rather short, wide, broadly conical. *Stamens* median to basal.

Core medium to large, axile; cells closed; core lines meeting or clasping. *Carpels* roundish to elliptical, rather flat, usually tufted. *Seeds* rather large, plump, obtuse to acute, moderately light brown.

Flesh white, sometimes streaked or stained with red, rather coarse, tender, juicy, agreeable mild subacid, very aromatic, good to very good.

Season last of September to December.

DUCHESS OF OLDENBURG.

This variety is often called Duchess or Duchess of Oldenburg, but the name now accepted for it by pomologists is Oldenburg, under which name it is described on page 150.



DUDLEY.

REFERENCES. 1. *Me. Sta. Rpt.*, 1891:97. 2. *U. S. Pom. Ppt.*, 1891:390. 3. Bailey, *Am. Hort.*, 1892:245. 4. Munson, *Me. Sta. Rpt.*, 1893:132. 5. Hoskins, *Amer. Gard.*, 14:299. 1893. 6. Munson, *Me. Sta. Rpt.*, 1902:83, 91. 7. Budd-Hansen, 1903:71. 8. Macoun, *Can. Dept. Agr. Rpt.*, 1903:95.

SYNONYMS. DUDLEY (2, 6, 8). DUDLEY WINTER (1, 4). DUDLEY WINTER (7). *Dudley's Winter* (6). NORTH STAR (3, 5). *North Star* (6, 7, 8).

A very hardy and productive variety which is being planted to a considerable extent in Northern New England. The fruit is pretty large, bright greenish-yellow washed and splashed with red, quite attractive in appearance and good in quality. Munson says that it is perhaps now more widely grown than any other of the newer sorts that have originated in New England. He considers it a valuable acquisition as a winter fruit for northern localities (6). As fruited at Geneva it is in season in September and October, although it may sometimes be kept into the winter. It is recommended for trial particularly where a very hardy apple of its season is desired.

Historical. A seedling of the Oldenburg, which originated with J. W. Dudley, Castle Hill, Aroostook county, Me. (1, 6). A few years ago it was introduced by a Rochester nursery under the name North Star but it was afterward found that this name had already been given to another variety and therefore the name Dudley Winter was retained for it, which, according to the accepted rules of nomenclature is shortened to Dudley.



DUDLEY (Reduced Size)

TREE.

Tree small, moderately vigorous to vigorous; branches short, moderately stout. *Form* very spreading and drooping, rather dense. *Twigs* below medium length to short, almost straight, moderately stout to rather slender; internodes short to medium. *Bark* brown tinged with clear bright red, with but little or no scarf-skin and but slightly pubescent. *Lenticels* rather conspicuous, clear in color, scattering, medium in size, oblong, raised. *Buds* medium or above, rather prominent, plump, obtuse to acute, free or nearly so, somewhat pubescent.

FRUIT.

Fruit medium to large, uniform. *Form* roundish conic to roundish oblate, symmetrical. *Stem* long, rather thick. *Cavity* acute to almost acuminate, rather deep, broad, sometimes lightly russeted, obscurely furrowed. *Calyx* medium to large, open or partly closed. *Basin* decidedly abrupt, moderately deep to deep, moderately broad, obscurely furrowed, wrinkled.

Skin thin, tender, smooth, bright pale yellow or whitish mostly covered with a bright pinkish-red blush striped and splashed with bright carmine and covered with light bloom. *Dots* scattering, light, small. *General appearance* red or red striped over contrasting yellow, attractive.

Calyx tube long, moderately wide, funnel-shape or sometimes conical. *Stamens* median to marginal.

Core almost axile, medium or below; cells closed or partly open; core lines clasping or nearly so. *Carpels* broadly elliptical, not emarginate, slightly tufted. *Seeds* large, wide, long, somewhat flat, obtuse to acute, dull dark brown.

Flesh tinged with yellow, firm, crisp, nearly fine-grained, tender, very juicy, aromatic, brisk subacid eventually becoming mild, very good.

Season September and October or sometimes later.

DYER.

REFERENCES. 1. Kenrick, 1835:60. 2. *Mag. Hort.*, 3:37. 1837. 3. Downing, 1845:83. 4. *Horticulturist*, 2:289. 1847. 5. Cole, 1849:111. 6. Thomas, 1849:153. fig. 7. Barry, 1851:283, 286. 8. Emmons, *Nat. Hist. N. Y.*, 3:39. 1851. 9. Elliott, 1854:75. 10. Hooper, 1857:30. 11. Hovey, *Mag. Hort.*, 27:70. 1861. fig. 12. *Am. Pom. Soc. Cat.*, 1862. 13. Warder, 1867:639. fig. 14. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 15. Bailey, *Am. Hort.*, 1892:238. 16. Hansen, *S. D. Sta. Bul.*, 76:47. 1902. 17. Budd-Hansen, 1903:72. fig.

SYNONYMS. *Bard Apple* (9, 11). *Beard Burden* (9, 11). *Bullripe* (9, 11). *Coe's Spice* (11). *DYER* (1, 3, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17). *Dyer* (4, 5, 7). *Golden Spice* (9, 11). *Mygatt's Bergamot* (9, 11). *POMME ROYAL* (2). *Pomme Royal* (6, 8, 9, 10, 11, 14, 16, 17). *POMME ROYALE* (4, 5, 7). *Pomme Royale* (3, 13). *Pomme Roye* (4). *Pomme Water* (9, 11). *Pomme-water* in Ill. (13). *Smithfield Spice* (3, 5, 10, 11). *Tompkins* (9, 11). *White Spice* (9, 11). *Woodstock* (1).

One of the very finest dessert apples but not a good commercial variety (14). The fruit is of medium size, greenish-yellow with a shade of red. The crop does not ripen evenly and it requires more than one picking. It comes in

season late in August or early in September and ripens continuously until midautumn. The tree is vigorous in the nursery but does not grow to be a large tree in the orchard. It succeeds better when topworked upon some hardier vigorous stock such as Tolman Sweet or Northern Spy. It is not long-lived but comes into bearing rather young and yields good crops biennially.

Historical. This variety has been supposed by some to be of French origin and was formerly known as Pomme Royale, but Hovey believed it to be an American apple (11). It was known in cultivation in Rhode Island during the Revolutionary War (4). It was named Dyer by the Massachusetts Horticultural Society more than fifty years ago and has retained that name. It is still occasionally listed by nurserymen (15). It is but little cultivated in New York and is now seldom, if ever, planted in this state.

FRUIT (3, 9, 11, 13).

Fruit medium or sometimes large. *Form* roundish, slightly oblate, regular or obscurely ribbed. *Stem* medium to long, slender. *Cavity* rather small, acute, moderately deep to deep, sometimes lipped. *Calyx* small, closed; lobes short to rather long, recurved. *Basin* medium to small, shallow to moderately deep, furrowed.

Skin smooth, clear pale yellow or greenish, more or less flecked and marbled with thin russet with a brownish blush on one cheek. *Dots* dark or russet.


Core medium size; cells open or closed; core lines clasping. *Seeds* numerous, plump, short, medium to small, pale.

Flesh yellowish-white, fine, very crisp, tender, aromatic, sprightly, mild subacid, highly flavored, very good to best.

Season September and October.

EARLY HARVEST.

- REFERENCES. 1. *Amer. Gard. Cal.*, 1806:584. 2. Cox, 1817:101. fig. 3. Thacher, 1822:129. 4. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 5. Wilson, 1828:136. 6. Fessenden, 1828:131. 7. *London Hort. Soc. Cat.*, 1831:No. 355. 8. Kenrick, 1832:26. 9. Floy-Lindley, 1833:84. 10. *Mag. Hort.*, 1:362. 1835. 11. Manning, 1838:45. 12. *Ib.*, *Mag. Hort.*, 7:51. 1841. 13. Downing, 1845:72. fig. 14. French, *Horticulturist*, 1:256. 1846. 15. Hovey, *Mag. Hort.*, 14:115. 1848. fig. 16. Cole, 1849:97. fig. 17. Walker, *Mag. Hort.*, 15:165. 1849. 18. Thomas, 1849:142. 19. Emmons, *Nat. Hist. N. Y.*, 3:16. 1851. 20. Barry, 1851:280. 21. Hovey, 1:75. 1851. col. pl. and fig. 22. *Am. Pom. Soc. Cat.*, 1852. 23. Elliott, 1854:84. fig. 24. Hooper, 1857:31, 107, 111. 25. Gregg, 1857:36. fig. 26. Warder, 1867:403. fig. 27. Fitz, 1872:143, 160, 172. 28. Downing, 1872:10 index, app. 29. *Ib.*, 1881:11 index, app. 30. Hogg, 1884:67. 31. Wickson, 1889:243. 32. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 33. Bailey, *Am. Hort.*, 1892:238. 34. Stinson, *Ark. Sta. An. Rpt.*, 7:44. 1894. 35. Burrill and McCluer, *Ill. Sta. Bul.*, 45:320. 1896. 36. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 3:7. 1896. figs. 37. *Can. Hort.*, 20:328. 1897. figs. 38. Alwood, *Va. Sta. Bul.*, 130:121. 1901. 39. Waugh, *Vt. Sta. An. Rpt.*, 14:292. 1901. 40. Budd-Hansen, 1903:73. fig.



EARLY HARVEST



EARLY HARVEST

SYNONYMS. *Early French Reinette* (2, 8, 13, 15, 18, 19, 21, 23, 30, 34). *EARLY HARVEST* (1, 7, 8, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40). *Early Harvest* (9, 23). *Early July Pippin* (24). *Early June* (23). *HARVEST* (23). *Harvest* (21). *July Early Pippin* (15, 21, 30). *JULY PIPPIN* (9). *July Pippin* (13, 15, 18, 19, 21, 23, 30, 34). *Large Early* (15, 21, 30). *Large Early Harvest* (15, 21). *LARGE EARLY HARVEST* (3). *Large White Juneating* (13, 15, 19, 21, 23, 30). *Maralandica* (29). *Oats* incorrectly (29). *Pomme d'Ete* of Canada (30). *Prince's Early Harvest* (8, 30). *PRINCE'S HARVEST* (2, 4, 5, 6). *Prince's Harvest* (8, 13, 15, 18, 19, 21, 23, 24, 30, 34). *Prince's Yellow Harvest* (30). *Tart Bough* (13, 15, 19, 21, 30). *Yellow Harvest* (13, 15, 16, 18, 19, 21, 23, 24, 30, 34). *Yellow Juneating* (23).

Fruit medium, pale yellow, sometimes with a faint blush, tender, sprightly subacid, and very good in quality. It is a desirable variety for the home orchard because it is one of the earliest of the summer apples, and is excellent for either dessert or culinary uses. It is not a desirable commercial variety because there is a comparatively large percentage of undersized or otherwise unmarketable fruit, the color is such that it shows bruises very readily and it keeps but a short time. The tree is a fairly good grower, moderately long-lived, comes into bearing rather early, is a biennial or almost annual cropper and moderately productive.

Historical. Early Harvest has been known in cultivation for more than a hundred years. Its origin is unknown but it is supposed to have originated in America. At one time it was quite extensively cultivated for local market in some localities but it is now seldom or never planted except for home use.

TREE.

Tree medium size, moderately vigorous. *Form* upright spreading or roundish, open. *Twigs* moderately long, curved, rather stout; internodes short. *Bark* dark brown with some olive-green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, round, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit usually medium or below but sometimes rather large, uniform in size and shape. *Form* oblate to nearly round, regular or slightly angular; sides slightly unequal. *Stem* medium in length, moderately thick. *Cavity* nearly acuminate, shallow, rather narrow to moderately broad, russeted and with outspreading, broken russet rays. *Calyx* small to medium, closed; lobes long, narrow. *Basin* shallow, moderately wide, obtuse, slightly wrinkled.

Skin thin, tender, very smooth, clear pale waxen yellow, occasionally with deeper yellow on exposed cheek, sometimes slightly blushed. *Dots* numerous, large and small, submerged or russet.

Calyx tube short, funnel-shape. *Stamens* medium.

Core medium size, somewhat abaxile; cells closed or slightly open; core lines clasping. *Carpels* slightly obovate. *Seeds* small to rather large, narrow, long, plump, acute.

Flesh white, not firm, rather fine, crisp, tender, juicy, at first briskly subacid but eventually becoming milder, and more agreeable for dessert. Good to very good.

Season late July and August.


EARLY JOE.

REFERENCES. 1. *N. Y. Agr. Soc. Trans.*, 1843:52. 2. *Mag. Hort.*, 9:469. 1843. 3. *N. Y. Agr. Soc. Trans.*, 1846:187. fig. 4. Hovey, *Mag. Hort.*, 12:474. 1846. 5. *Ib.*, 13:159. 1847. fig. 6. Smith, *Horticulturist*, 1:386. 1847. fig. 7. *Cultivator*, 4:310. 1847. fig. 8. *Mag. Hort.*, 14:539. 1848. 9. Cole, 1849:105. fig. 10. Thomas, 1849:137. fig. 11. Emmons, *Nat. Hist. N. Y.*, 3:19. 1851. 12. Barry, 1851:280. 13. Elliott, 1854:75. fig. 14. Gregg, 1857:36. 15. Downing, 1857:76. 16. Hooper, 1857:32. 17. *Am. Pom. Soc. Rpt.*, 1860:240. 18. Warder, 1867:513. 19. Fitz, 1872:152, 172. 20. Leroy, 1873:404. figs. 21. *Ill. Hort. Soc. Rpt.*, 1877:124. 22. Hogg, 1884:68. 23. Wickson, 1889:243. 24. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 25. Bailey, *Am. Hort.*, 1892:238. 26. Macoun, *Can. Dept. Agr. Rpt.*, 1901:96. 27. Waugh, *Vt. Sta. An. Rpt.*, 14:292. 1901. 28. Budd-Hansen, 1903:74. fig.


SYNONYMS. EARLY JOE (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28). JOE PRÉCOCE (20).

Fruit medium to small, red striped, excellent in flavor and quality for dessert use; season, August and early September. The tree is small to medium in size, slow growing, moderately long-lived, comes into bearing rather young and is a reliable biennial cropper. The fruit hangs pretty well to the tree until it is quite ripe. The crop contains a large percentage of undersized or otherwise unmarketable fruit. Recommended for the home orchard, but not for commercial planting.

Historical. Originated with Northern Spy and Melon in the orchard of Heman Chapin, East Bloomfield, Ontario Co., N. Y. This orchard was planted with seedling trees grown from seeds brought from Salisbury, Conn. about the year 1800. In October, 1843, Early Joe was exhibited at the fair of the New York State Agricultural Society, Rochester, N. Y., by Jonathan Buel of East Bloomfield (1). It has been widely disseminated and is still listed by nurserymen (25) but it is not cultivated extensively in any locality.



EARLY JOE



TREE.

Tree moderately vigorous, dwarfish with short, moderately stout, crooked branches. *Form* rather flat, spreading. *Twigs* short, straight, stout, with large terminal buds; internodes short. *Bark* dark brown, lightly streaked with scarf-skin; pubescent. *Lenticels* scattering, medium size, oblong, slightly raised. *Buds* medium size, plump, acute, free, slightly pubescent.

FRUIT.

Fruit small to medium, uniform in size and shape. *Form* oblate conic to conic, somewhat ribbed, rather symmetrical. *Stem* medium to long, rather slender to moderately thick. *Cavity* acute, shallow to medium in depth, rather broad, symmetrical, sometimes thinly russeted. *Calyx* medium size, closed or slightly open. *Basin* small to medium, usually shallow, medium in width or rather narrow, somewhat abrupt, smooth or slightly wrinkled.

Skin thin, tender, smooth, pale greenish-yellow, irregularly and obscurely striped and splashed with dull, dark red, in highly colored specimens becoming deeply blushed on the exposed cheek. *Dots* russet and greenish or nearly white.

Calyx tube medium in length, rather wide, broadly conical. *Stamens* median or basal.

Core medium to rather small, axile; cells slightly open or closed; core lines clasping. *Carpels* broadly obcordate to elliptical, decidedly concave. *Seeds* small to medium, rather wide, short, obtuse to acute.

Flesh tinged with yellow, fine, crisp, very tender, very juicy, mild subacid, very good to best.

Season August and September.

EARLY PENNOCK.

REFERENCES. 1. Humrickhouse, *Mag. Hort.*, 12:472. 1846. fig. 2. Cole, 1849:104. 3. Emmons, *Nat. Hist. N. Y.*, 3:14. 1851. 4. Barry, 1851:332. 5. Hooper, 1857:33, 106, 110. 6. Gregg, 1857:36. 7. Downing, 1857:137. 8. *Am. Pom. Soc. Cat.*, 1862. 9. Warder, 1867:594. fig. 10. Fitz, 1872:145. 11. *Ill. Hort. Soc. Rpt.*, 1875:49. 12. Thomas, 1875:191. 13. Downing, 1881:11 index, app. 14. Bailey, *Am. Hort.*, 1892:238. 15. Budd-Hansen, 1903:74.

SYNONYMS. *August Apple* (7). *EARLY PENNOCK* (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15). *Heicke's Summer Queen* (13). *Harmony* (7, ? of the south, 9). *Indian Queen* (7). *N. J. Red Streak* (7). *Shaker's Yellow* (7, 9). *Sleeper's Yellow* (5). *Warren Pennock* (5, 7).

Fruit large, showy, yellow covered with mixed striped red, but often the yellow predominates. *Flesh* yellow, moderately juicy, subacid, coarse, suitable for culinary use but not esteemed for dessert; season August. *Tree* hardy, a biennial cropper and moderately productive. Not recommended for planting in New York.

Historical. Origin unknown. It was first brought to notice in Ohio more than fifty years ago (1) where it was widely disseminated from some of the nurseries of that state. At one time it was being planted to a limited extent in New York but it has been almost wholly discarded.

EARLY RIPE.

REFERENCES. 1. Warder, 1867:717. 2. Downing, 1869:156. 3. Fitz, 1872:151. 4. *Am. Pom. Soc. Cat.*, 1873. 5. Bailey, *An. Hort.*, 1892:238. 6. *Ill. Sta. Bul.*, 45:320. 1896. 7. *Mich. Sta. Bul.*, 143:200. 1897. 8. Thomas, 1897:634. 9. Alwood, *Va. Sta. Bul.*, 130:121. 1901. 10. *Kan. Sta. Bul.*, 106:52. 1902. 11. Budd-Hansen, 1903:74.

SYNONYMS. None.

Fruit of good medium size, yellowish-green, subacid, good for culinary use. The tree is a good grower, comes into bearing young, and yields full crops in alternate years.

Historical. The locality of its origin is unknown but it is supposed to have originated in Pennsylvania (2).

TREE.

Tree large, vigorous with moderately long, stout branches. *Form* upright spreading, rather dense, top roundish. *Twigs* long, stout, curved; internodes medium. *Bark* brown tinged with olive-green, lightly streaked with scarf-skin; heavily pubescent. *Lenticels* scattering, small to medium, oblong, slightly raised. *Buds* medium size, broad, plump, obtuse, appressed, pubescent.

FRUIT.

Fruit medium to above, fairly uniform in size but not in shape. *Form* roundish oblate somewhat inclined to conic, irregular, broadly ribbed. *Stem* often bracted, medium in length or short, thick. *Cavity* acute or approaching acuminate, usually shallow, rather broad, sometimes russeted. *Calyx* rather small, closed. *Basin* obtuse, usually very shallow, moderately wide, somewhat wrinkled.

Skin light yellowish-green. *Dots* numerous, small, pale gray or russet.

Calyx tube rather narrow, funnel-form. *Stamens* median.

Core medium to rather large, abaxile; cells closed or partly open; core lines clasping. *Carpels* broadly roundish, emarginate. *Seeds* medium size, plump, obtuse.

Flesh white, quite firm, moderately coarse, crisp, tender, juicy, briskly subacid, becoming rather mild subacid when fully ripe, fair to good.

Season August.

EARLY STRAWBERRY.

REFERENCES. 1. Manning, 1838:46. 2. Downings, 1845:73. *fig.* 3. *Horticulturist*, 1:145. 1846. 4. *Mag. Hort.*, 14:488. 1848. *fig.* 5. Thomas, 1849:139. *fig.* 6. Cole, 1849:101. *fig.* 7. Barry, 1851:280. 8. *Am. Pom. Soc. Cat.*, 1852. 9. Elliott, 1854:76. 10. Hooper, 1857:31, 106, 109. 11. Gregg, 1857:36. 12. *Horticulturist*, 14:425. 1859. 13. Warder, 1867:514. *fig.* 14. Downing, 1869:157. 15. Fitz, 1872:143, 151, 160. 16. Leroy, 1873:310. *figs.* 17. Downing, 1881:12 index, app. 18. Wickson, 1889:243. 19. Watts, *Tenn. Sta. Bul.*, 5:78. 1890. 20. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 21. Bailey, *An. Hort.*, 1892:238. 22. Budd-Hansen, 1903:75. *fig.*



EARLY RIPE

SYNONYMS. AMERICAN RED JUNEATING (1). *American Red Juneating* (4, 5, 9, 13, 14, cit. 5, ? 2). *De La Madeleine Rouge* (16). EARLY STRAWBERRY (2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22). *Early Strawberry* (16). *FRAISE* (16). *Louis XVIII* (16). *Red Juneating* (3, 6, 9, 10, 14, 15, 16, err. 4, err. of some American gardens, 2). *St. John Strawberry* (17). *Striped Shropshire* (17). *Tennessee Early Red* (17).

Fruit of a very attractive bright deep red color, very desirable for dessert and good also for culinary uses. The tree is medium in size, a moderate grower, upright when young, but eventually becoming roundish and somewhat spreading. It is hardy, healthy, comes into bearing young and yields moderate to good crops biennially or almost annually. It is not a very satisfactory variety for commercial planting because the demand for it is mostly limited to local markets, a relatively high percentage of the apples are undersized or otherwise unmarketable and the fruit does not keep well. On account of its productiveness and high quality it is a desirable variety for the home orchard.

Historical. Early Strawberry is an American fruit which is said to have originated in the vicinity of New York (2, 3). It was formerly known to some under the name Red Juneating or American Red Juneating. The name Red Juneating has been applied also to the Margaret. In 1846 Downing published the following observations concerning these two varieties (3). "The Early Strawberry has a long stalk, and is a high colored fruit, striped with dark red. The Early Red Margaret has a short stalk and is a dull colored fruit, with faint red stripes. We have had both fruits in bearing this year, and have compared them for several years past. The Early Red Margaret is correctly shown in the beautiful colored plates of Ronald's *Pyrus Malus Brentfordensis*, and in the *Pomological Magazine*. Our Early Strawberry apple is not described in any European work that we have seen. It is greatly superior to the Early Red Margaret in productiveness, and especially in *long keeping* and *ripening gradually*, qualities that are rare in early apples and for which the market dealers in New York rate the Strawberry very highly."

Early Strawberry has been extensively disseminated and is generally catalogued by nurserymen throughout the apple-growing regions of America (21).

Fruit.

Fruit below medium to medium, pretty uniform in shape and size. *Form* roundish conic or roundish, regular or somewhat ribbed; sides often unequal. *Stem* long and rather slender, often clubbed. *Cavity* acute or approaching acuminate, deep, broad, symmetrical, sometimes with faint radiating rays of russet. *Calyx* rather small, closed or sometimes open; lobes long, narrow. *Basin* small, shallow to moderately deep, narrow, obtuse, slightly furrowed.

Skin rather thick, tough, smooth, waxy, entirely red or yellow nearly covered with a rich dark red, mottled and irregularly striped and splashed with deeper red. *Dots* minute, grayish.

Calyx tube short, moderately wide, conical or approaching funnel-shape, with fleshy pistil point projecting into the base. *Stamens* median.

Core large, axile or somewhat abaxile; cells usually open, sometimes partly closed; core lines nearly meeting. *Carpels* broadly roundish to elliptical, much concave, emarginate. *Seeds* medium or above, wide, plump, obtuse, dark brown.

Flesh whitish-yellow often with streaks of red, moderately coarse, crisp, moderately tender, juicy, subacid, aromatic, sprightly, very good.

Season August.

EGG TOP.

REFERENCES. 1. *Mag. Hort.*, 10:210. 1844. 2. Elliott, 1854:169. 3. Hooper, 1857:34. 4. Warder, 1867:717. 5. Downing, 1869:159. 6. Thomas, 1875:498.

SYNONYMS. *Early June* (3). *EGG TOP* (1, 2, 4, 5, 6). *Eggtop* (3). *Eve APPLE* (3). *Eve* (2, 5). *Round Top* (2, 5). *Sheepnose* (2, 5). *Wine* of some (2, 5).

Fruit similar in shape to Black Gilliflower but not so large, somewhat streaked and shaded with red, pleasant flavored but not high in quality; good for dessert but not for cooking. Tree large, moderately vigorous and a regular and abundant bearer. In season from late fall to midwinter.

Historical. An old variety of uncertain origin. A few trees of it are occasionally found in the oldest orchards but it is now nearly obsolete.

ELGIN PIPPIN.

REFERENCES. 1. Bailey, *An. Hort.*, 1892:238. 2. Clayton, *Ala. Sta. Bul.*, 47:6. 1893. 3. Thomas, 1897:634. 4. Beach and Clark, *N. Y. Sta. Bul.*, 248:118. 1904.

SYNONYMS. None.

Fruit of good size and attractive appearance for a yellow apple. It evidently belongs to the Fall Pippin class but it does not closely resemble that variety and is not superior to it. Although it has much merit it does not excel other varieties of its season and is not recommended for planting in New York. The tree is a strong grower and productive.

Historical. Origin Alabama (2, 3). Downing questions whether it is identical with the White Spanish Reinette but it appears to be distinct from that variety.¹ Warder describes another variety under the name Elgin Pippin which we have not seen.²

TREE.

Tree vigorous, upright; branches long, moderately stout. *Form* very much spreading and somewhat drooping, rather dense. *Twigs* medium in

¹ Downing, 1869:404.

² Warder, 1867:717.

APPLES
EARLY STRAWBERRY



size, curved, stout; internodes long. *Bark* brown, tinged with olive-green, mottled with scarf-skin, slightly pubescent. *Lenticels* scattering, large, oblong, raised. *Buds* large, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit above medium to large, averages nearly large. *Form* oblate to roundish conic, distinctly ribbed, irregular; sides often unequal, sometimes elliptical, not uniform. *Stem* medium to very long, slender. *Cavity* acute to acuminate, medium in depth to deep, rather broad to narrow, usually with outspreading russet rays, sometimes decidedly compressed, sometimes lipped. *Calyx* medium to large, closed; lobes often leafy, long, acuminate. *Basin* sometimes oblique, shallow to moderately deep, moderately wide to rather narrow, obtuse to rather abrupt, often prominently and irregularly ribbed.

Skin thin, moderately tender, greenish yellow becoming deep yellow, sometimes with a faint bronze blush in the sun. *Dots* scattering, russet, or submerged and whitish.

Calyx tube rather large, moderately wide, conical or approaching funnel-form. *Stamens* median to basal.

Core abaxile, medium to large; cells unsymmetrical, usually wide open; core lines meeting or slightly clasping. *Carpels* irregular, often somewhat ovate, much concave, tufted. *Seeds* often abortive, dark, rather large, long, flat, acute to acuminate.

Flesh whitish tinged with yellow, firm, moderately fine, crisp, tender, juicy, subacid, good.

Season September to December. Commercial limit, October.

ENGLISH PIPPIN.

REFERENCES. 1. Gibb, *Am. Pom. Soc. Rpt.*, 1887:50. 2. Troop, *Ind. Sta. Bul.*, 53:123. 1894. 3. Budd, *Ia. Sta. Bul.*, 31:328. 1895. 4. Beach, *W. N. Y. Hort. Soc. Rpt.*, 1896:51. 5. Troop, *Ind. Sta. An. Rpt.*, 12:79. 1899. 6. Budd, *Ia. Sta. Bul.*, 41:85. 1899. 7. Beach and Clark, *N. Y. Sta. Bul.*, 248:118. 1904.

SYNONYMS. ENGLISH PIPPIN (1, 2, 3, 4, 5, 6, 7). *Englischer Pepping* (1). *No. 587* (1). *Pepping Englischer* (1).

A Russian apple inferior in quality to standard varieties of its season. Not recommended for planting in New York.

FRUIT.

Fruit large. *Form* variable, roundish oblate to oblate conic or inclined to oblong truncate, ribbed; sides sometimes unequal. *Stem* short, usually not exerted. *Cavity* acuminate, rather wide, moderately deep, russeted and with outspreading russet. *Calyx* medium size, closed; lobes usually erect or connivent. *Basin* medium in width to rather narrow, deep, abrupt, irregularly furrowed.

Skin yellow or greenish, sometimes with shade of brownish-red overspread with thin white bloom. *Dots* minute, pale or russet.

Calyx tube cone-shape or approaching funnel-form. *Stamens* median.

Core large, somewhat abaxile; cells partly open; core lines meeting. *Seeds* medium size, obtuse.

Flesh tinged with yellow, moderately juicy, fine-grained, sprightly subacid, fair to good in quality.

Season late September to November.

ENGLISH SWEET.

Ramsdell or Ramsdell Sweet has been described by some leading pomologists under the name English Sweet. Ramsdell *Sweet* is the name accepted for this variety in the catalogue of the American Pomological Society, and it is generally known among nurserymen and fruit growers as Ramsdell or Ramsdell Sweet. For a description of this variety, the reader is referred to Ramsdell Sweet, page 175.

FALL GREENING.

REFERENCES. 1. Warder, 1867:718. 2. Downing, 1869:167.

SYNONYMS. None.

This variety originated at Claverack, Columbia county, N. Y. (1). According to Downing the tree is moderately vigorous and very productive. The fruit medium size, greenish-yellow, subacid, good to very good. In season from December to February.

Downing also mentions another Fall Greening of similar color, sprightly subacid, in season from September to November; tree a poor grower.

We have not seen either of these varieties.

FALL HARVEY.

REFERENCES. 1. Manning, 1838:48. 2. Downing, 1845:84. 3. Cole, 1849:117. 4. Hovey, *Mag. Hort.*, 15:537. 1849. *fig.* 5. Thomas, 1849:155. 6. Emmons, *Nat. Hist. N. Y.*, 3:47. 1851. *col. pl.* No. 74 and *fig.* 7. Barry, 1851:284. 8. Elliott, 1854:132. 9. Gregg, 1857:42. 10. Warder, 1867:482. 11. *Am. Pom. Soc. Cat.*, 1869. 12. Hoskins, *Amer. Gard.*, 15:272. 1894. 13. Wagh, *Vt. Sta. An. Rpt.*, 14:296. 1901. 14. Budd-Hansen, 1903:79.

SYNONYMS. FALL HARVEY (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14). *Fall Harvey* (13). *Fall Pippin* erroneously (13). HARVEY (13).

Fruit large, yellow, showy, similar in general appearance to Fall Pippin but less desirable where that variety can be grown. According to Hoskins (12) it is hardy in Northern Vermont and has there proved an excellent annual bearer. He considers it valuable for all Northern New England and Canada. Cole (3) observes that the fruit is "fine and fair but not first-rate, and rather apt to fall, or to rot on the tree." Elliott (8) says that it is not productive but Hovey (4) and Cole (3) call it a vigorous grower and a good bearer. In the nursery it may be easily known from Fall Pippin by its yellow and more slender shoots (4).

FRUIT (2, 3, 4, 10).

Fruit large. *Form* roundish, slightly oblate, regular or somewhat ribbed at the base; crown large, slightly oblique. *Stem* short, stout or rather slen-

der. *Cavity* rather shallow to deep, wide, uneven. *Calyx* small to large, closed. *Basin* narrow to rather wide, shallow, wrinkled.

Skin smooth, pale yellow to deep yellow, sometimes blushed. *Dots* small, gray or russet.

Core large, abaxile. *Seeds* medium size, acute, short, plump.

Flesh whitish or tinged with yellow, a little coarse, crisp, juicy, rich sub-acid, with high flavor, very good quality.

Season October to December.

FALL JENNETING.

REFERENCES. 1. *Mag. Hort.*, 12:474. 1846. 2. Hovey, *Ib.*, 13:161. 1847. fig. 3. Elliott, 1854:132. 4. Downing, 1857:213. 5. *Am. Pom. Soc. Cat.*, 1862. 6. Warder, 1867:533. fig. 7. Thomas, 1875:498. 8. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 9. Bailey, *Am. Hort.*, 1892:239. 10. Budd-Hansen, 1903:80.

SYNONYMS. FALL GENETING (6). *Fall Gennetting* (3). FALL JENETTING (8). FALL JENNETING- (1, 2, 4, 5, 7, 10). FALL JENNETTING (3, 9).

Fruit often large but variable in size, a considerable portion of the crop being undersized or otherwise unmarketable. It is of a green or yellowish color, shows bruises quite readily, does not stand shipping very well and is not a good keeper. It is very good for culinary uses and acceptable for dessert; being of light weight, it is less desirable than some other varieties for evaporating. The tree is an exceedingly strong grower and long-lived, eventually becoming very large, tall and spreading. It is hardy, healthy, comes into bearing rather young and is a reliable cropper, yielding moderately good to heavy crops biennially or nearly annually. There is apt to be considerable loss from premature dropping of the fruit. Not recommended for planting in New York.

Historical. It appears that this variety was brought into Western New York from Athens on the Hudson, about 1827 (1, 2). Elliott (3) in 1854 called it an old Connecticut variety. It has been grown more largely in portions of the Hudson valley and of Western New York than it has in other sections of the state but it has not been cultivated extensively in any locality. It is still listed by nurserymen (9) but in New York it is now seldom planted and is gradually going out of cultivation.

TREE.

Tree large, vigorous or very vigorous. *Form* spreading or somewhat roundish. *Twigs* moderately long, curved, moderately stout; internodes medium. *Bark* dark brown lightly streaked with gray scarf-skin; slightly pubescent. *Lenticels* scattering, medium size, oblong, slightly raised. *Buds* medium size, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit above medium to large. *Form* roundish oblate inclined to conic, slightly ribbed at the base; sides unequal. *Stem* medium to short, moderately thick. *Cavity* acuminate, deep, wide, rather symmetrical, with outspreading rays of russet. *Calyx* below medium to rather large, closed or somewhat open; lobes long, narrow, acute, reflexed. *Basin* rather small, shallow, narrow, somewhat furrowed and wrinkled.

Skin thin, tough, smooth, pale greenish-yellow with faint brownish-red or bronze bluish. *Dots* moderately numerous, rather inconspicuous, sometimes russet but more often whitish and submerged. *Prevailing effect* yellow.

Calyx tube rather long, narrow funnel-shape. *Stamens* median to basal.

Core small, axile to somewhat abaxile; cells symmetrical, closed; core lines clasping. *Carpels* somewhat roundish to broadly ovate. *Seeds* light brown, rather small, moderately narrow, plump, acute.

Flesh tinged with yellow, moderately firm, fine, crisp, tender, juicy, sprightly, subacid, good.


Season late September to December.

FALL ORANGE.

REFERENCES. 1. Thomas, *Cultivator*, 5:246. 1848. 2. *Ib.*, 1849:155. 3. Elliott, 1854:149. 4. Gregg, 1857:42. 5. Downing, 1857:143. 6. *Am. Pom. Soc. Cat.*, 1862. 7. Warder, 1867:718. 8. Downing, 1869:168. 9. *Ib.*, 1872:10 index, app. 10. Bailey, *Am. Hort.*, 1892:239. 11. Burrill and McCluer, *Ill. Sta. Bul.*, 45:321. 1896. 12. Budd-Hansen, 1903:80. fig.

SYNONYMS. FALL ORANGE (1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12). *Fall Orange* (3). *Hogpen* (8). *Holden* (5, 8). *Holden Pippin* (8). *Hoypen* (5). *Jones' Pippin* (5, 8). *Long Island* (8). *Long Island Graft* (9). *N. Y. Bellflower* (9). ORANGE (3). *Orange* (8). *Red Check* (8). *Speckled* (9). *Westbrook* (9). *White Graft* of Wis. (8). *White Newell* (9).

When well grown Fall Orange is of good size, yellow or greenish, with occasionally a shade of red, agreeable subacid, and very good in quality for culinary use. When it becomes fully ripe so that its acidity is subdued it is an excellent dessert apple. The tree is thrifty, hardy, long-lived and a regular biennial cropper, often yielding pretty heavy crops. It is not generally regarded as a good commercial variety because its color is yellow, the fruit is rather tender and a poor shipper, and with heavy crops there is apt to be a comparatively large percentage of fruit that is undersized or otherwise unmarketable. It is in season from late September to early winter; sometimes a portion of the fruit is kept till spring.



FALL ORANGE

It resembles Autumn Swaar considerably in general appearance; for a comparison of the two varieties the reader is referred to the description of Autumn Swaar, pages 11, 12.

Historical. Fall Orange was described by Thomas in 1848 as a new or newly-introduced variety of unknown origin (1). In 1857 Downing gave its origin as Holden, Mass. (5). Since its introduction it has been sparingly disseminated in various parts of the state. Although it is still listed by some nurserymen (10) it is now seldom planted in New York.

FRUIT.

Fruit above medium to large; fairly uniform in size but not in shape. *Form* roundish conic, irregular. *Stem* medium to rather short, slender. *Cavity* acute to acuminate, deep, medium in width, regular or slightly compressed, often russeted and with outspreading russet rays. *Calyx* medium to large, open or sometimes nearly closed. *Basin* uneven, one side projecting higher than the other, moderately deep to deep, narrow to moderately wide, abrupt, furrowed.

Skin pale yellow or greenish, sometimes with brownish blush. *Dots* numerous, large and small, russet or sometimes reddish, areolar.

Calyx tube rather large, usually long, conical to funnel-form with fleshy pistil point projecting into the base; the lower part of the funnel cylinder is sometimes enlarged. *Stamens* median or below.

Core medium to rather small, axile; cells symmetrical; core lines meeting when the tube is short, clasping when it is long. *Carpels* elliptical to nearly cordate. *Seeds* not numerous, rather dark brown, medium to below, plump, obtuse.

Flesh white, moderately fine, crisp, rather tender, juicy, subacid, aromatic, very good.

Season late September to early winter.

FALL PIPPIN.

REFERENCES. 1. *Amer. Gard. Cal.*, 1806:585. 2. Coxe, 1817:109. fig. 3. Cobbett, 1821:par. 300. 4. Thacher, 1822:120. 5. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 6. Fessenden, 1828:131. 7. Kenrick, 1832:32. 8. Floy-Lindley, 1833:85. 9. *Mag. Hort.*, 1:326, 364. 1835. 10. Downing, 1845:84. 11. Floy-Lindley, 1846:412 app. 12. Phoenix, *Horticulturist*, 1:361. 1847. 13. Downing, *Ib.*, 3:345. 1849. Cole, 1849:117. fig. 15. Thomas, 1849:155. fig. 16. Phoenix, *Horticulturist*, 4:472. 1850. 17. Barry, 1851:283. 18. Emmons, *Nat. Hist. N. Y.*, 3:44. 1851. fig., col. pl. No. 18. 19. *Am. Pom. Soc. Cat.*, 1852. 20. Elliott, 1854:78. fig. 21. Gregg, 1857:42. 22. Hooper, 1857:34, 106, 110. 23. Warder, 1867:571. fig. 24. Downing, 1869:169. fig. 25. Todd, 1871:311. fig. 26. Fitz, 1872:143, 149, 162. 27. Hogg, 1884:74. 28. Wickson, 1889:244. 29. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 30. Bailey, *Am. Hort.*, 1892:239. 31. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 8:7. 1901. figs. 32. Budd-Hansen, 1903:80. fig. 33. Beach and Clark, *N. Y. Sta. Bul.*, 248:119. 1904.


SYNONYMS. *American Fall* (7). *AUTUMN PIPPIN* (4). *Cathead* incorrectly (24). *Cat Head* (25). *Cobbett's Fall* (7). *Cobbett's Fall Pippin* (27). *Concombre Ancien* (7). *De Rateau* (7). *D'Espange* (7). *Episcopal* (24, 25). *FALL PIPPIN* (1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33). *Fall Pippin* (4). *Golden Pippin* (22, 25, erroneously 2, 24). *Holland Pippin* (2, 5, err. 15). *Philadelphia Pippin* (20, 24, 25). *Pound Pippin* (24, 25). *Pound Royal* (25, of some 24). *Prince's large Pippin of N. Y.* (2). *Reünete Blanche d'Espagne* (7). *Summer Pippin* (2). *Van Duzyn's Pippin* (2). *Van Dyn's Pippin* (5). *York Pippin* (24, 25).

Fruit large, and, when fully ripe, of an attractive yellow color. The flesh is tender, rich and very good in quality, being excellent for dessert but especially desirable for culinary uses. The tree is a strong grower, hardy and very long-lived, eventually becoming large or very large. Since the foliage and fruit are both quite subject to the attacks of the apple-scab fungus, thorough preventive treatment for this disease is necessary in order to grow Fall Pippin successfully for commercial purposes.¹ The crop does not ripen uniformly, some of the fruit being ripe, well colored and ready for immediate use in September, while at the same time a considerable portion of the crop is still hard and green. When grown under favorable conditions and properly handled some portion of the crop may keep till midwinter or later, but even carefully selected fruit cannot be relied upon to hold in common storage till December 1st without considerable loss. In cold storage it may be held till January or February (33). Fall Pippin is generally in pretty good demand in local markets, and in portions of Eastern New York it is being used to a limited extent for the early export trade. It is one of the most desirable varieties of its season for the home orchard.

HOLLAND PIPPIN AND FALL PIPPIN COMPARED.

From the time of Coxé (2), Fall Pippin has by some been called Holland Pippin. The Holland Pippin indeed much resembles Fall Pippin, but it differs from it in being in season from mid-August

¹ Directions for treating apple scab are given in the reports of this Station for 1899:397-418, and for 1903:321-386.



FALL PIPPIN (Reduced Size)

to midautumn and in being more roundish and less flattened, and in having a short, thick stem which is not exerted.

Historical. Origin unknown. Downing (10) held the opinion that Fall Pippin is an American variety and probably a seedling raised in this country from either the White Spanish Reinette or the Holland Pippin, both of which it resembles. It has been widely disseminated from the Atlantic to the Pacific in most of the important apple-growing regions of the continent and is still listed by many nurserymen (30). We are unable to determine how long this variety has been in cultivation. In some New York orchards trees of it are found which are more than one hundred years old. It is quite generally grown for home use throughout New York, and also to a limited extent for market, but it is now seldom planted.

TREE.

Tree large, moderately vigorous to very vigorous, with large, long branches which eventually become somewhat drooping. *Form* spreading or roundish and rather dense. *Twigs* moderately long, curved, stout and with large terminal buds; internodes medium. *Bark* dark reddish-brown, somewhat tinged with green, heavily coated with gray scarf-skin; much pubescent. *Lenticels* scattering, medium size, oval, raised. *Buds* medium size, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit large or very large, pretty uniform in size and shape. *Form* roundish to roundish oblate, sometimes a little inclined to conic, or sometimes slightly oblong and truncate, often obscurely ribbed. *Stem* medium to rather long, thick. *Cavity* acute or approaching acuminate, moderately deep, moderately narrow to rather wide, symmetrical or sometimes compressed, russeted. *Calyx* medium to large, open; lobes separated at the base, moderately long, rather narrow, acuminate. *Basin* medium in depth to deep, moderately narrow to rather wide, abrupt, wavy, slightly wrinkled.

Skin, thin, smooth, at first greenish-yellow but becoming clear yellow, sometimes faintly blushed. *Dots* numerous, small, pale and submerged or russet.

Calyx tube large, wide, long, conical to nearly funnel-form. *Stamens* median to basal.

Core medium size, somewhat abaxile; cells symmetrical, closed or partly open; core lines meeting or clasping. *Carpels* roundish, emarginate, tufted. *Seeds* rather dark brown, medium size, somewhat acute, plump.

Flesh whitish or tinged with yellow, moderately firm, rather fine, tender, very juicy, agreeable subacid, somewhat aromatic, very good.

Season late September to January.

FALL WINE.

REFERENCES. 1. *Mag. Hort.*, 14:114. 1848. 2. *Cole*, 1849:108. 3. *Ib.*, 1849:109. 4. *Elliott*, 1854:77. 5. *Horticulturist*, 10:87. 1855. 6. *Hooper*,

1857:36. 7. Downing, 1857:78. 8. *Am. Pom. Soc. Rpt.*, 1860:50, 240. 9. Warder, 1867:434. *fig.* 10. Downing, 1869:170. *fig.* 11. Fittz, 1872:153. 12. Thomas, 1875:200. 13. Barry, 1883:338. 14. Wickson, 1889:244. 15. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 16. Bailey, *An. Hort.*, 1892:239. 17. Budd-Hansen, 1903:81. *fig.*

SYNONYMS. FALL WINE (1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17). *House* (10). *Hower* (10). *Musk Spice* (10). *Ohio Wine* (7, 10, 13). *Sharpe's Spice* (7, 10). *Sweet Wine* (7, 10). *Uncle Sam's Best* (7, 10). *WINE* (2). *Wine of Cole* (4).

Fruit medium or above, bright red, attractive; in season from September to early winter. The tree is of medium size with rather drooping branches, moderately vigorous, healthy, moderately long-lived and yields good to heavy crops biennially. The fruit is apt to drop to a considerable extent before it is fully mature. It is very tender, not a good shipper and not a desirable commercial variety. It is not much valued for culinary use but it is generally esteemed wherever it is known on account of its excellent dessert qualities. It is not recommended for commercial planting in New York.

Fall Wine should not be confused with Twenty Ounce which in some sections of New York is known under the name of Wine apple.¹ It is also quite distinct from the true Wine which is a large, showy apple that ripens in midautumn and often keeps well through the winter.²

Historical. The origin of this variety is unknown. Elliott (4) says that it was introduced into the West from the garden of Judge Jonathan Buel, Albany, N. Y., about 1832. It has been a favorite variety in many parts of the Middle West but has not been extensively cultivated in New York and is now seldom or never planted in this state.

FRUIT.

Fruit medium or above. *Form* roundish oblate, somewhat ribbed; sides often unequal. *Stem* medium to rather long. *Cavity* obtuse to somewhat acute, wide, shallow to moderately deep, sometimes lipped. *Calyx* small to medium, closed or partly open; lobes long, narrow, reflexed. *Basin* deep, wide or medium in width, rather abrupt, furrowed.

Skin clear yellow washed with red which on the exposed cheek deepens to a beautiful bright blush, indistinctly striped with carmine. *Dots* yellowish-brown or russet.

Calyx tube long, narrow, funnel-form. *Stamens* median.

Core medium, axile; cells symmetrical, closed or slightly open; core lines clasping. *Carpels* broadly roundish, tufted. *Seeds* medium size, rather wide, acute.

Flesh tinged with yellow, tender, juicy, aromatic, very mild subacid or almost sweet; very good for dessert but only fair for culinary use.

Season September to January.

¹ See description of Twenty Ounce, page 227.

² See description of Wine apple in Volume I of this Report, page 373.

FAMEUSE.

REFERENCES. 1. Forsyth, 1824:101. 2. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 3. *London Hort. Soc. Cat.*, 1831:No. 757. 4. Kenrick, 1832:33. 5. Floy-Lindley, 1833:16. 6. Manning, 1838:55. 7. *Ib.*, *Mag. Hort.*, 7:45. 1841. 8. Downing, 1845:91. 9. *Horticulturist*, 1:257. 1846. 10. Thomas, *Cultivator*, 5:306. 1848. *fig.* 11. *Ib.*, 1849:147. *fig.* 12. Cole, 1849:118. 13. Emmons, *Nat. Hist. N. Y.*, 3:36. 1851. *col. pl.* No. 48. 14. Hovey, 2:7. 1851. *col. pl.* and *fig.* 15. *Mag. Hort.*, 17:15. 1851. *fig.* 16. Goodrich, *Ib.*, 17:122. 1851. 17. *Am. Pom. Soc. Cat.*, 1852. 18. Elliott, 1854:77. 19. Gregg, 1857:42. 20. Hooper, 1857:36, 107, 110. 21. Lothrop, *Mag. Hort.*, 32:363. 1866. 22. Warder, 1867:505. 23. Todd, 1871:25. *fig.* 24. Fitz, 1872:145. 25. Leroy, 1873:483. 26. *Montreal Hort. Soc. Rpt.*, 1876:12. 27. Jack, *Ib.*, 6:61. 1880. 28. *Ib.*, 8:140. 1881-82. 29. Hogg, 1884:58. 30. Hoskins, *Rural N. Y.*, 47:249. 1888. 31. Wickson, 1889:245. 32. *Montreal Hort. Soc. Rpt.*, 14:88, 89. 1889. 33. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290. 34. *Montreal Hort. Soc. Rpt.*, 15:19, 26. 1890. 35. Bailey, *An. Hort.*, 1892:239. 36. *Ib.*, 1892:249. 37. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57. 38. *Amer. Gard.*, 14:425. 1893. 39. Burrill and McCluer, *Ill. Sta. Bul.*, 45:321. 1896. 40. *Can. Hort.*, 20:304. 1897. *figs.* 41. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 5:12. 1898. *figs.* 42. *Can. Hort.*, 22:226. 1899. 43. Craig, *Amer. Gard.*, 20:27. 1899. 44. *Vt. Sta. Bul.*, 83:83-92. 1900. 45. *Can. Hort.*, 24:236. 1901. 46. Waugh, *Vt. Sta. An. Rpt.*, 14:293. 1901. 47. Budd-Hansen, 1903:82. *fig.* 48. Beach and Clark, *N. Y. Sta. Bul.*, 248:120. 1904. 49. Scriber, *Can. Hort.*, 28:277. 1905.

SYNONYMS. *Chimney Apple* (14, 16, 22, 25). *DE NEIGE* (3, 25, 29). *De Neige* (14, 15). *du Marechal* (25). *FAMEUSE* (1, 2, 4, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 28, 30, 31, 32, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48). *Fameuse* (5, 8, 25, 29, 36). *La Belle Fameuse* (43). *La Fameuse* (29). *Neige* (2). *Neige-Framboise de Gielen* (25). *POMME DE NEIGE* (5, 8). *Pomme de Neige* (4, 9, 10, 11, 13, 14, 15, 16, 18, 20, 22, 23, 41). *Pomme de Niège* (1, 7, 12). *Pomme Fameuse* (3). *Pomme de Fameuse* (14, 15). *Sanguineus* (8, 14, 15, 18, 23, 25). *SNOW* (36). *Snow* (2, 10, 12, 14, 16, 18, 20, 21, 22, 23, 25, 31, 33, 41, 43, 46, 47, 48).


Fameuse is one of the most desirable dessert apples of its season. It is very beautiful in appearance and the flesh is white, tender and excellent in flavor and quality for dessert. It is decidedly inferior to other varieties of its season for culinary purposes. It is well known in market, and during its season, which extends from October to the holidays, it usually sells above average market prices, particularly if well colored and free from scab or other imperfections. The fruit is often badly injured by the apple-scab fungus, but this may readily be controlled by proper preventive treatment.¹ It keeps well in cold storage. Some report that if free

¹ *N. Y. Sta. An. Rpt.*, 18:399-418. 1899. *Ib.*, 22:321-386. 1903.

from scab it may be held as long as Rhode Island *Greening* (48). In the Champlain district and in portions of the St. Lawrence valley it is one of the most important varieties found in commercial orchards. Generally speaking, it grows to a higher degree of perfection in those districts than it does in other apple-growing regions of New York. In the more southern sections of the state it appears to succeed best in the high elevations and on light well drained soil with clay subsoil. The tree is of medium size, a moderate grower, hardy, healthy, rather long-lived, and a reliable cropper yielding good to heavy crops biennially or sometimes nearly annually. The fruit hangs pretty well to the tree. Fruit of marketable grades is fairly uniform in size, but there is a considerable amount of it that is too small for market. In those portions of New York south of Lake Champlain and the St. Lawrence valley the fruit does not usually develop its best color, and in some seasons, and particularly in unfavorable localities, it is so poorly colored as to be quite unsatisfactory. In such locations it is better for the fruit grower to plant McIntosh instead of Fameuse if he desires to grow a variety of the Fameuse group.

Historical. Waugh (44) gives the following historical account of the Fameuse. "The history of Fameuse is obscure, probably beyond clearing up satisfactorily, but extremely interesting as far as we know or can guess at it. The turning point of speculation for years has been as to whether the variety is of American or European origin. One of the most interesting contributions to this discussion was made by Mr. Chauncey Goodrich, (16) of Burlington, Vt., in 1851. We quote the following extracts from this article:

'It is here one of the most common as well as oldest varieties; hundreds of barrels are sold in a single season in this town alone. . . . All American writers call it a Canadian apple; of this I think there is no proof. One hundred and twenty years since, the French planted this variety on the eastern shore of Lake Champlain, opposite Fort Frederick on Crown Point, at a place called 'Chimney Point'—more than fifty years before any other permanent settlement. From these old trees cions have been scattered through Vermont, and called the Chimney apple. A very intelligent and highly educated French seigneur residing on an old seignory eighty miles below Quebec informed me that this was one of the first varieties of apples planted on the place; that the trees were very old and were brought from France. The early French settlers planted the same variety at Ogdensburg, Detroit, and other places on Lakes Erie and Ontario, where it is still known as the 'Snow Apple'; also at Kaskaskia, Illinois, more than one hundred



FAMEUSE

and fifty years since, where the old trees are still productive, and apples from them are sent to St. Louis, &c. The same apple may be found in France, and in London of the growth of France.'

"It is hardly to be supposed that a seedling apple was produced in Canada at so early a day as to be distributed more than a thousand miles in every settlement made by the French, one hundred and fifty years since.

"Another fact tending to suggest a European origin for Fameuse is that it is usually found in the old gardens, in company with well known European varieties of pears, apples and other fruits.

"On the other hand the testimony of European pomologists is mostly against the theory of a European origin. The variety is known in the larger collections of all the countries of Europe, just as Ben Davis is, and has been known there for many years. But most European authors unhesitatingly assign a Canadian origin to the variety; and the variety seems too little known, too little appreciated, and too little at home with European surroundings for us to believe it originated there. Those who call it a European apple usually assign its nativity to France; but Leroy (25) the greatest of all French, and perhaps of all European authorities, did not know the variety. He says that Le Lectier cultivated the Pomme de Neige (synonym of Fameuse) at Orleans (France) before 1628; but Leroy did not know whether or not this was the same Pomme de Neige grown in Canada. In fact he says, 'I have never, up to the present time, met this apple Pomme de Neige on our soil. In place of it they have always sent me Calvill de Neige, ripening from January to March.' Most of the so-called Snow apples of Europe, in fact, are white skinned and totally different from the Snow, or Fameuse, of America.

"It is agreed that, whether the Fameuse came from Europe or not, it was distributed by the earliest of the French missionaries and planted by the first settlers. Quebec was founded shortly before 1600 and Montreal in 1641. The *seigniorie du Cote de Beaupre*, said to be the oldest seigniorie in Quebec, was granted in 1636 and promptly colonized. Thus we have almost a hundred years of French settlement and missionary activity prior to 1700, the approximate date at which, according to Mr. Goodrich, the Fameuse was brought to Vermont. This seems to allow ample time for a Canadian origin for the variety and for its wide distribution in Quebec, Ontario and the Northern states.

"The early distribution of apples, either from Europe to Canada, or from place to place on this continent, was accomplished chiefly, almost exclusively, by seeds. Some of the missionaries knew the art of grafting, but there was small encouragement to practice it. From these considerations, and others which cannot be fully argued here, the writer is firmly convinced that the Fameuse originated in Canada from seed brought from France."

TREE.

Tree vigorous, with long, moderately stout branches. *Form* upright spreading or roundish, rather dense. *Twigs* medium size, curved, stout; internodes short. *Bark* dark brown tinged with red, lightly coated with scarf-skin; pubescent. *Lenticels* scattering, small, round or somewhat oval, slightly raised. *Buds* medium size, flat, obtuse, free, pubescent.

FRUIT.

Fruit hardly averages medium but sometimes is above medium size. *Form* roundish inclined to conic, sometimes a little oblate, regular, uniform, symmetrical. *Stem* medium to short but sometimes long, rather slender. *Cavity* acute to somewhat acuminate, moderately deep to deep, rather wide, often gently furrowed, sometimes partly russeted but generally smooth and red or greenish. *Calyx* rather small, usually closed. *Basin* medium in width and depth, rather small, somewhat abrupt, obscurely furrowed or wrinkled, often having mammiform protuberances.

Skin thin, tender, smooth, light bright red deepening to almost purplish black in highly colored specimens with a somewhat striped appearance toward the apex. In less highly colored specimens the striped effect is more noticeable. *Dots* few, scattering, light.

Calyx tube narrow, funnel-form. *Stamens* median or somewhat basal.

Core medium to rather small, axile; cells closed; core lines clasping. *Carpels* symmetrical, roundish or inclined to elliptical, somewhat emarginate, mucronate. *Seeds* dark, long, rather narrow, acute to acuminate.

Flesh white, sometimes streaked or stained with red, very tender, juicy, subacid becoming very mild subacid or sweetish, aromatic, very good for dessert.

Season October to midwinter.

STRIPED FAMEUSE. A variety has sometimes been propagated and disseminated under the name Fameuse which is recognized as Striped Fameuse. The tree is a thriftier grower in the nursery than the true Fameuse, but the fruit is less desirable, being inferior in color but similar in all other respects to Fameuse. It is mottled or thinly washed with bright red over a pale yellow background, striped and splashed with carmine.

OTHER VARIETIES OF THE FAMEUSE GROUP.

Waugh (44) remarks that one of the striking things about the Fameuse type is that it has the strong tendency to reproduce itself from seed. This has been taken advantage of in the last fifty years, and apples of the Fameuse type have been grown from seed by the hundred and planted in the orchard. This practice has prevailed largely in Quebec in neighborhoods where nurseries were scarce and grafted nursery trees expensive or unknown. He concludes, therefore, that the modern Fameuse apples are most certainly not all from the same original seed, the conspicuous variations among them being thus accounted for at least in part. He further observes that seedlings of the Fameuse often show so much



FANNY

departure from the common characters of Fameuse as to be readily recognized as something different. Such seedlings are generally accepted as new varieties, and in cases where they show conspicuous merit they are separately propagated by grafting, and eventually receive special names of their own. He then lists several named varieties of the Fameuse group, including Bloom, Brilliant, Canada Baldwin, Fameuse Green, Fameuse Noire, Fameuse Sucre, La Victoire, Louise, McIntosh, Hilaire and Shiawassee. The more important of these are described under their respective names in this volume.

FANNY.

REFERENCES. 1. Downing, 1869:173. *fig.* 2. Thomas, 1875:499. 3. *Am. Pom. Soc. Cat.*, 1877:8. 4. Barry, 1883:333. 5. *Am. Pom. Soc. Rpt.*, 1887:93. 6. Bailey, *Am. Hort.*, 1892:239. 7. *Rural N. Y.*, 55:642, 706, 707. 1896. *fig.* 8. *Ib.*, 56:292. 9. Budd-Hansen, 1903:83. *fig.*

SYNONYMS. None.

This is an attractive bright red fruit of good dessert quality. It begins to ripen about the first of September, and continues in season till late fall. The tree is quite vigorous, comes into bearing moderately early and is a reliable annual cropper. Some regard it as desirable for commercial planting (7, 8), but it has not proved so at this Station because it is somewhat deficient in size. On account of its beauty and excellent dessert quality it is worthy of being classed among the varieties desirable for the home orchard.

Historical. Originated with Dr. John K. Eshelman, Lancaster Co., Pa. (7). It has been as yet but little disseminated in New York.

TREE.

Tree moderately vigorous with moderately long, stout branches. *Form* flat, spreading, open. *Twigs* long, curved, moderately stout; internodes long. *Bark* brown tingled with olive-green, heavily streaked with scarf-skin; much pubescent. *Lenticels* quite numerous, medium size, round, not raised. *Buds* medium size, plump, obtuse, free, heavily pubescent.

FRUIT.

Fruit above medium to below medium size, pretty uniform in size and shape. *Form* roundish, slightly oblate varying to somewhat oblong or ovate,

regular or slightly ribbed. *Stem* short to medium, slender. *Cavity* acute to nearly obtuse, medium in width and depth, sometimes russeted. *Calyx* small to medium, closed or partly open; lobes rather short, narrow, acute. *Basin* shallow to medium in depth, moderately wide, rather abrupt, usually furrowed.

Skin thin, tender, smooth, clear yellow mostly overlaid with bright red indistinctly striped with carmine. *Dots* small, yellowish.

Calyx tube rather wide, slightly funnel-form to conical with pistil point projecting into the base. *Stamens* median to marginal.

Core below medium, somewhat abaxile; cells open; core lines clasping. *Carpels* broadly ovate to elliptical, slightly emarginate. *Seeds* medium to large, moderately wide, flat, plump, acute.

Flesh whitish slightly tinged with yellow, moderately firm, fine, very tender, juicy, mild subacid, good to very good.

Season September to November or later.

FISHKILL.

REFERENCES. 1. Downing, 1869:176. 2. *N. Y. Sta. An. Rpt.*, 11:222. 1892. 3. Waugh, *1st. Sta. An. Rpt.*, 14:294. 1901.

SYNONYMS. FISHKILL (2, 3). FISHKILL BEAUTY (1). *Fishkill Beauty* (2).

Fruit large, rather attractive in appearance, but not good enough in quality to displace standard sorts of its season either for culinary or dessert uses. Downing remarks that it is apt to decay on the tree (1), but this has not proved true in our experience with the variety. The tree comes into bearing rather early and is almost an annual bearer, yielding fair to good crops of uniformly large fruit. Should it possess superior hardiness it may be worthy of trial in those regions where this character is a prime requisite.


Historical. Origin Fishkill, N. Y.

TREE.

Tree vigorous with long, slender, curved branches; laterals willowy, long, slender. *Form* upright spreading or roundish, rather dense. *Twigs* short to medium, straight, moderately slender, rather geniculate; internodes medium to long. *Bark* brown tinged with clear reddish brown, mottled with scarfskin; slightly pubescent. *Lenticels* numerous, small or below medium, generally elongated and narrow, usually not raised. *Buds* small to above medium, plump, obtuse to somewhat acute, usually free or nearly so.

FRUIT.

Fruit very large to large, uniform in size and shape. *Form* nearly round varying to somewhat oblate or oblate conic, regular, obscurely ribbed. *Stem*



FISHKILL

medium to long, thick to slender. *Cavity* varying from somewhat obtuse to slightly acuminate, moderately deep to deep, rather broad, usually symmetrical, sometimes lipped, sometimes lightly russeted. *Calyx* rather small, closed or slightly open. *Basin* shallow, usually moderately wide, obtuse, angularly furrowed and wrinkled, with some tendency to develop mammiform protuberances.

Skin a little rough, thick, tough, dull yellow washed with dull or brownish-red, in highly colored specimens deepening to a bright red blush, mottled, striped and splashed with deep carmine, often overspread with thin whitish bloom. *Dots* numerous, conspicuous, medium size, russet. *Prevailing effect* attractive although rather dull red.

Calyx tube rather long, wide, varying from elongated conical to funnel-shape. *Stamens* median to marginal.

Core below medium to rather large, abaxile; cells usually symmetrical, open; core lines clasping. *Carpels* roundish to broadly obcordate, somewhat emarginate, slightly tufted. *Seeds* light brown, below medium to rather large, narrow, rather long, plump, varying from obtuse to acuminate.

Flesh whitish or tinged with yellow, firm, coarse, moderately crisp, moderately tender, juicy, mild subacid, fair to good in flavor and quality.

Season November to February.

FORD.

REFERENCES. 1. Downing, 1857:144. 2. Warder, 1867:719. 3. Thomas, 1875:499.

SYNONYMS. None.

The fruit of Ford is described as large, roundish, yellow; flesh solid, rather acid but of high flavor and good quality. Season October to January (1, 3). We do not know this variety and so far as we can discover it is no longer propagated. According to Downing it originated in Canaan, Columbia county, N. Y. (1).

FRANCHOT.

REFERENCE. 1. Downing, 1869:182.

SYNONYMS. None.

According to Downing this variety originated in Otsego county, N. Y. The tree is productive, the fruit medium size, yellow, shaded and splashed with red; flesh pleasant, aromatic, good. Season October to January (1). So far as we can learn it is not now being propagated.

FULLERTON SWEET.

REFERENCE. 1. Downing, 1869:185.

SYNONYMS. None.

A variety of unknown origin which has been fruited in Orange county (1). The fruit much resembles Autumn Bough. It is below medium, pale yellow; flesh tender, sweet, very good; season October and November. We do not know this variety and so far as we can learn it is not being propagated.

GARDEN ROYAL.

REFERENCES. 1. Manning, *Mag. Hort.*, 13:438. 1847. 2. Hovey, *Ib.*, 14:18. 1848. *fig.* 3. Thomas, *Cultivator*, 5:212. 1848. 4. Cole, 1849:106. *fig.* 5. Thomas, 1849:139. *fig.* 6. Barry, 1851:284. 7. *Am. Pom. Soc. Cat.*, 1854. 8. Elliott, 1854:81. 9. *Mag. Hort.*, 21:63. 1855. 10. Downing, 1857:79. 11. Warder, 1867:719. 12. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 13. Bailey, *Am. Hort.*, 1892:239. 14. *Rural N. Y.*, 60:247. 1901. 15. Budd-Hansen, 1903:88. *fig.* 16. Farrand, *Mich. Sta. Bul.*, 205:44. 1903.

SYNONYMS. None.

Garden Royal is not a good variety for commercial planting because the fruit is too small and its season early and short, but by many it is considered one of the very best dessert apples of late summer and early autumn. The fruit is of regular form, very handsome deep yellow striped with orange-red and dark crimson. The flesh is very tender, aromatic and with a delicate, pleasant acid flavor; season, August and September. The tree when full grown is of medium size, moderately vigorous, with roundish head. It appears to be hardy, healthy and long-lived, comes into bearing young and is a reliable biennial cropper.

Historical. Origin Sudbury, Mass. (2). It is occasionally found in this state in home orchards. It is now little propagated by nurserymen and seldom planted.

FRUIT.

Fruit medium or below. *Form* round, slightly oblate, often a little inclined to conic, regular or obscurely ribbed. *Stem* short to medium, straight, rather slender. *Cavity* acute sometimes approaching acuminate, rather deep and broad, slightly furrowed, often faintly russeted. *Calyx* small to above medium, open or partly closed; lobes often separated at base, rather short, acute. *Basin* moderately shallow, rather wide, obtuse to somewhat abrupt, slightly wrinkled.

Skin thin, greenish-yellow, sometimes entirely overspread with red, irregularly striped and splashed with carmine. *Dots* numerous, rather conspicuous, medium or above, often irregular, russet or yellowish.

Calyx tube medium size, funnel-shape. *Stamens* median to nearly marginal.

Core small, axile; cells closed or slightly open; core lines clasping. *Carpels* small, elliptical, emarginate.

Flesh tinged with yellow, fine, tender, juicy, agreeable mild subacid, aromatic, very good.

Season late August and September.

GARDNER SWEET PEARMAIN.

REFERENCE. 1. Downing, 1869:188.

SYNONYMS. None.

A Long Island variety, the fruit of which, according to Downing (1), is medium in size, nearly covered with red; flesh whitish, sweet, good; season September. This variety is unknown to us; so far as we have been able to learn it is no longer propagated.

GENESEE FLOWER.

REFERENCE. 1. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:290.

SYNONYMS. *Demary*. FLOWER (OF GENESEE) (1). *Flower of Genesee*. *Hawkins Pippin*. *Pride of Genesee*.

Fruit large, remarkably uniform in size and shape, with very few culls. In color and general appearance it resembles Pumpkin Sweet, commonly called Pound Sweet. It is not an apple of high dessert quality, but is very good for culinary uses, having a pleasant mild subacid flavor. It does not always sell well because the color is green and the variety is not generally well known. With some growers the tree is not a satisfactory cropper, but others find it a regular and abundant bearer, yielding good to heavy crops almost annually, and regard the variety as desirable for commercial planting.

Historical. This variety appears to be known by the name Genesee Flower more than by any other, but Mr. Nelson Bogue of Batavia informs us that it is also known locally under the various names Flower of Genesee, Pride of Genesee, Hawkins Pippin, and Demary. He states that the original tree, now about sixty years old, is still standing on the old Demary farm, in the town of Alexander, Genesee county, N. Y. The cultivation of this variety appears to be confined principally to the counties of Genesee, Wyoming and Orleans, and it does not appear to be increasing.

FRUIT.

Fruit large. *Form* roundish, somewhat oblate. *Stem* short, rather slender. *Cavity* broad, moderately deep, russeted. *Calyx* medium size, partly open. *Basin* shallow, irregular, often distinctly ridged.

Skin light green mingled with light yellow. *Dots* numerous, pale, with a few that are large and russet.

Flesh nearly white, mild subacid, good for culinary use but not much esteemed for dessert.

Season late September to November.

GIDEON.

REFERENCES. 1. Gideon, *Am. Pom. Soc. Rpt.*, 1885:26. 2. *Mo. Hort. Soc. Rpt.*, 1886:233. 3. *Am. Pom. Soc. Cat.*, 1889:6. 4. Bailey, *Am. Hort.*, 1892:239. 5. Beach, *N. Y. Sta. An. Rpt.*, 13:580. 1894. 6. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 4:2. 1897. figs. 7. Waugh, *Vt. Sta. Bul.*, 61:30. 1897. figs. 8. *Am. Gard.*, 22:132. 1901. figs. 9. *Me. Sta. An. Rpt.*, 18:83. 1902. 10. Hansen, *S. D. Sta. Bul.*, 76:51. 1902. fig. 11. Budd-Hansen, 1903:88. fig. 12. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:12. 1903. 13. Farrand, *Mich. Sta. Bul.*, 205:44. 1903. 14. Beach and Clark, *N. Y. Sta. Bul.*, 248:121. 1904.

SYNONYMS. GIDEON (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14). *Gideon White* (14).

Fruit decidedly attractive in general appearance being of good size, clear waxen yellow often with a bright pinkish blush but it is deficient in quality and apt to be defective in that the flesh becomes withered and discolored about the core as soon as the fruit reaches maturity. It is not a good variety for holding in cold storage as it stands heat poorly before going into storage and when in storage goes down quickly. In ordinary storage it is inferior to Hubbardston in keeping qualities, being in season from October to December or possibly later (14). The tree is of an exceptionally fine habit in the nursery, and in the orchard it is an upright vigorous grower, very hardy and healthy, comes into bearing young and yields full crops biennially or nearly annually. It is not recommended for planting in New York except it be as a stock upon which to topwork less hardy varieties.

Historical. Originated by Peter M. Gideon, Excelsior, Minn., from crab seed. The following is his statement of its parentage. "The Gideon is a seedling of the small crab; the seed came from Boston, where the tree was surrounded in the orchard by Blue Pearmain. I consider the tree a cross between those two. The tree resembles Blue Pearmain; seed was taken from the crab" (1).

TREE.

Tree medium to large, vigorous or moderately vigorous. *Form* at first upright but becoming spreading and open. *Twigs* short, curved, stout with large terminal buds; internodes medium. *Bark* brown mingled with some olive-green, lightly streaked with scarf-skin; heavily pubescent. *Lenticels* quite numerous, medium to small, oblong, slightly raised. *Buds* prominent, large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit usually above medium to large, uniform in size. *Form* roundish conical or somewhat ovate, sometimes inclined to oblong, angular. *Stem* long to below medium, slender. *Cavity* acute or somewhat acuminate, deep, broad to medium in width, sometimes lightly russeted. *Calyx* small, closed; lobes medium in length, rather narrow, acute, reflexed. *Basin* small to medium, sometimes oblique, shallow or very shallow, narrow to medium in width, rather obtuse, somewhat wrinkled, usually with narrow but not prominent ridges.

Skin rather thin, glossy, clear pale waxen yellow, sometimes with beautiful pink blush on exposed cheek. *Dots* light, submerged, inconspicuous, except where the skin is blushed.

Calyx tube short, narrow, often funnel-shape with very short, truncate cylinder. *Stamens* marginal to median.

Core medium size, axile or abaxile; cells closed or open; core lines meeting the limb or clasping the cylinder. *Carpels* round to broadly ovate or elliptical, emarginate, tufted. *Seeds* rather large, irregular, medium in width, rather long, not very plump, acute to acuminate, tufted, light brown.

Flesh whitish or tinged with yellow, of rather soft loose texture, a little coarse, crisp, juicy, brisk subacid to mild subacid, fair to good.

Season October.

GINNIE.

REFERENCES. 1. *N. Y. Sta. An. Rpt.*, 2:35. 1883. 2. Beach and Paddock, *N. Y. Sta. An. Rpt.*, 14:252, 257. 1895. 3. Beach and Clark, *N. Y. Sta. Bul.*, 248:121. 1904.

SYNONYMS. AUNT GINNIE (1, 2). Aunt Ginnie (3). GINNIE (3).

An autumn apple of good color and good quality. Tree rather upright, moderately vigorous, moderately productive. It comes into bearing early and yields some fruit nearly every year. Not recommended for planting in New York.

Historical. Received from Ellwanger and Barry, Rochester, N. Y., in 1883 for testing at this Station. We do not know its origin. So far as we have learned it is practically unknown among New York fruit growers.

FRUIT.

Fruit medium to large. *Form* oblate conic, broad and flattened at the base, obscurely ribbed. *Stem* medium, usually not exerted. *Cavity* large, acuminate, broad, deep, with conspicuous, broad, irregular, outspreading russet rays. *Calyx* small, nearly closed. *Basin* shallow to medium in depth, rather narrow to moderately wide, furrowed and wrinkled.

Skin yellow nearly covered with bright red and marked with broad stripes and splashes of bright carmine. *Dots* minute, russet. *Prevailing effect* red striped.

Calyx tube funnel-form. *Stamens* basal or nearly so.

Core very large, abaxile; cells pretty symmetrical, open; core lines clasp the funnel cylinder. *Carpels* broadly roundish, emarginate, tufted. *Seeds* medium in size, rather wide, obtuse to somewhat acute.

Flesh whitish, rather coarse, moderately juicy, aromatic, subacid, good to very good.

Season late September to early winter; commercial limit November in common storage (3).

GLADSTONE.

REFERENCES. 1. Hogg, 1884:150. 2. *N. Y. Sta. An. Rpt.*, 11:224. 1892. 3. Hansen, *S. D. Sta. Bul.*, 76:52. 1902.

SYNONYMS. GLADSTONE (2, 3). MR. GLADSTONE (1).

Fruit of good size, and when highly colored rather attractive. In general appearance it is intermediate between Oldenburg and Gravenstein, perhaps resembling Gravenstein in color more than Oldenburg. It is not equal to either of these varieties in quality. The tree comes into bearing young, is an annual cropper and productive. Not recommended for planting in New York.

Historical. This is a comparatively recent introduction from England. It has not been extensively disseminated and so far as we can learn its cultivation is not increasing in this country.

TREE.

Tree rather small, moderately vigorous with short, stout branches. *Form* spreading and inclined to droop. *Twigs* moderately long, curved, stout, with large terminal buds; internodes medium in size. *Bark* brown, tinged with olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* numerous, conspicuous, medium size, round, slightly raised. *Buds* medium size, broad, flat, obtuse, appressed, slightly pubescent.

FRUIT.

Fruit medium to large, uniform. *Form* roundish oblate, slightly conic, obscurely ribbed; sides usually unequal. *Stem* medium in length, moderately slender. *Cavity* acuminate to acute, rather wide, moderately deep to deep, sometimes with outspreading russet. *Calyx* below medium to rather large, closed or somewhat open. *Basin* rather small, shallow to medium in depth, medium in width to rather wide, a little abrupt, slightly furrowed and wrinkled.

Skin thin, smooth, greenish-yellow or pale yellow, more or less thinly overspread with red, irregularly mottled, splashed and distinctly striped with pinkish carmine. *Dots* rather numerous, inconspicuous, light colored, submerged.

Calyx tube rather short, cone-shape to funnel-form. *Stamens* median.


Core medium to rather large, usually axile; cells symmetrical, closed or slightly open; core lines clasping. *Carpels* broadly roundish, very slightly emarginate. *Seeds* rather dark brown, very wide, flat, obtuse to acute.

Flesh slightly tinged with yellow, moderately firm, a little coarse, crisp, tender, juicy, mild subacid, fair in quality.

Season September and October.

GLORIA MUNDI.

REFERENCES. 1. *Dom. Encyc.*, 1804 (cited by 4). 2. *Am. Gard. Cal.*, 1806:385. 3. *Coxe*, 1817:117. fig. 4. *Thacher*, 1822:131. 5. *Buel, N. Y. Bd. Agr. Mem.*, 1826:476. 6. *Fessenden*, 1828:129. 7. *London Hort. Soc. Cal.*, 1831:No. 447. 8. *Kenrick*, 1832:49. 9. *Floy-Lindley*, 1833:86. 10. *Downing*, 1845:110. fig. 11. *Cole*, 1849:122. 12. *Thomas*, 1849:182. 13. *Phœnix. Horticulturist*, 4:470. 1850. 14. *Emmons, Nat. Hist. N. Y.*, 3:41. 1851. col. pl. No. 35 and fig. 15. *Elliott*, 1854:174. 16. *Mag. Hort.*, 23:83. 1857. 17.



GLADSTONE

Hooper, 1857:39. 18. Warder, 1867:719, 725. 19. Downing, 1869:191. 20. Fitz, 1872:168. 21. Leroy, 1873:407. fig. 22. Downing, 1881:11 index, app. 23. Hogg, 1884:85. 24. Wickson, 1889:245. 25. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 26. Green, *Rural N. Y.*, 49:604. 1890. 27. Bailey, *Am. Hort.*, 1892:240. 28. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 29. *Can. Hort.*, 23:75. 1900. 30. Bruner, *N. C. Sta. Bul.*, 182:20. 1903.

SYNONYMS. *American Gloria Mundi* (4, 7, 21). *American Mammoth* (7, 10, 15, 17, 19, 21). *Baltimore* (10, 12, 14, 21, 23, 26, of some 19). *Baltimore Pippin* (15, 17, 19). *Belle Dubois* (23). *Belle Josephine* (19, 21). *Copp's Mammoth* (22). *Glazenwood* (10). *Glazenwood Gloria Mundi* (19, 21, 23). *GLORIA MUNDI* (1, 7, 10, 11, 14, 16, 17, 19, 22, 23, 24, 25, 27, 28, 29, 30). *Gloria Mundi* (5, 10, 12, 13, 15, 18, 20, 21, 26). *Impératrice Joséphine* (21). *JOSÉPHINE* (21). *Kinderhook Pippin* (14). *Mammoth* (21, 23). *MAMMOTH PIPPIN* (2, 3, 4, 5, 6, 8, 9, 12, 13, 15, 18, 20, 26). *Mammoth Pippin* (7, 10, 11, 14, 16, 17, 19, 21). *Melon* (21). *Mississippi?* (19). *Monstreuse Pippin* (21). *Monstrous Pippin* (23). *Mountain Flora* (22). *N. Y. Gloria Mundi* (3, 7, 8, 9, 10, 15, 19, 21). *Ox Apple* (10, 12, 14, 15, 19, 23, 26). *Pound* (17). *Vandyne Apple* (9).

Fruit of largest size; seldom cultivated except for exhibition; suitable only for culinary purposes. The tree is large, vigorous, spreading, hardy and long-lived. It has been commonly held to be unproductive, but a few fruit growers in Southeastern New York report that it is a good bearer and a profitable commercial variety. Season October to January.

Historical. The exact place of origin of this apple seems doubtful. In 1804 Mease stated, "It originated on the farm of Mr. Crooks, near Red Hook in New York" (1). Thirteen years later Cox credited it with a Long Island origin (3). All that can be stated with certainty is that it is an old variety known in parts of Eastern New York, New Jersey and Pennsylvania in the beginning of the last century.

FRUIT.

Fruit very large or large, uniform. *Form* roundish with truncate ends, slightly conical, ribbed; sides usually unequal. *Stem* medium to short, moderately thick. *Cavity* large, acuminate, moderately deep to deep, broad, furrowed and compressed, sometimes slightly russeted. *Calyx* medium to large, open or partly closed; lobes separated at base, short, narrow. *Basin* large, moderately deep to deep, rather wide, somewhat abrupt, sometimes compressed, furrowed and wrinkled.

Skin greenish-yellow sometimes with faint bronze blush. *Dots* small to medium, often areolar with russet center, or light colored and submerged. *Prevailing effect* yellow or greenish.

Calyx tube very large, long, wide, broadly conical extending to core or approaching cylindrical below. *Stamens* median.

Core medium to rather large, usually decidedly abaxile; cells symmetrical, open; core lines usually clasping. *Carpels* broadly roundish to somewhat elliptical, slightly tufted. *Seeds* moderately dark brown, medium to rather small, rather narrow, short, plump, obtuse to acute, sometimes tufted.

Flesh slightly tinged with greenish-yellow, coarse, moderately crisp, rather tender, juicy, rather mild subacid, fair or nearly good in quality.

Season October to January.

THE GOLDEN PIPPINS.

The name Golden Pippin has been applied to several distinct varieties, the most important of which are mentioned below.

The Golden Pippin of England is a small, yellowish apple with shade of red, which is in season from November to March. This has already been described in Volume I, page 141.

The Golden Pippin of Westchester County, also known under the name of American Golden Pippin and by various other synonyms, is described as Golding, page 82, which is the name now accepted for this variety by pomologists. In addition to the varieties described below under the name Golden Pippin there are several other sorts which have been known under this name.

(I) GOLDEN PIPPIN.

REFERENCES. 1. Downing, 1869:195. 2. Thomas, 1875:500. 3. Bailey, *An. Hort.*, 1892:240. 4. Ragan, *U. S. B. P. I. Bul.*, 56:123. 1905. 5. *Ib.*, 56:347. 1905.

SYNONYMS. *Butter Pippin* (1). GOLDEN PIPPIN (1, 2, 3, 4, 5). *Large Golden Pippin* (1). *Mammoth* (1, 2). *Pound Royal* (2, of some 1). *York Pippin* (1, 5).

This variety belongs in the Fall Pippin group of apples. In Central and Western New York it is often called York Pippin. The fruit is large, coarse-grained, with a very pleasant flavor, and is suitable for either dessert or culinary uses. As compared with Fall Pippin the fruit is harder, keeps longer and stands shipping better in hot weather. In Western New York its season in ordinary storage extends from about September 20 to January 1. It holds its flavor and quality well for a late fall and early winter sort. It frequently brings better prices than Fall Pippin, and we are informed it is sometimes exported. The tree is more spreading than that of Fall Pippin, and is perhaps somewhat hardier. It is generally healthy, vigorous, quite long-lived, and when full grown becomes a pretty large tree. It is a reliable cropper, bearing good to heavy crops biennially or sometimes annually. There is apt to be considerable loss by premature dropping of the fruit, and unless proper preventive measures are taken the crop may be seriously injured by apple scab and codling moth, but with proper treatment

these pests may be kept under good control. Golden Pippin is grown to a considerable extent in various parts of New York state and in New England. In some sections it is regarded as one of the most desirable of the fall varieties for commercial planting.

Historical. The origin of this variety is unknown. It has long been in cultivation. In New York it is now found mostly in old orchards and, generally speaking, its cultivation is not increasing.

FRUIT.

Fruit very large or large, pretty uniform in size and shape. *Form* roundish to roundish oblate, sometimes slightly conic, often with a broad flat base and broadly ribbed toward the apex. *Stem* long to medium, thick, sometimes swollen. *Cavity* acute to acuminate, medium in depth, broad, usually rather symmetrical, sometimes lipped, russeted and often with heavy, outspreading russet rays. *Calyx* rather large, closed. *Basin* deep, moderately wide to wide, abrupt, slightly furrowed, sometimes irregularly compressed.

Skin rather tender, green or yellowish changing to a deeper and rather attractive yellow when fully mature, sometimes with bronze blush and russet flecks. *Dots* small to rather large and conspicuous, greenish and submerged or with russet point.

Calyx tube wide, conical. *Stamens* median to somewhat basal.

Core rather small, somewhat abaxile; cells open; core lines meeting or slightly clasping. *Carpels* roundish or approaching elliptical, sometimes obovate, heavily tufted. *Seeds* few, often not perfectly developed, medium size, irregular, rather dark brown, rather plump, acute.

Flesh yellowish, coarse, rather tender, juicy, agreeable mild subacid, somewhat aromatic, good to very good.

Season late September to December or January.

(II) GOLDEN PIPPIN.

REFERENCES. 1. Downing, 1869:194. 2. Bailey, *An. Hort.*, 1892:240.

SYNONYMS. GOLDEN PIPPIN (1, 2). *Pittstown Pippin* (1).

Downing states that this is an old apple grown many years ago in Adams, Mass. Tree upright and vigorous; fruit rather large, roundish oblate, sometimes conic, greenish-yellow with blush; flesh yellowish, sprightly subacid, good for cooking; season September and October (1).

It appears that this variety is no longer propagated in New York.

GOLDEN REINETTE.

REFERENCES. 1. Budd, *Ia. Agr. Coll. Bul.*, 1885:32, 35, 38. 2. *Ib.*, 1890:25. 3. *Ib.*, *Ia. Sta. Bul.*, 19:538. 1892. 4. Munson, *Me. Sta. An. Rpt.*, 1896:74, 80. 5. Lyon, *Mich. Sta. Bul.*, 152:221. 1898. 6. Munson, *Me. Sta. An. Rpt.*, 18:83. 1902. 7. Ragan, *U. S. B. P. I. Bul.*, 56:124. 1905.

SYNONYMS. No. 51 *Vor.* (1, 2). No. 10 (7). *Solotoc renet* (1). *Zolotoi renet* (7).

In addition to the old English variety known as Golden Reinette which has been described in Volume 1, page 142, there are at least two Russian apples that have been disseminated in this country under this name. One is a variety described by Munson as a promising autumn apple for Northern Maine, in season from September to December; fruit small, golden yellow washed and splashed with carmine (4, 6). What appears to be the same variety was received by this Station from Professor Budd, Ames, Ia., in 1890. It is decidedly inferior to the standard varieties of its season, which are in general cultivation throughout New York. Possibly on account of superior hardness it may have some value in the more northern or elevated regions of the state.

TREE.

Tree moderately vigorous with short, moderately stout, curved branches. *Form* upright spreading or roundish, rather dense. *Twigs* long, curved, stout; internodes medium. *Bark* brown, mingled with reddish-brown, mottled with scarf-skin, slightly pubescent. *Lenticels* numerous, conspicuous, large, oblong, raised. *Buds* prominent, large, broad, long, plump, acute, free, slightly pubescent.

FRUIT.

Fruit medium to below. *Form* oblate conic to strongly roundish conic, flattened at the base, ribbed, rather irregular; sides unequal. *Stem* short, thick. *Cavity* acuminate, medium in depth to rather deep, moderately broad, often compressed, usually not russeted. *Calyx* large, open or partly closed, leafy; lobes usually separated at base, long, acute to acuminate. *Basin* shallow to moderately deep, medium in width to narrow, usually abrupt, slightly furrowed and wrinkled.

Skin pale greenish-yellow with faint splashes and stripes of red.

Calyx tube large, wide, broadly conical. *Stamens* basal to median.

Core medium size, often abaxile; cells usually unsymmetrical, closed or open; core lines clasping or nearly meeting. *Carpels* broadly roundish to elliptical. *Seeds* medium brown, medium to below, plump, rather obtuse.

Flesh yellowish, crisp, juicy, mild subacid.

Season September to December.

The other Russian apple, above referred to, was disseminated some years ago by Professor Budd. In 1885 he remarked that he was sending out, under the name Golden Reinette (51 Vor.), a variety having fruit medium to large, golden in color, fine-grained, juicy, subacid, almost best in quality (1). In 1890 he stated: "This has proven a fine tree on a great variety of soils where the air had free circulation. Its northern limit is not yet known, but I have not known its wood colored at Ames when the Wealthy by its side in nursery was nearly killed. Fruit medium to large, golden yellow, fine-grained, subacid, and nearly best in quality. Season here, December to February, depending on time of picking and mode of storage" (2). In 1892 Budd further reported: "This has not proven true to name as received from the Bogdanoff estates in Russia. It is a member of the Anis family, of fine size and excellent quality. Season late fall, and early winter north" (3).

GOLDEN SWEET.

REFERENCES. 1. Kenrick, 1832:37. 2. Downing, 1845:84. 3. Phoenix, *Horticulturist*, 1:361. 1846. 4. *N. Y. Agr. Soc. Trans.*, 1846:189. 5. Thomas, 1849:136. 6. Cole, 1849:102. fig. 7. Barry, 1851:280. 8. Emmons, *Nat. Hist. N. Y.*, 3:40. 1851. fig. 9. Elliott, 1854:81. fig. 10. Gregg, 1857:37. fig. 11. Hooper, 1857:41. 12. *Am. Pom. Soc. Cat.*, 1862. 13. Warder, 1867:551. fig. 14. Downing, 1869:196. fig. 15. Wickson, 1889:244. 16. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 17. Bailey, *Am. Hort.*, 1892:240. 18. Waugh, *Vt. Sta. An. Rpt.*, 14:295. 1901. 19. Budd-Hansen, 1903:90. fig.

SYNONYMS. *Early Golden Sweet* (14). *GOLDEN SWEET* (2, 3, 4, 5, 6, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19). *GOLDEN SWEETING* (7, 10). *Orange Sweet* (6, 9). *ORANGE SWEETING* (1). *Orange Sweetening* (2, 13, 14). *Trenton Early* (9). *Yellow Sweeting* ? (1).

Fruit of good medium size, attractive clear yellow when fully mature, rich, sweet, very good in flavor and quality. Cultivated principally for home use. Of no commercial value except that it is sold in limited quantities in local markets. In season from the middle of August to the last of September. The tree is a good grower, healthy, hardy, moderately long-lived, comes into bearing rather young and yields moderate to heavy crops biennially.

Historical. An old Connecticut variety (2). Its exact origin is unknown. It has been pretty generally disseminated throughout the state but is nowhere grown extensively. It is listed by nearly all nurserymen (17).

TREE.

Tree large, vigorous. *Form* roundish spreading, inclined to droop, dense. *Twigs* long, curved, slender; internodes long. *Bark* brown, lightly mottled with scarf-skin; pubescent. *Lenticels* quite numerous, medium size, oval, slightly raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit below medium to nearly large, uniform in size and shape. *Form* roundish to roundish oblate or somewhat ovate, regular or faintly ribbed. *Stem* very long, moderately thick. *Cavity* acute, of medium depth, medium in width to rather narrow, symmetrical, usually partly russeted, often with outspreading russet rays. *Calyx* medium to small, closed; lobes medium in length, narrow, acute. *Basin* shallow to moderately deep, narrow to medium in width, somewhat obtuse, smooth, symmetrical, furrowed.

Skin thin, tender, smooth, waxy, yellowish-green becoming clear pale yellow when fully mature.

Calyx tube medium in width, cone-shape to truncate funnel-form. *Stamens* median.

Core medium to rather small, abaxile; cells often unsymmetrical, open; core lines clasping the funnel cylinder. *Carpels* ovate. *Seeds* medium to rather small, rather narrow, angular, acute, medium brown.

Flesh yellowish-white, firm, fine-grained, moderately tender, juicy, very sweet, aromatic, good to very good.

Season mid-August to late September.

GOLDEN WHITE.

REFERENCES. 1. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:447. 2. Brodie, *Montreal Hort. Soc. Rpt.*, 1885:72. 3. Hoskins, *Mich. Hort. Soc. Rpt.*, 1886:221. 4. *Montreal Hort. Soc. Rpt.*, 1886-7:16. 5. *Can. Hort.*, 13:157, 332. 1890. 6. Beach and Paddock, *N. Y. Sta. An. Rpt.*, 14:252. 1895. 7. Beach, *W. N. Y. Hort. Soc. Rpt.*, 1896:50. 8. Buad-Hansen, 1903:91.

SYNONYMS. *Beel Solotofskaja* (1). GOLDEN WHITE (1, 2, 3, 4, 5, 6, 7, 8). No. 978 (3, 5, 8). No. 979 (8). No. 981 (8).

Fruit of medium size, greenish-yellow, streaked with bright red in the sun, subacid, fair quality; season September. The tree comes into bearing rather young, and yields full crops biennially. Evidently not desirable for planting in New York.

Historical. A Russian apple received for testing at this Station from T. H. Hoskins, Newport, Vt., in 1888.

TREE.

Tree rather small, a slow grower with short, stout branches. *Form* spreading, open. *Twigs* short, curved, stout, with large terminal buds; internodes short. *Bark* dull brown mingled with olive-green, coated with gray scarf-skin; heavily pubescent. *Lenticels* scattering, large, oval or elongated, raised. *Buds* large, prominent, broad, plump, obtuse, free, much pubescent.

FRUIT.

Fruit medium size. *Form* oblate inclined to conic, flattened at the base, ribbed. *Stem* small to medium. *Cavity* small, acuminate, narrow, moderately shallow, russeted. *Calyx* open or partly open. *Basin* large, irregular, moderately deep, wrinkled.

Skin greenish-yellow nearly overlaid with red and striped with carmine. *Dots* numerous, large, light. *Prevailing effect* red or striped red.

Calyx tube large, cone-shape to funnel-form.


Core medium size, somewhat abaxile; cells usually symmetrical, closed or partly open; core lines clasping. *Carpels* roundish, emarginate, tufted. *Seeds* above medium, plump, wide, obtuse.

Flesh white with faint salmon tinge, fine-grained, moderately juicy, subacid, fair or sometimes good in quality.

Season September and October.

GOLDING.

REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 479. 2. Kenrick, 1835:64. 3. Downing, 1857:79. fig. 4. *Mag. Hort.*, 30:162. 1864. 5. Warder, 1867:636. fig. 6. Thomas, 1875:226. 7. *Am. Pom. Soc. Cat.*, 1877:6.



GOLDEN SWEET

8. Barry, 1883:342. 9. Lyon, *Mich. Sta. Bul.*, 105:108. 1894. 10. *Am. Pom. Soc. Cal.*, 1897:113. 11. Budd-Hansen, 1903:92. fig. 12. Ragan, *U. S. B. P. I. Bul.*, 56:125. 1905.

SYNONYMS. *American Golden* (12). *AMERICAN GOLDEN PIPPIN* (2, 4, 5, 7, 8). *American Golden Pippin* (3, 6, 10, 11, 12). *Golden Apple* (12). *GOLDEN PIPPIN* (3, of Westchester Co., 6). *Golden Pippin* (5, of Westchester Co., 8 and 12). *GOLDING* (1, 9, 10, 11, 12). *Newtown Greening* (3, 12). *N. Y. Greening* (3, 5, 6, 12). *Ribbed Pippin* (3, 12).

Fruit medium or above, yellow, sometimes with shade of brownish-red; flesh rather coarse, aromatic, subacid and excellent in quality for either dessert or culinary uses. It is much subject to scab and a comparatively large percentage of the crop is apt to be unmarketable unless thorough treatment is given to prevent injury from insects and fungus diseases. The tree is rather large, vigorous, hardy, long-lived and moderately productive yielding moderate to heavy crops biennially. In some sections it is regarded as a pretty good commercial variety particularly in portions of Eastern New York.

Historical. In 1857 Downing wrote (3) regarding this variety, that although it was one of the finest American fruits and an old variety, it was not generally known. It was said to have been cultivated in Westchester and adjoining counties for more than fifty years where it was considered profitable for market and superior for family use. So far as we can learn *Golding* is now seldom or never planted in New York.

GRANDMOTHER.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 1881-82:39, 73. 2. Budd, *Ia. Hort. Soc. Rpt.*, 1882:78. 3. *Montreal Hort. Soc. Rpt.*, 1883:105. fig. 4. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:443. 5. *Am. Pom. Soc. Rpt.*, 1883:72. 6. Budd, *Ia. Agr. Coll. Bul.*, 1883:27. 7. *Ib.*, 1885:18. 8. Schroeder, *Montreal Hort. Soc. Rpt.*, 12:71. 1886-7. 9. *Ia. Agr. Coll. Bul.*, 1890:24. 10. Bailey, *Am. Hort.*, 1892:240. 11. Budd, *Ia. Sta. Bul.*, 19:542. 1892. 12. Munson, *Me. Sta. An. Rpt.*, 12:74, 80. 1896. 13. Thomas, 1897:277. fig. 14. Macoun, *Can. Dept. Agr. Rpt.*, 1899:79. 15. Troop, *Ind. Sta. An. Rpt.*, 12:80. 1899. 16. Hansen, *S. D. Sta. Bul.*, 76:54. 1902. fig. 17. Budd-Hansen, 1903:92.

SYNONYMS. *Baboushkino* (1). *Babuscheno* (16, 17). *BABUSCHKINO* (3). *Habushkino* (4, 7). *Bogdanoff* (14). *Bogdanoff Steklianka* (14). *GRANDMOTHER* (2, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17). *Grandmother* (3). *GRAND MOTHER* (1, 10). *No. 6 M* (1, 2, 6, 9, 11). *No. 6* (8). *No. 469* (1, 2, 3, 6, 7, 11, 12, 16, 17). *No. 84 Vor.* (9, 11). *Red Reinette* (14).

Fruit of good medium size, greenish-yellow, sometimes with slight blush. It shows the marks of handling readily, is not very uniform in size nor does it excel in quality. Season late fall and early winter. The tree is a good thrifty grower, comes into bearing young, is reliably productive and is almost an annual bearer, yielding moderate to good crops. It does not appear to be valuable for planting in New York.

Historical. A Russian variety which has been disseminated for trial in various sections of the country. It has been thus far but little planted in this state.

TREE.

Tree large, rather vigorous with long, stout branches. *Form* spreading, rather flat, open. *Twigs* medium length, curved, moderately stout; internodes short. *Bark* dark brown or reddish-brown, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, medium size, roundish, raised. *Buds* medium to large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit medium to rather large. *Form* ovate to roundish conic, quite flat at base, a little angular; sides unequal; fairly uniform. *Stem* short or very short, rather thick. *Cavity* acute to acuminate, narrow, moderately deep to shallow or scarcely depressed, much russeted and often with outspreading russet. *Calyx* large to medium, closed or somewhat open; lobes medium in length, broad, acute. *Basin* deep to medium in depth, medium in width to rather wide, abrupt, usually furrowed and wrinkled.

Skin smooth, light greenish-yellow, sometimes with faint blush. *Dots* numerous, inconspicuous, light and submerged, or sometimes areolar with dark center.

Calyx tube medium in length, wide, broadly conical. *Stamens* basal.



Core medium to small or abortive, axile or abaxile; cells symmetrical, usually closed; core lines meeting. *Carpels* variable, irregular, broadly ovate or obovate, emarginate, slightly tufted. *Seeds* below medium to rather large, rather dark brown, plump, obtuse to acute.

Flesh with slight green or yellow tinge, moderately firm, coarse, neither crisp nor tender, very juicy, subacid to briskly subacid, slightly aromatic, fair to good in flavor and quality.

Season November to January.

GRAVENSTEIN.

REFERENCES. 1. Forsyth, 1824:105. 2. *Pom. Mag.*, 3:No. 98. 1830. col. pl. 3. *London Hort. Soc. Cat.*, 1831:No. 489. 4. Kenrick, 1832:34. 5. Floy-Lindley, 1833:52. 6. Manning, 1838:53. 7. *Ib.*, *Mag. Hort.*, 7:45. 1841. 8. Downing, 1845:85. fig. 9. *Horticulturist*, 1:196. 1846. 10. Cole, 1849:110. fig. 11. Thomas, 1849:149. fig. 12. *Horticulturist*, 5:200. 1850. 13. Barry, 1851:284. 14. Emmons, *Nat. Hist. N. Y.*, 3:32, 104. 1851. fig., col. pl. No. 28. 15. Hovey, 2:15. 1851. col. pl. 16. *Am. Pom. Soc. Cat.*, 1852. 17. Elliott, 1854:83. 18. Gregg, 1857:43. fig. 19. Hooper, 1857:41, 107, 111. 20. De Wolf and Hovey, *Mag. Hort.*, 23:444. 1857. 21. Warder, 1867:487. fig. 22. Fitz, 1872:143, 149, 152. 23. Leroy, 1873:338. fig. 24. *Horticulturist*, 30:32. 1875. 25. Downing, 1881:11 index, app. 26. Hogg, 1884:96. 27. Wickson, 1880:243. 28. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:202. 29. Bailey, *Am. Hort.*, 1892:240. 30. Stinson, *Ark. Sta. An. Rpt.*, 6:55. 1893. 31. Gaucher, 1894:No. 10. col. pl. 32. Saunders, *Ont. Fr. Gr. Assn. An. Rpt.*, 27:51. 1895. 33. *Amer. Gard.*, 16:381. 1895. fig. 34. Burrill and McCluer, *Ill. Sta. Bul.*, 45:303. 1896. 35. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 36. *Can. Hort.*, 22:133. 1899. 37. Van Deman, *Rural N. Y.*, 59:20. 1900. 38. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 8:8. 1901. figs. 39. Waugh, *Vt. Sta. An. Rpt.*, 14:295. 1901. 40. *Can. Hort.*, 25:167. 1902. fig. 41. Budd-



GRAVENSTEIN

Hansen, 1903:93. fig. 42. Bruner, *N. C. Sta. Bul.*, 182:20. 1903. 43. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:43. 1903. 44. Beach and Clark, *N. Y. Sta. Bul.*, 248:122. 1904.

SYNONYMS. *Blumen-Calvill* (31). *Calville de Gravenstein* (31). *Calville Grafensteiner* (23, 31). *Diel's Sommerkönig* (31). *Early Congress* (25). *Grafen-Apfel* (31). *Grafensteiner* (15, 23, 31). *Grave Slige* (23). *Grave Slije* (8, 15, 17, 31). GRAVENSTEIN (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44). *Gravenstein* (31). GRAVENSTEINER (31). *Gravensteiner* (23). *Gravenstine* (31). *Ohio Nonpareil* (17, 23, 31). *Paradies Apfel* (31). *Pomme Graefenstein* (31). *Prinzessin-Apfel* (31). *Ripp Apfel* (31). *Sabine* (31). *Sabine of the Flemings* (15, 23). *Strohmer* (31). *Stromling* (31). *Tom Harryman* (31).

Gravenstein is an apple of good size, attractive appearance and excellent quality. For culinary purposes it is perhaps unexcelled by any variety of its season. It often sells at comparatively high prices. In many sections of the state the tree is regarded as not quite hardy, being somewhat subject to sunscald and canker. It comes into bearing moderately early, is quite productive and a pretty reliable cropper. The crop ripens continuously during a period of several weeks and should have two or three pickings. Beginning in the latter half of September it continues in season till early November. When properly handled a considerable portion of the fruit may remain apparently sound much later than this but the color fades and the quality and flavor deteriorate. As compared with other varieties of its class it stands up well in good dry cold storage (44). There is apt to be considerable loss from the dropping of the fruit before it is properly colored and there is also a rather high percentage of low grade or unmarketable fruit. In spite of these serious faults its cultivation in commercial orchards is gradually increasing in some sections of the state, particularly in portions of the Hudson valley where fruit growers find it a desirable apple for both domestic and foreign markets.

Historical. The following excellent account of the history of this variety was given by Hovey in 1851 (15). "The origin of the Gravenstein remains in some doubt. It is said to have been originally found in the Duke of Augustinberg's garden at Gravenstein, in Holstein, and that the original tree was growing there in the middle of the last century; another statement is, that it derived its name from being found in the garden of the castle of Gräfenstein, in Sleswick; and Diel says, that it was supposed by some to

have been introduced from Italy. Be this as it may, it is a common apple throughout Germany and Sweden, and was received from thence into the English collections. It is undoubtedly of similar origin with the Red Astrachan and Duchess of Oldenburg, possessing the peculiar habit of growth, form of foliage, and texture of the fruit, which distinguish the German apples.

"At what time it was first introduced into our gardens we are not aware. But as neither Coxe or Thacher describe or name it, we suppose it was some time subsequent to the account given of it in the Transactions of the London Horticultural Society in 1822. It is at the present time considerably cultivated, though not to the extent its merits deserve."

In 1857 Captain DeWolfe stated that the Gravenstein was imported by him from Denmark in May, 1826 (20). In a letter dated October 11, 1820, published in the New England Farmer, Judge Buel, of Albany, called attention to the importation of Gravenstein and other German apples, trees of which he had presented to the members of the Massachusetts Horticultural Society. Hovey states that in view of this it appears that Gravenstein was imported to the vicinity of Albany probably prior to 1826 (20). After these early importations Gravenstein gradually found its way into cultivation in various portions of the country. For many years it has been pretty generally disseminated through New York state, but in most localities it is grown to a limited extent only.

TREE.

Tree large, vigorous. *Form* upright spreading to roundish, open. *Twigs* medium to long, curved, moderately stout; internodes long. *Bark* brownish-red, mingled with olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* very scattering, medium to small, oblong, not raised. *Buds* medium in size, plump, acute, free, pubescent.

FRUIT.

Fruit large to above medium, fairly uniform in size but not in shape. *Form* oblate to roundish, somewhat irregular, broad at the base, slightly angular about the basin. *Stem* short to medium, thick. *Cavity* rather large, acute to acuminate, moderately deep to deep, rather narrow to broad, irregularly russeted. *Calyx* large, open or sometimes closed; lobes large, long, very broad, acute. *Basin* irregular, medium in depth to deep, medium to wide, obtuse to somewhat abrupt, wrinkled.

Skin thin, tender, slightly rough, greenish-yellow to orange-yellow overlaid with broken stripes of light and dark red. *Dots* few, small, light. *Prevailing effect* yellow striped.

Calyx tube large, conical to funnel-shape. *Stamens* median.

Core medium in size, strongly abaxile; cells open; core lines clasping the funnel cylinder. *Carpels* broadly ovate, emarginate. *Seeds* medium to large, medium in width, rather long, plump, acute to acuminate, medium brown.

Flesh yellowish, firm, moderately fine, crisp, moderately tender, juicy, sprightly subacid, aromatic, very good to best.

Season late September till early November.

RED TYPES OF THE GRAVENSTEIN.

Several instances are known where bud sports of the Gravenstein have originated which bear highly colored red fruit, but in other respects are quite like the typical Gravenstein. Some of these sports have neither been described nor propagated but others of them have been introduced into cultivation under distinct names. Gaucher and Leroy each describe a Red Gravenstein.¹ A Red Gravenstein which originated in Nova Scotia is now cultivated under the name Banks. For an account of this variety the reader is referred to Banks, page 14.

GREAT MOGUL.

REFERENCES. 1. Regel, 1:453. 1868. 2. Budd, *Ia. Agr. Coll. Bul.*, 1885:24, 35, 36, 37. 3. Lyon, *U. S. Pom. Bul.*, 2:141. 1888. 4. Budd, *Ia. Sta. Bul.*, 18:519. 1892. 5. Troop, *Ind. Sta. Bul.*, 53:124. 1894. 6. Stinson, *Ark. Sta. An. Rpt.*, 9:105. 1896. 7. Troop, *Ind. Sta. Rpt.*, 12:80. 1899. 8. *Ib.*, *Ia. Sta. Bul.*, 41:85. 1899. 9. Ragan, *U. S. B. P. I. Bul.*, 56:128. 1905.

SYNONYMS. GREAT MOGUL (1, 2, 3, 4, 5, 6, 7, 8, 9). 54 M (3). *Grosser Mogul* (2, 9). *Vilikui Mogul* (2, 9).

Fruit large, greenish, blushed and striped with red, somewhat resembling Alexander in type but tending to be more oblong, less broadly striped with carmine, and on the whole less attractive in color and form. Tree a fine grower, comes into bearing rather young, is an annual bearer and productive. It is not recommended for growing in New York.

Historical. An apple of Russian origin which was introduced into the United States about twenty-five years ago (1, 2, 3).

TREE.

Tree a good grower when young but when full grown is rather below medium size. *Form* open, spreading, rather drooping with rather short stout branches and drooping laterals. *Twigs* below medium to short, stout, irregularly geniculate. *Bark* clear brownish-red to very dark brown almost black, mottled lightly with gray scarf-skin, slightly pubescent. *Lenticels* very conspicuous, medium to large, oblong, generally elongated and russeted. *Buds* very prominent, large, broad, plump, acute, free, lightly attached to the bark, scales not well united, pubescent.

FRUIT.

Fruit large to very large. *Form* roundish ovate, sometimes varying to oblong conic or to oblate conic, slightly angular. Usually the fruit is pretty regular in form, shape and size. *Stem* medium to nearly long, rather thick, often clubbed or swollen. *Cavity* usually very acuminate, sometimes acute, moderately deep, moderately wide, sometimes with outspreading russet rays, smooth, symmetrical, often lipped. *Calyx* medium to rather large, closed or

¹ Leroy, 1873:339. Gaucher, 1894:No. 11.

slightly open, lobes long, broad, acute. *Basin* rather small, often oblique, narrow to medium in width, shallow to moderately deep, somewhat abrupt, smooth or slightly wrinkled, symmetrical.

Skin rather thick, tough, smooth, somewhat waxy, pale greenish-yellow more or less overspread with rather bright pinkish-red, often indistinctly striped with carmine. *Dots* inconspicuous, numerous, small, scattering, gray.

Calyx tube medium, elongated, conical or somewhat funnel-form with wide limb and fleshy pistil point projecting into the base. *Stamens* marginal to median.

Core somewhat abaxile; cells often unsymmetrical, varying from wide open to nearly closed; core lines slightly clasping. *Carpels* broadly ovate, elongated, slightly emarginate. *Seeds* numerous, compactly filling the cells, medium to large, variable in shape, rather short, very wide, plump, obtuse or sometimes acute, light brown.

Flesh nearly white, slightly tinged with yellow, not very firm, moderately fine, not crisp, tender, juicy, subacid mingled with sweet, fair to good in quality.

Season October to December.

GREEN SEEK-NO-FURTHER.

REFERENCES. 1. Coxé, 1817:131. fig. 2. Kenrick, 1832:53. 3. Thomas, 1849:181. fig. 4. Emmons, *Nat. Hist. N. Y.*, 3:38. 1851. fig. 5. Elliott, 1854:137. 6. Warder, 1867:720. 7. Downing, 1869:202.

SYNONYMS. *Autumn Seeknofurther* (4). *Bracy's Seek-no-further* (5). *Flushing Seek-no-further* (7). GREEN SEEK-NO-FURTHER (3, 5, 6, 7). SEEKNOFURTHER (4). *Seeknofurther* (5, 7, of Coxé 3). SEEK-NO-FURTHER (1). *White Seek-no-further* (7). WINTER SEEK-NO-FURTHER (2).

A large yellowish-green apple with faint blush of orange-red, very good in quality; season early winter. The tree is a rather slow grower, but eventually forms a regular, compact head and is quite productive. Desirable for the home orchard (1, 7.)

Historical. The earliest description of this variety which we find is that given by Coxé (1) who remarks that it is a native of one of the eastern states. Downing states that it originated in the garden of William Prince, Flushing, N. Y. (7). It is now seldom found in cultivation in New York.

FRUIT.

Fruit large. *Form* roundish conic or a little inclined to oblate conic, ribbed. *Stem* short to medium, thick. *Cavity* large, acute to somewhat acuminate, deep, broad, more or less marked with faint greenish-russet. *Calyx* moderately large, closed or slightly open. *Basin* deep, rather wide, abrupt, furrowed and wrinkled.

Skin moderately thick, tough, greenish-yellow or yellow with faint orange-red blush. *Dots* variable, large and small, often irregular, areolar with russet center or whitish and submerged.

Calyx tube large, moderately long, wide, conical. *Stamens* median.

Core small, axile; cells symmetrical, closed; core lines meeting or slightly clasping. *Carpels* roundish to broadly ovate, tufted. *Seeds* rather numerous, above medium, narrow, long, acute to acuminate, tufted.

Flesh yellowish-white, moderately coarse, crisp, tender, very juicy, sprightly, rich subacid, very good.

Season October to January.

GROSH.

REFERENCES. 1. *Western Horticultural Review*, 1853 (cited by 4). 2. *Gard. Monthly*, 1861:124 (cited by 3). 3. Kenrick, *Mag. Hort.*, 29:73. 1863. fig. 4. Warder, 1867:464, 735. fig. 5. Warder, 1867:720. 6. Downing, 1869:205. 7. *Ib.*, 1869:89. 8. *Ib.*, 1872:37 app. 9. *Am. Pom. Soc. Cat.*, 1877:16. 10. Barry, 1883:356. 11. Thomas, 1885:528. 12. Bailey, *An. Hort.*, 1892:252. 13. *Ill. Sta. Bul.*, 45:345. 1896. 14. *Am. Pom. Soc. Cat.*, 1897:13. 15. Lyon, *Mich. Sta. Bul.*, 143:200. 1897. 16. Van Deman, *Rural N. Y.*, 58:722. 1899. fig. 17. Ragan, *U. S. B. P. I. Bul.*, 56:132. 1905.

SYNONYMS. BEAUTY OF THE WEST (7, 13). *Beauty of the West* (17). *Big Rambo* (4, 8, 16, 17). *Cummings Rambo* (17). *English Rambo* (16). *French Rambo* of some (17). GROSH (5, 6, 14, 15, 16, 17). *Grosh* (9): *Grosh's Mammoth* (17). *Large Rambo* (17). *Large Summer Rambo* (17). *Lohringer Rambour* (17). *Mammoth Rambo* (17). *Monstrous Rambo* (17). *Musgrove* (16). *Musgrove's Cooper* (4, 7, 8, 17). *Naylor Rambo* (17). OHIO BEAUTY (2). *Ohio Beauty* (4, 7, 8, 16, 17). *Pickaway Rambo* (17). *Rambour Lorraine* (17). *Summer Rambo* (9, 17). *Sweet Rambo* incorrectly (17). WESTERN BEAUTY (3, 4, 8, 9, 10, 12). *Western Beauty* (7, 13, 16, 17, ? 14).

Fruit large, uniform and when well colored rather attractive, being mottled and striped with red. The flesh is tender, sprightly, pleasant subacid, in season from September till early winter. The tree is a strong grower, comes into bearing early and is a reliable cropper, yielding moderate to good crops almost annually. There is apt to be considerable loss from premature dropping of the fruit. So far as we can learn this variety has been as yet but little grown in New York. It appears to be worthy of further testing.

Summer Rambo much resembles this variety in general appearance but ripens about a month earlier.

Historical. Origin unknown. It was first brought to notice in Ohio, where it has been much grown under the name of Western Beauty.

TREE.

Tree vigorous with very long, moderately stout, curved branches. *Form* upright spreading to roundish, open. *Twigs* long to below medium, somewhat

curved, moderately stout; internodes medium or below. *Bark* clear reddish-brown mingled with olive-green, lightly streaked with scarf-skin; pubescent near tips. *Lenticels* clear brownish, conspicuous, quite numerous, medium or above, roundish, raised. *Buds* medium to large, prominent, broad, plump, obtuse, free or nearly so, slightly pubescent.

FRUIT.

Fruit large or very large, very uniform in size and shape. *Form* roundish oblate to somewhat conical, regular or sometimes elliptical. *Stem* short to above medium, moderately thick. *Cavity* acuminate to acute, deep, wide, often somewhat compressed, smooth and green or sometimes russeted, symmetrical or gently furrowed, sometimes lipped. *Calyx* above medium to very large, usually somewhat open disclosing the yellow calyx tube; lobes separated at the base, rather long, often leafy, acute to acuminate. *Basin* often oblique, large, medium in depth and width to deep and rather wide, abrupt, usually somewhat furrowed.

Skin thick, tough, waxy, greenish-yellow becoming clear bright yellow when fully mature, washed and mottled with bright red and striped and splashed with carmine, except in highly-colored specimens the yellow predominates. *Dots* numerous, small to rather large, pale gray or russet, often areolar or whitish and submerged.

Calyx tube rather large, wide at the top, conical or approaching funnel-form. *Stamens* nearly basal to above median.

Core small to medium, axile or somewhat abaxile with hollow cylinder in the axis; cells symmetrical, closed or partly open; core lines clasping. *Carpels* broadly roundish to ovate, emarginate, often tufted. *Seeds* moderately numerous, moderately dark brown, often abortive, medium size, moderately wide, obtuse to acute.

Flesh whitish, slightly tinged with yellow, rather firm, medium to rather coarse, crisp, tender, juicy, sprightly subacid, a little aromatic, good or sometimes very good.

Season September to January.

GRUNDY.

REFERENCES. 1. Hansen, S. D. *Sta. Bul.*, 76:56. 1902. 2. Budd-Hansen, 1903:95. 3. *Jewell Nursery Co. Cat.*, 1903:7. 4. Ragan, U. S. B. P. I. *Bul.*, 55:133. 1905.

SYNONYMS. GRUNDY (1, 2, 3, 4). *Thompson Seedling No.* 38 (1, 2, 3, 4).

Fruit large, regular, subacid, yellow marbled with red. Season September and October. Tree vigorous, spreading, productive. Originated from seed taken from New York to Grundy county, Iowa, by Mrs. J. S. B. Thompson in 1861. It has received favorable notice as a hardy variety in that region, but has not been sufficiently tested in this state to determine its value here (2).



HAAS

HAAS.

REFERENCES. 1. *Horticulturist*, 25:55. 1870. 2. Downing, 1872:10 app. fig. 3. Barry, 1883:345. 4. Thomas, 1885:512. 5. *Can. Hort.*, 11:69, 73. 1888. 6. Bailey, *Am. Hort.*, 1892:240. 7. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. 8. Stinson, *Ark. Sta. Bul.*, 43:103. 1896. 9. Munson, *Me. Sta. Rpt.*, 1896:70. 10. Waugh, *Vt. Sta. An. Rpt.*, 14:296. 1901. 11. Munson, *Me. Sta. Bul.*, 82:95. 1902. 12. Dickens and Greene, *Kan. Sta. Bul.*, 106:53. 1902. 13. Hansen, *S. D. Sta. Bul.*, 76:56. 1902. 14. Budd-Hansen, 1903:96. fig. 15. Farrand, *Mich. Sta. Bul.*, 205:44. 1903. 16. Beach and Clark, *N. Y. Sta. Bul.*, 248:123. 1904.

SYNONYMS. FALL QUEEN (2, 3, 7). *Fall Queen* (4, 5, 14, 16). *Gros Pomier* (2, 14). *Gros Pommier* (2). HAAS (5, 6, 9, 10, 11, 12, 13, 14, 15, 16). *Haas* (2, 3, 7). HASS (8). *Horse*, of some (2). Hoss (1). Huss (2).

Fruit of good medium size and very attractive bright red color but only fair in quality. With some growers it has proved profitable because the tree comes into bearing young and is very thrifty, hardy and productive, but on account of the inferior quality of its fruit doubtless it will be eventually wholly supplanted by better kinds.

Historical. Originated on the grounds of Gabriel Cerré, St. Louis, Mo. (2). It has been widely disseminated throughout the Middle West and Southwest where it is recognized as one of the hardiest of American apples. In New York it has been planted to a limited extent only and its cultivation is not increasing.

TREE.

Tree large, very vigorous with long, slender branches. *Form* at first comparatively tall and upright but becoming spreading or roundish. *Twigs* long, curved, slender with large terminal buds; internodes long. *Bark* brown or reddish-brown, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, medium size, oval, not raised. *Buds* medium to large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit medium to above. *Form* oblate a little inclined to conic, somewhat ribbed; sides usually unequal. *Stem* medium to short, thick, often partly red. *Cavity* acute to acuminate, deep, broad, usually symmetrical, more or less covered with thin greenish-russet. *Calyx* small to medium, closed or nearly so; lobes separated at base, short, narrow, acuminate. *Basin* moderately narrow, rather deep, abrupt, smooth or slightly furrowed and wrinkled.

Skin thin, tough, smooth, yellow, mottled, washed and nearly covered with deep bright red or brownish-red, striped and splashed with deep carmine. *Dots* small to rather large, inconspicuous, numerous, pale or russet. *Prevailing effect* red striped with carmine.

Calyx tube very variable, rather long and wide, conical or approaching funnel-form. *Stamens* median or below.

Core below medium to above, somewhat abaxile; cells symmetrical, open or sometimes closed; core lines clasping. *Carpels* broadly roundish to elliptical. *Seeds* dark brown, medium to large, of medium width, plump, acute.

Flesh white, often stained with red, firm, moderately fine, a little tough, moderately juicy to juicy, sprightly subacid, aromatic, a little astringent, poor to fair or sometimes nearly good.

Season October to early winter. In common storage the ordinary commercial limit is November (16).

HAGLOE.

REFERENCES: 1. Cox, 1817:107. fig. 2. Thacher, 1822:126. 3. Fessenden, 1828:131. 4. Downing, *Horticulturist*, 3:249. 1848. 5. Thomas, 1849:141. 6. Downing, 1854:146. 7. Elliott, 1854:137. 8. Hooper, 1857:42. 9. *Horticulturist*, 14:425. 1859. 10. *Am. Pom. Soc. Cat.*, 1862. 11. Warder, 1867:596. 12. Barry, 1883:334. 13. Bailey, *Am. Hort.*, 1892:240. 14. *Am. Pom. Soc. Cat.*, 1899:17. 15. Budd-Hansen, 1903:96.

SYNONYMS. HAGLOE (7, 8, 11, 12, 13, 14, 15). HAGLOE CRAB (1, 2, 3), but incorrectly. SUMMER HAGLOE (4, 5, 6, 9, 10). *Summer Hagloe* (7).

In 1817 Cox gave the following description of this variety (1). "The fruit, when fully ripe, has a yellow ground streaked with bright red—the size about middling, the form round, flat at the ends; the stalk large—the flesh remarkably soft and woolly, but not dry—the taste acid, but highly flavoured. * * * It ripens in August and September; keeps a long time without rotting—it bears abundantly and early; the growth of the tree is very uncommon; thick strong shoots; buds, particularly at the extremity of the branches, very large; the colour of the wood dark—the size of the tree small: the Hagloe is an uncommonly fine cooking apple; and from its great beauty and large size, added to its abundant bearing, is a valuable market fruit."

The tree is not a very good grower but comes into bearing rather young and yields moderate to good crops annually or nearly annually. The quality of the fruit is such that it is valued chiefly for culinary use and market. It is of good size and pretty uniform but the color is predominantly pale yellow rather faintly striped with red. It is not sufficiently attractive for a good market sort.

Historical. It is now held that Hagloe originated in America (14, 15). Cox and some later writers confused this variety with the English cider fruit known as Hagloe Crab but eventually this error was discovered (4) and the name Summer Hagloe came to be commonly accepted among pomologists for this variety, under which name it was listed by the American Pomological Society in 1862 (10). In 1899 the name was changed to Hagloe (14) in the catalogue of the American Pomological Society. This variety is but little known in New York.

TREE.

Tree rather small, a slow grower with moderately long, crooked branches. *Form* flat, spreading, rather dense. *Twigs* short, straight, stout with large

terminal buds; internodes medium. *Bark* brown with some olive-green, lightly mottled with scarf-skin; pubescent near tips. *Lenticels* scattering, large to medium, oblong, raised. *Buds* medium to large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit medium to large, pretty uniform. *Form* roundish truncate to roundish conic, ribbed; sides often unequal. *Stem* short to medium, rather thick. *Cavity* acute, medium in width and depth, symmetrical, russeted and with rather irregular, outspreading russet. *Calyx* below medium to above, closed or open; lobes often separated at the base, narrow, acuminate. *Basin* moderately deep, narrow to medium in width, abrupt, nearly smooth.

Skin rather tender, smooth, somewhat glossy, pale greenish-yellow washed and mottled with pinkish-red marked with splashes and narrow stripes of bright carmine. *Dots* light, inconspicuous. *Prevailing effect* yellow, faintly striped.

Calyx tube moderately long, wide, conical. *Stamens* basal.

Core usually small, axile to abaxile; cells usually open; core lines clasping. *Carpels* broadly ovate, emarginate, tufted. *Seeds* rather light brown, small to medium, roundish, very plump, obtuse.

Flesh white, moderately fine, tender, rather juicy, sprightly subacid, good for culinary purposes.

Season late August and September.

HARVEST REDSTREAK.

REFERENCES. 1. Downing, 1857:214. 2. Warder, 1867:436. 3. Downing, 1869:211. 4. Thomas, 1875:501. 5. Ragan, U. S. B. P. I. Bul., 56:138. 1905.

SYNONYMS. *Early Red Pippin* (5). *Early Redstreak* (2). *Early Red Streak* (3). HARVEST REDSTREAK (2, 5). HARVEST RED STREAK (1, 3, 4). *Striped Harvest* (3, 5).

Fruit of medium size, smooth, greenish-yellow or whitish striped and splashed with red. *Flesh* whitish, coarse, subacid, good for culinary use; season August and September. It is not sufficiently attractive in color for a good market variety. The tree is medium to large, with round head, moderately vigorous to very vigorous and yields good to heavy crops biennially.

Historical. This is an old variety of unknown origin. It is rarely found in New York and is now seldom or never planted.

HASKELL.

REFERENCES. 1. Manning, *Mag. Hort.*, 6:172. 1840. 2. *Ib.*, 7:45. 1841. 3. Thomas, 1849:145. fig. 4. Cole, 1849:108. fig. 5. *Cultivator*, 6:342. 1849. 6. Barry, 1851:284. 7. Elliott, 1854:137. 8. Gregg, 1857:43. 9. Downing, 1857:82. 10. Warder, 1867:385. 11. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 12. Bailey, *Am. Hort.*, 1892:240. 13. Hoskins, *Rural N. Y.*, 53:278. 1894. 14. Powell and Fulton, U. S. B. P. I. Bul., 48:44. 1903. 15. Beach and Clark, *N. Y. Sta. Bul.*, 248:124. 1904.

SYNONYMS. HASKELL (14, 15). HASKELL SWEET (1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13). *Haskell Sweet* (4, 14, 15). SASSAFRAS SWEET (4). *Sassafras Sweet* (6, 7, 8, 9).

Fruit large, of good appearance for a greenish-yellow apple, sweet, excellent for culinary use, in season from September to late fall or early winter. The crop does not ripen uniformly. The earliest ripening fruit becomes fully mature in September while at the same time others are green and hard. In ordinary storage the commercial limit appears to be early November and in cold storage the middle of January (15). The tree is a thrifty grower, comes into bearing moderately young and yields full crops biennially. Desirable for the home orchard.

Historical. Origin, Ipswich, Mass. (2, 5). It is not commonly known in New York. It is occasionally listed by nurserymen (12) but is now seldom planted.

TREE.

Tree large, vigorous; branches long, moderately stout with numerous small spurs. *Form* upright spreading or roundish, rather open. *Twigs* moderately long, curved, moderately stout; internodes long. *Bark* brown, heavily mottled with scarf-skin; pubescent. *Lenticels* quite numerous, small, round, not raised. *Buds* medium size, plump, obtuse, free, pubescent.

FRUIT.

Fruit large or above medium, uniform in size and shape. *Form* roundish or oblate, regular. *Stem* short, usually not exerted. *Cavity* acute to acuminate, deep or moderately deep, moderately narrow to rather wide, somewhat furrowed, russeted and with some out-spreading russet rays. *Calyx* large, closed. *Basin* wide, moderately deep, rather abrupt, smooth or somewhat wrinkled.

Skin greenish-yellow, more or less dotted and flecked with russet, occasionally with a bronze bluish. *Dots* numerous, large, dark.

Calyx tube large, cone-shape to funnel-form. *Stamens* median.


Core rather small, axile; cells symmetrical, closed; core lines clasping. *Carpels* broad at the middle narrowing toward base and apex, emarginate. *Seeds* medium size, short, plump, obtuse.

Flesh yellowish, a little coarse, moderately crisp, tender, very sweet, aromatic, very good in flavor and quality.

Season September to late fall or early winter.

HAWLEY.

REFERENCES. 1. Hovey, *Mag. Hort.*, 13:112, 535. 1847. fig. 2. *Cultivator*, 4:114. 1847. fig. 3. Leavenworth, *Horticulturist*, 2:27. 1847. fig. 4. *Cultivator*, 5:246. 1848. 5. Cole, 1849:112. fig. 6. Thomas, 1849:156.



HAWLEY

fig. 7. Emmons, *Nat. Hist. N. Y.*, 3:48. 1851. col. pl. No. 24, fig. 8. Hovey, 2:39. 1851. col. pl. and fig. 9. Barry, 1851:284. 10. *Horticulturist*, 7:484. 1852. 11. *Am. Pom. Soc. Cat.*, 1852. 12. *Mag. Hort.*, 19:68. 1853. 13. Elliott, 1854:137. 14. *Mag. Hort.*, 20:241. 1854. 15. Downing, 1857:82. 16. Gregg, 1857:44. fig. 17. Hooper, 1857:43. 18. *Horticulturist*, 13:481. 1858. 19. Warder, 1867:410. fig. 20. Leroy, 1873:375. fig. 21. Bailey, *Am. Hort.*, 1892:240. 22. Waugh, *Vt. Sta. An. Rpt.*, 14:296. 1901.

SYNONYMS. *Douse* (3, 5, 8, 10, 13, 20). *Dows* (1, 3, 8, 13, 20). *Dowse* (2, 4, 6, 9, 10, 16, 20, 22). HAWLEY (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22). *Howley* (20).

Hawley is a very large apple of the Fall Pippin type in color, size and form. When the color is fully developed it is a handsome yellow. Season September and October. It is of delicious dessert quality and desirable for the home orchard but not well adapted for market because the tree is not very productive and the fruit often is scabby and sometimes it water-cores and rots at the core. The tree is a moderate grower in the nursery but in the orchard it is rather vigorous, medium in size to large, hardy and rather long-lived. It does not come into bearing very young. When mature it bears quite regularly but is usually a light or moderate cropper.

Historical. Originated on the farm of Mr. Mathew Hawley, New Canaan, N. Y., about 1750, from seeds which Mr. Hawley obtained from Milford, Conn. (1, 3). The original tree lived nearly a century. The variety gradually became disseminated throughout New York state. It has long been known in cultivation in different parts of the state, particularly in Columbia, Onondaga, Cayuga, Tompkins, Seneca and Monroe counties. It is now rarely listed by nurserymen and is seldom planted.

FRUIT.

Fruit large or very large, pretty uniform in size and shape. *Form* nearly globular to somewhat oblate or slightly conic, sometimes inclined to elliptical, more or less distinctly ribbed. *Stem* medium in length, rather slender. *Cavity* acute to nearly obtuse, deep, wide and with outspreading russet rays. *Calyx* below medium to rather large, partly closed; lobes often separated at base, reflexed, wide, acute. *Basin* moderately deep to deep, medium to wide, very abrupt, often decidedly furrowed.

Skin fair, smooth, waxy, rather thin, pale green deepening to yellow as it matures, sometimes showing a faint brownish blush, with scattering russet dots and flecks especially toward the cavity.

Calyx tube large, wide, cone-shape, yellow or brownish.

Core below medium to above; cells closed; core lines meeting. *Carpels* rather flat, tufted, roundish, emarginate. *Seeds* few, obtuse. When well developed they are medium in size, but often some are abortive.

Flesh tinged with yellow, soft, very tender, rather fine-grained, juicy, rich, mild subacid, very good especially for dessert.

Season September to November or later.

HAWTHORNDEN.

REFERENCES. 1. Hooker, *Pom. Lond.*, 1813:T. 44 (cited by 2). 2. Forsyth, 1824:106. 3. *Pom. Mag.*, 1828:No. 34. col. pl. 4. *London Hort. Soc. Cat.*, 1831:No. 530. 5. *Mag. Hort.*, 1:326. 1835. 6. Manning, 1838:48. 7. *Ib.*, *Mag. Hort.*, 7:45. 1841. 8. Downing, 1845:86. 9. Cole, 1849:113. 10. Hovey, *Mag. Hort.*, 17:18. 1851. fig. 11. Barry, 1851:284. 12. Emmons, *Nat. Hist. N. Y.*, 3:40. 1851. 13. *Mag. Hort.*, 19:174. 1853. 14. Elliott, 1854:171. 15. Gregg, 1857:43. 16. Hooper, 1857:43. 17. *Am. Pom. Soc. Cat.*, 1862. 18. Warder, 1867:410. fig. 19. Leroy, 1873:376. fig. 20. Thomas, 1875:501. 21. *Montreal Hort. Soc. Rpt.*, 1:10. 1876. 22. *Ib.*, 5:24. 1879. 23. Hogg, 1884:105. 24. *Rural N. Y.*, 45:233. 1886. figs. 25. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 26. Bailey, *Am. Hort.*, 1892:241. 27. Gaucher, 1894:No. 13. col. pl. 28. Dempsey, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. fig. 29. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354.

SYNONYMS. *Hawley* (27) but erroneously. *HAWTHORNDEN* (2, 6, 7). *Hawthorndean* (23). *HAWTHORNDEN* (1, 3, 4, 5, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29). *Lincolnshire Pippin* (27). *Lord Kingston* (27). *Maiden's Blush* erroneously (10). *Old Hawthorndean* (23, 27). *Pomme de Hawthornden* (27). *Red Hawthornden* (10). *Shoreditch White* (27). *Weisser Hawthornden* (27). *Wheeler's Kernel* (27). *White Apple* (27). *White Hawthorndean* (23, 27). *White Hawthornden* (3, 4, 8, 10, 12, 14).

A Scotch variety which has done fairly well in various parts of this country from New England to California. Warder (18) says of it: "This famous Scotch fruit appears to do very well in this country, but it must yield the palm to its American cousin and representative, the Maiden's Blush, which possesses all its good qualities as a market and kitchen fruit, with attractive appearance."

The tree is said to be a vigorous grower and a biennial cropper, and the fruit is above medium to large, regular, fair, white sometimes nearly overspread with faint blush that deepens to bright red in the sun. *Flesh* mild subacid, not of high flavor, pleasant, good. *Season* September and October.

We are not acquainted with this variety. It appears to be but little known among New York fruit growers. It is still occasionally listed by nurserymen (26).

HIBERNAL.

REFERENCES. 1. Budd, *Ia. Hort. Soc. Rpt.*, 1880:525. 2. Gibb, *Montreal Hort. Soc. Rpt.*, 1881:156. 3. Tuttle, *Ib.*, 8:136. 1881-82. 4. *Ib.*, 1883:98. 5. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:440. 6. Budd, *Ia. Agr. Coll. Bul.*, 1885:15. 7. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:15. 8. Schroeder, *Ib.*, 1886-87:71. 9. Craig, *Ib.*, 1886-87:103. 10. Budd, *Ia. Agr. Coll. Bul.*, 1890:20. 11. *Can. Hort.*, 13:216. 1890. 12. Budd, *Ia. Sta. Bul.*, 19:537. 1892. 13. Bailey, *Am. Hort.*, 1892:241. 14. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57, 58. 15.

Freeborn, *Nat. Nurseryman*, 1894:132. 16. *Can. Hort.*, 17:7. 1894. 17. *Gard. and For.*, 8:340. 1895. 18. Munson, *Me. Sta. Rpt.*, 1896:74. 19. Waugh, *Vt. Sta. Bul.*, 61:30. 1897. 20. *Am. Pom. Soc. Cat.*, 1897:13. 21. Troop, *Ind. Sta. An. Rpt.*, 12:80. 1899. 22. Waugh, *Vt. Sta. An. Rpt.*, 14:296. 1901. 23. Macoun, *Can. Dept. Agr. Bul.*, 37:39. 1901. 24. Hansen, *S. D. Sta. Bul.*, 76:57. 1902. *fig.* 25. Munson, *Me. Sta. An. Rpt.*, 1902:83, 87. 26. Budd-Hansen, 1903:98. *fig.*

SYNONYMS. HIBERNAL (1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26). *Hibernal* (8). No. 378 (1, 2, 4, 5, 6, 10, 12, 15, 18, 25, 26). *Orsimui* (4, 5, 6). OSMOE (8). *Romna* (23).

A Russian variety which is proving valuable in portions of the Upper Mississippi valley and the Northwest because of its ability to withstand the rigorous climatic conditions of those regions. Hansen says of it: "This variety represents what is probably the hardiest type of the Russian race of apples; there are several sorts closely resembling, or identical with, *Hibernal*. Tree vigorous, very spreading, productive. The strong spreading growth makes it especially desirable as a stock for top-grafting, probably the best we have at the present time. Fruit large, irregular, oblate to roundish oblate conical; skin thick; surface greenish-yellow, with a dull bronze mixed red on sunny side, with a few dull crimson splashes; dots white, minute, obscure, often some large russet dots; cavity regular, medium deep, with a large patch of russet radiating out irregularly over nearly the entire base, this is a marked characteristic; stem medium, often short; basin narrow, rather shallow, wrinkled; calyx half open or open. Core closed, meeting; tube funnel-shaped; stamens median; seeds few; flesh acid, with some astringency, juicy, good for cooking. Early winter" (24, 26).

Macoun reports "Flesh yellowish, crisp, tender, juicy, acid; core small; quality above medium; season September to November. Tree very hardy, a strong, spreading grower, and very productive. Although not a good dessert fruit this is a fine cooking apple and on account of its great hardness and productiveness is one of the best of the Russian apples" (23).

HICKS.

REFERENCES. 1. Hicks, *Horticulturist*, 21:333. 1866. *fig.* 2. Downing, 1869:215. 3. Burrill and McCluer, *Ill. Sta. Bul.*, 45:308, 316, 324. 1896. 4. Thomas, 1897:639.

SYNONYMS. *Buckram* (2, 3). *HICKS* (1, 2, 3, 4).

A sweet apple of medium size, yellowish striped and splashed with crimson; season middle of August. It is but little known except in certain localities on Long Island. Not recommended for general planting in this state.

Historical. Hicks originated as a chance seedling and was brought to notice by Isaac Hicks, North Hempstead, L. I. (1, 2).

HIGHTOP SWEET.

REFERENCES. 1. Thacher, 1822:128. 2. Hovey, *Mag. Hort.*, 14:390. 1848. 3. Cole, 1849:97. 4. Phoenix, *Horticulturist*, 4:472. 1850. 5. Emmons, *Nat. Hist. N. Y.*, 3:16. 1851. *fig.* 6. Elliott, 1854:139. 7. *Am. Pom. Soc.*

Cat., 1856. 8. *Mag. Hort.*, 22:181. 1856. 9. Downing, 1857:151. 10. Gregg, 1857:40. 11. Warder, 1867:553. 12. *Ill. Hort. Soc. Rpt.*, 1869:33. 13. Fitz, 1872:145. 14. Thomas, 1875:189. 15. Downing, 1881:11 index, app. 16. Barry, 1883:333. 17. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 18. Bailey, *Am. Hort.*, 1892:250. 19. Munson, *Me. Sta. Rpt.*, 1893:132. 20. Hoskins, *Rural N. Y.*, 1894:248. 21. Budd-Hansen, 1903:99. fig.

SYNONYMS. *Early Sweet* (15). *HIGH TOP SWEET* (19). *High Top Sweet* (13). *HIGHTOP Sweet* (21). *HIGH TOP SWEETING* (1, 5, 6, 8, 12, 15). *High Top Sweeting?* (2, 3). *SUMMER SWEET* (3). *Summer Sweet* (5, 6, 9, 14). *SUMMER SWEETING* (2). *SWEET JUNE* (4, 10, 13, 18). *Sweet June* (6, 9, 11, 14, 16, 19, 20, 21).

Fruit of medium size; flesh yellowish, very sweet, rich and of very good quality; season July and August. Tree upright, vigorous, very productive.

Historical. In 1822 Thacher (1) remarked: "This tree, it is believed, is peculiar to the old Plymouth colony. The first settlers, either from choice, or for want of other varieties, cultivated it more generally than any other apple. It is now much on the decline. The fruit is under the middle size; of a yellowish colour, pleasant taste; but chiefly used for baking, and for drying. It is ripe in August, and is not long preserved. The tree is remarkable for its long upright stem."

It appears that this variety was introduced into Ohio from Connecticut and Massachusetts and afterward disseminated westward under the name of Sweet June. In 1892 Bailey (18) found that although various nurserymen were offering Sweet June for sale none of them mentioned Hightop Sweet. Some have held that the Sweet June of the West is not identical with the Hightop Sweet of Massachusetts (12). We have not had the opportunity of determining whether this is true, but if they are identical it appears strange that the name Hightop Sweet should be entirely dropped by those who are propagating it in the West. This variety is but little known in New York.

HILAIRE.

REFERENCES. 1. *1a. Hort. Soc. Rpt.*, 1879:453. 2. *Montreal Hort. Soc. Rpt.*, 1886 87:95. 3. Bailey, *Mich. Sta. Bul.*, 31:54. 1887. 4. Downing, 1881:102 app. fig. 5. Waugh, *Vt. Sta. Bul.*, 83:87. 1900.

SYNONYMS. *Cabane du Chien* (3, 5, 6). *Fameuse Baldwin*, of some (5). *St. HILAIRE* (1, 2, 3, 4, 5, 6).

An apple which resembles Fameuse in the color of its skin and in the color and texture of its flesh, but the flesh has more of a sprightly acid flavor and the fruit keeps better than that of Fameuse. Waugh reports (5) that it is not now grown in the vicinity of its origin. Professor U. P. Hedrick of the Michigan Agricultural College, who supplied the fruit for the following description, states that as grown in Michigan the variety is hardy, productive and gives promise of being a valuable acquisition.

Historical. This is said to have originated in the orchard of Alexis Dery, Quebec (4). Probably a seedling of Fameuse (2). So far as we know it is not grown in New York.

TREE.

Tree large, vigorous; a heavy alternate bearer (2). *Twigs* short, curved, slender; internodes short. *Bark* dark brown, with light coat of streaked scarf-skin, slightly pubescent. *Lenticels* scattering, small, oblong, raised. *Buds* small, plump, acute, free, slightly pubescent.

FRUIT.

Fruit medium to above, uniform in size. *Form* oblate to roundish-oblate, rather irregular. *Stem* medium, moderately slender. *Cavity* acuminate or acute, moderately deep and broad, not russeted, symmetrical. *Calyx* medium, usually closed; lobes broad, obtuse. *Basin* medium in depth to shallow, moderately wide to rather narrow, rather abrupt, slightly wrinkled, symmetrical.

Skin thin, tender, smooth, pale yellow or whitish almost completely overspread with attractive red of the Fameuse hue becoming as highly colored as the Fameuse or McIntosh and covered with faint bloom; stripes obscure if any. *Dots* very numerous, small, red, sometimes gray or russet. *Prevailing effect* brilliant deep pinkish-red deepening to purplish-red.

Calyx tube long, rather narrow, funnel-shape. *Stamens* median to basal.

Core nearly axile, small to medium; cells closed or partly open; core lines clasping the funnel cylinder. *Carpels* round, slightly emarginate. *Seeds* dark, numerous, medium to large, wide, obtuse to acute.

Flesh whitish sometimes tinged with red, fine, crisp, tender, juicy, sprightly subacid, good to very good.

Season November to January. A better keeper than Fameuse.

HILTON.

REFERENCES. 1. Downing, 1857:151. 2. Warder, 1867:721. 3. Thomas, 1875:502.

SYNONYMS. None.

This variety originated in Columbia county, N. Y. According to Downing (1) the tree is vigorous and productive: the fruit large, yellowish-green, subacid, excellent for culinary purposes. Season September and October.

So far as we can discover this variety is not now known in cultivation.

HOADLEY.

REFERENCES. 1. Goff, *Wis. Sta. An. Rpt.*, 11:347. 1894. 2 *Ib.*, *Am. Pom. Soc. Rpt.*, 1899:236.

SYNONYMS. None.

A variety of the Oldenburg type which much resembles Oldenburg except that its season is about a month later. It is decidedly attractive in general appearance and of good quality for culinary purposes. The tree is a moderate grower, comes into bearing early and so far as tested here is very productive. It appears to be worthy of testing where an apple of this type is desired.

Historical. Received from the Wisconsin Experiment Station in 1896 for testing at this Station.

TREE.

Tree moderately vigorous. *Form* upright spreading when young. *Twigs* short, straight, stout; internodes medium. *Bark* brown and reddish-brown, lightly streaked with scarf-skin, slightly pubescent. *Lenticels* scattering, medium size, round, not raised. *Buds* medium to large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit above medium to large, sometimes very large. *Form* roundish oblate inclined to conic, a little angular; sides unequal. *Stem* short, thick to slender. *Cavity* acute to obtuse, deep, broad, pretty symmetrical, thinly and irregularly russeted. *Calyx* rather large, closed or partly open; lobes long, broad, acute to obtuse. *Basin* deep to medium in depth, rather narrow to moderately wide, abrupt, slightly furrowed.

Skin moderately thick, tough, attractive yellow or greenish-yellow largely overspread with rather light bright red, mottled and irregularly striped and splashed with carmine. *Dots* inconspicuous, small, submerged, pale.

Calyx tube variable, short, rather wide, funnel-shape, sometimes broadly conical with core lines meeting. *Stamens* median to nearly marginal.

Core medium size, abaxile; cells open; core lines meeting or slightly clasping. *Carpels* broadly cordate or elliptical, slightly tufted. *Seeds* medium or below, wide, moderately long, usually plump, rather obtuse, dark colored.

Flesh tinged with yellow, pretty firm, a little coarse, crisp, tender, very juicy, brisk subacid, good.

Season late September to November.

HOG ISLAND SWEET.

REFERENCES. 1. Downing, 1857:152. 2. Warder, 1867:721. 3. Downing, 1872:10 index, app. 4. Thomas, 1875:502. 5. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292.

SYNONYMS. HOG ISLAND SWEET (1, 2, 3, 4, 5). *Sweet Pippin* (1). *Van Kleeck's Sweet* (3).

A beautiful and excellent sweet apple in season from September to early winter. It is an old variety which had its origin on Hog Island, near Long Island, N. Y. (1). It is but little known. The tree is vigorous and productive.

FRUIT.

Fruit medium to large, pretty uniform. *Form* roundish conical to oblate, broad at the base, regular or obscurely ribbed. *Stem* short to medium, moderately thick. *Cavity* acute, deep, rather broad, heavily russeted and with outspreading russet rays. *Calyx* medium to large, closed or partly open; lobes short, broad, acute. *Basin* shallow to medium in depth, narrow to moderately wide, abrupt, broadly furrowed.

Skin thick, rather tough, somewhat roughened with flecks and patches of russet, pale yellow or greenish washed and mottled with red overlaid with



HOADLEY

broad and narrow stripes and splashes of carmine. *Dots* numerous, small, russet. *Prevailing effect* bright striped red over a yellow background.

Calyx tube long, wide, conical to funnel-shape. *Stamens* basal to median.

Core small to medium, abaxile to nearly axile; cells somewhat unsymmetrical, open; core lines slightly clasping the funnel cylinder or meeting. *Carpels* roundish obovate to elliptical, emarginate. *Seeds* medium or below, sometimes tufted, rather wide, rather short, plump, acute to somewhat obtuse.

Flesh tinged with yellow, moderately coarse, crisp, tender, juicy, very sweet, somewhat aromatic, good to very good.

Season September to early winter.

HOLLAND PIPPIN.

REFERENCES. 1. Cox, 1817:109. fig. 2. Downing, 1845:86. 3. *Ib.*, *Horticulturist*, 3:345. 1848. 4. Thomas, 1849:156. fig. 5. Cole, 1849:110. 6. Downing, Chas., *Horticulturist*, 8:196. 1853. 7. Elliott, 1854:138. 8. Hovey, *Mag. Hort.*, 22:555. 1856. fig. 9. Hooper, 1857:45. 10. Gregg, 1857:37. 11. *Am. Pom. Soc. Cat.*, 1862. 12. Warder, 1867:506. 13. Wickson, 1889:244. 14. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 15. Bailey, *Am. Hort.*, 1892:241. 16. Beach and Clark, *N. Y. Sta. Bul.*, 248:124. 1904.

SYNONYMS. FALL PIPPIN (1). *French Pippin* (7). HOLLAND PIPPIN (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16). *Pie Apple* (2, 7, 8). *Reinette d'Hollande* (2). *Reinette d'Holland* (7). *Summer Pippin* (2, 7, 8).

There are two varieties in cultivation in New York under the name Holland Pippin. One is a winter apple in season from late autumn to April or May which has already been described under the name Holland Winter, Vol. I., page 159. The other begins to ripen earlier than Fall Pippin and is in season during September and October. This variety was formerly confused by some with the Fall Pippin. The following comparison of the two varieties was given by Downing in 1848 (3).

"The Holland Pippin, though considerably resembling this apple in the growth of the tree, and size and shape of the fruit, is a totally distinct apple from the Fall Pippin. In fact, while the Fall Pippin is one of the best autumn table apples (at least in this district), the Holland Pippin is of very inferior quality for dessert, and is, in fact, only a *cooking* apple. As a kitchen fruit, however, it is one of the most valuable *summer* fruits we know — for it bears regularly and well, comes into use at the beginning of August, and continues fit for pies, tarts, and sauce, *until* October, when the Fall Pippin *begins* to ripen. The Holland Pippin is fit for use while

the skin is quite green, but the Fall Pippin, not until it turns quite yellow. Finally, the stalk of the Holland Pippin is short, and set in a wide cavity, while that of the Fall Pippin is large, and set in a cavity often narrow, and comparatively shallow. With these points of difference, these two apples ought not to be confounded."

Holland Pippin is grown to a limited extent for market. It appears to be more valued for this purpose in certain portions of the Hudson valley than in other sections of the state. The fruit is large and when kept free from scab its general appearance is good for a green apple. The crop ripens unevenly. Some of the apples ripen early and are correspondingly short-lived while others ripen later and keep correspondingly later. It varies greatly in keeping qualities in different seasons, some years keeping well till late fall or early winter (16). The tree is a good grower, hardy or nearly so, healthy, pretty long-lived and generally quite productive yielding moderate to heavy crops biennially or sometimes annually.

Historical. Origin unknown. It is an old variety which has long been in cultivation in this and adjoining states. It is still listed by nurserymen but it is not being planted to any considerable extent.

TREE.

Tree large or moderately large, vigorous. *Form* spreading or roundish. *Twigs* medium to long, curved, stout; internodes medium. *Bark* dark brown, heavily coated with gray scarf-skin; pubescent. *Lenticels* scattering, small to medium, oval, not raised. *Buds* medium size, plump, obtuse, free, pubescent.

FRUIT.

Fruit usually large or very large, sometimes medium, pretty uniform in size and shape. *Form* roundish often decidedly flattened at the end varying to oblate conic, obscurely ribbed. *Stem* medium to short, usually rather slender. *Cavity* acute or sometimes acuminate, medium in width to broad, moderately shallow to deep, usually covered with thick outspreading russet. *Calyx* pubescent, medium to small, closed or partly open; lobes rather long, acute. *Basin* usually rather shallow but varying to moderately deep, medium in width to rather narrow, abrupt to somewhat obtuse, ridged and wrinkled.

Skin thin, tough, nearly smooth, rather pale yellow or greenish with more or less of a brownish-red blush which is conspicuously marked with large, irregular, areolar dots. *Dots* numerous, large and small, often submerged and greenish.



HOOK

Calyx tube wide, broadly conical to truncate funnel-form. *Stamens* below median to basal.

Core medium to large, abaxile; cells unsymmetrical, open; core lines meeting or somewhat clasping. *Carpels* broad, narrowing toward base and apex, slightly emarginate, a little tufted. *Seeds* medium size, rather narrow, acute to somewhat acuminate.

Flesh nearly white, medium to slightly coarse-grained, moderately crisp, rather tender, very juicy, brisk subacid, good for culinary uses.

Season September and October.

HOOK.

REFERENCE. 1. (?) *Mich. Pom. Soc. Rpt.*, 1880:183.

SYNONYMS. None.

Fruit of good medium size, rather attractive pale yellow with tender flesh of mild subacid flavor. It is in season during October and November. It is especially esteemed for dessert use.

Historical. This variety was received from Schoharie county, N. Y., where it is well known and has the reputation of being one of the most desirable dessert apples of its season. We have not yet been able to determine its origin, nor the extent of its distribution.

FRUIT.

Fruit medium size, pretty uniform in shape and size. *Form* roundish ovate or inclined to oblong conic, sometimes roundish truncate. *Stem* short, rather slender. *Cavity* acuminate, moderately shallow to deep, broad, often lipped, smooth or with some outspreading russet rays. *Calyx* medium size, usually somewhat open; lobes rather narrow to wide, acute to acuminate. *Basin* medium in width and depth, obscurely furrowed and wrinkled.

Skin smooth, clear yellow or greenish-yellow. *Dots* inconspicuous, minute, usually submerged.

Calyx tube large, conical or approaching funnel-form. *Stamens* median or below.

Core rather small, axile; cells symmetrical, closed or a little open; core lines meeting or somewhat clasping. *Carpels* smooth, flat, obcordate to elliptical, emarginate. *Seeds* medium brown, rather large, wide, flat, obtuse.

Flesh whitish with slight tinge of yellow, very tender, fine-grained, juicy, mild subacid, somewhat aromatic, very good.

Season October and November.

HOWARD BEST.

REFERENCES. 1. Bailey, *Mich. Sta. Bul.*, 31:52. 1887. 2. *N. Y. Sta. An. Rpt.*, 11:223. 1892.

SYNONYMS. HOWARD'S BEST (2). HOWARD'S BEST RUSSIAN (1).

This is an attractive apple which bears a decided resemblance to Alexander in form, color and quality. The tree is a moderate grower, comes into bearing rather young and yields moderate to good crops almost annually. We have not yet determined whether or not it is superior to Alexander.

Historical. Received in 1892 for testing at this Station from C. G. Patten, Charles City, Ia. (2).

TREE.

Tree moderately vigorous with short, moderately stout, crooked branches. *Form* rather flat and spreading, inclined to droop. *Twigs* short, straight, slender with large terminal buds; internodes short to medium. *Bark* dull brown tinged with olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* scattering, small, oblong, not raised. *Buds* small, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit above medium, usually large to very large. *Form* oblate conic, rather flat at the base, somewhat ribbed, symmetrical. *Stem* medium to long, thick. *Cavity* acute, deep, medium in width to wide, symmetrical, sometimes compressed, heavily russeted. *Calyx* large, open; lobes separated at base, short, narrow, acute. *Basin* moderately shallow to rather deep, rather narrow, abrupt, distinctly furrowed to nearly smooth.

Skin rather tough, smooth, waxy, light yellowish-green, mottled, striped and splashed with bright, dark red over a large part of the surface. *Dots* indistinct, medium size, gray, scattering.

Calyx tube short, wide, conical or approaching funnel-form. *Stamens* basal to median.

Core medium size, nearly axile; cells closed or slightly open; core lines clasping or nearly meeting. *Carpels* very broadly ovate or inclined to elliptical, slightly emarginate, tufted. *Seeds* medium to small, wide, short, rather flat, obtuse.

Flesh slightly tinged with yellow, firm, rather coarse, tender, very juicy, sprightly subacid, fair to good.

Season September and October.

HUNTER PIPPIN.

REFERENCE. 1. Downing, 1876:54 app.

SYNONYMS. None.

A chance seedling which originated in Westport, N. Y. Downing describes the tree as moderately vigorous and a good annual bearer; fruit medium size, whitish-yellow; flesh brisk subacid; season August (1).

So far as we can learn this variety is not being propagated.

HURLBUT.

REFERENCES. 1. Cole, 1849:118. *fig.* 2. Emmons, *Nat. Hist. N. Y.*, 3:31. 1851. *col. pl.* and *fig.* 3. Downing, 1857:155. 4. *Am. Pom. Soc. Cat.*, 1862. 5. Warder, 1867:722. 6. Thomas, 1875:201. 7. Barry, 1883:347. 8. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 9. Bailey, *Am. Hort.*, 1892:241. 10. Mun-

son, *Me. Sta. Rpt.*, 1893:133. 11. Dempsey, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. 12. Burrill and McCluer, *Ill. Sta. Bul.*, 45:326. 1896. 13. Munson, *Me. Sta. An. Rpt.*, 18:89. 1902. 14. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:45. 1903. 15. Beach and Clark, *N. Y. Sta. Bul.*, 248:126. 1904.

SYNONYMS. HURLBURY (2). HURLBUT (1, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15). *Hurlbut Stripe* (3). HURLBUTT (5).

The general appearance of Hurlbut is good yet it is not particularly attractive either in size or color nor does it take first rank for either home use or market. The tree is a strong grower, comes into bearing moderately young and yields heavy crops biennially. Because of the tendency of the tree to overproduction in bearing years the size of the fruit is in many cases reduced and a considerable portion of it is too small to be marketable. Hurlbut has proved to be a profitable variety with some fruit growers but although it has long been known in cultivation and has been quite widely disseminated it has nowhere gained prominence as a commercial variety.

Historical. In 1849 Cole (1) wrote: "The original tree is still flourishing on the farm of General Leonard Hurlbut, Winchester, Conn." It is still quite frequently listed by nurserymen (9) but is not being planted to any considerable extent.

TREE.

Tree rather large, vigorous or moderately vigorous. *Form* spreading or roundish and somewhat inclined to droop. *Twigs* moderately long, slightly curved, moderately stout to rather slender; internodes below medium to short. *Bark* dark brown to clear brownish-red, heavily mottled with scarf-skin; slightly pubescent. *Lenticels* scattering, small, round, slightly raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit below medium to above, fairly uniform in size and shape. *Form* roundish, slightly oblate or inclined to oblate conic, somewhat angular, rather symmetrical. *Stem* medium to short, rather slender. *Cavity* acute, shallow to medium in depth, medium in width, symmetrical or compressed, usually covered with heavy outspreading russet. *Calyx* medium or below, usually closed; lobes medium to long, narrow, acute. *Basin* shallow to medium in depth, narrow to nearly medium in width, somewhat abrupt, smooth or slightly wrinkled.

Skin thick, tough, smooth, greenish-yellow largely overspread with brownish-red or dull red, splashed and striped with carmine. *Dots* scattering, inconspicuous, usually submerged, sometimes russet.

Calyx tube very short, wide, truncate conical with fleshy pistil point projecting into the base. *Stamens* marginal.

Core medium size, axatile; cells wide open to nearly closed; core lines

meeting or slightly clasping. *Carpels* nearly round. *Seeds* numerous, rather large, moderately wide, long, rather plump, acute.

Flesh white or yellowish, moderately firm, rather fine, tender, crisp, very juicy, aromatic, mildly subacid, good to very good.

Season variable; October to December or January (15).

ISHAM.

REFERENCES. 1. Bailey, *Am. Hort.*, 1892:242. 2. Hoskins, *Rural N. Y.*, 53:310. 1894. 3. *Am. Pom. Soc. Cat.*, 1899:17. 4. Hansen, *S. D. Sta. Bul.*, 76:60. 1902. fig. 5. *Kan. Sta. Bul.*, 106:53. 1902. 6. Budd-Hansen, 1903:105.

SYNONYMS. ISHAM SWEET (1, 2, 3, 4, 5). ISHAM *Sweet* (6).

A red sweet apple of medium size, in season in late fall and early winter. It has been grown to some extent in some of the Western states but it has not been much tested in New York and it is doubtful whether it is desirable for planting in any portion of this state.

Historical. Isham originated from seed of Bailey *Sweet*. It was introduced about 1864 by F. K. Phoenix, Delavan, Wis. (2).

TREE.

Tree moderately vigorous. *Form* upright spreading. *Twigs* medium to long, rather stout, in some cases quite blunt at the tips, straight or nearly so; internodes rather long. *Bark* reddish-brown overlaid with heavy grayish scarf-skin, not pubescent or very sparingly so. *Lenticels* rather inconspicuous, rather scattering, irregular in size and shape, not raised. *Buds* large, prominent, fleshy, heavily pubescent, adhering to bark.

FRUIT (4, 6).

Fruit medium. *Form* roundish, slightly tapering. *Stem* short. *Cavity* regular, acute, with much radiating russet. *Calyx* open; segments flat, convergent. *Basin* very shallow, minutely wrinkled.

Skin yellowish-green mostly covered with brownish-red, solid and mixed on sunny side, striped and broadly splashed on the shady side. *Dots* distinct, russet, numerous, minute; a few large russet dots.

Calyx tube funnel-shape. *Stamens* median.

Core closed; cells round, entire. *Seeds* long, large, flat.

Flesh very yellow with yellow veinings, firm, very sweet, very good.

JACK.

REFERENCES. 1. (?) Hooper, 1857:46. 2. (?) Downing, 1869:206. 3. (?) *Ill. Sta. Bul.*, 45:334. 1896.

SYNONYMS. (*Early Jack* 1)? (JACK APPLE 1)? (OSKALOOSA 2, 3)?

Fruit of good medium size, yellow; flesh very tender, rich, mild subacid. It is highly esteemed for its excellent dessert quality but it is too tender to stand shipping very well and on account of its irregular shape and yellow

color it is not sufficiently attractive for market purposes. The tree is not a vigorous grower and has rather slender twigs.

Possibly this is identical with Oskaloosa which has Jack as a synonym, (2, 3) but we have been unable to obtain fruit of Oskaloosa and the available descriptions of that variety are so meager that it is impossible to determine whether or not it is identical with the variety described above.

Historical. Jack is grown to a very limited extent in East Bloomfield, Ontario county, N. Y. We have not obtained it from any other locality. We have been unable to learn where it originated, or whether it is the Jack mentioned by Hooper (1).

TREE.

Tree not very vigorous with moderately long, slender, crooked branches. *Form* at first upright spreading but becoming roundish and rather dense. *Twigs* long, straight, slender; internodes short. *Bark* brown or reddish-brown, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous, rather conspicuous, medium size, oblong, slightly raised. *Buds* small, plump, obtuse, appressed, slightly pubescent.

FRUIT.

Fruit medium or above, sometimes rather large, not uniform in shape or size. *Form* oblate varying to roundish oblate or to oblate conic, very irregular, obscurely ribbed, often with the sides somewhat furrowed and unequal. *Stem* short to medium, slender. *Cavity* acuminate, usually deep, medium in width, russeted, with concentric russet lines often extending beyond the cavity, sometimes lipped. *Calyx* medium size, usually open. *Basin* small to medium, rather shallow to moderately deep, narrow to medium in width, unsymmetrical, irregularly furrowed or nearly smooth.

Skin rather thin, waxy, glossy, attractive yellow with shades of green, sometimes faintly tinged with red and marbled with whitish scarf-skin over the base somewhat after the manner of Yellow Newtown. *Dots* mostly small and depressed mingled with a few that are larger, scattering and irregular with russet center.

Calyx tube medium in width and length, conical to somewhat funnel-form. *Stamens* median or below.

Core medium in size, somewhat abaxile; cells usually symmetrical, somewhat open; core lines clasping. *Carpels* elliptical, deeply emarginate. *Seeds* medium or below, wide, short, rather flat, obtuse, mingled with light and dark brown.

Flesh tinged with yellow, moderately firm, moderately crisp or breaking, very tender, moderately juicy, very mild subacid, very good for dessert.

Season October and November.

JARVIS.

REFERENCE. 1. N. Y. Sta. An. Rpt., 11:223. 1892.

SYNONYMS. Crandall Seedling (1). JARVIS (1). No. 25 (1).

Fruit large and when well colored partly overspread and striped with red; flesh tender, juicy, subacid, pleasant but not superior in flavor or

quality; season late September to early winter. It is possibly desirable for local market but it is not recommended for general cultivation.

Historical. The original tree is standing near Ithaca, N. Y., on land once owned by a Mr. Jarvis from whom the variety takes its name. It is grown to a limited extent in the vicinity of Ithaca but so far as we know is not cultivated in any other portion of the state. Received for testing at this Station in 1892 from C. B. Crandall.

JEFFERIS.

REFERENCES. 1. Thomas, 1849:149. 2. *Mag. Hort.*, 18:491. 1852. 3. *Am. Pom. Soc. Cat.*, 1854. 4. Elliott, 1854:139. 5. *Mag. Hort.*, 21:62. 1855. 6. Gregg, 1857:44. 7. Downing, 1857:83. *fig.* 8. Hooper, 1857:47. 9. *Mag. Hort.*, 24:109. 1858. 10. Hoffs, *N. A. Pom.*, 1860. *col. pl.* 11. *Horticulturist*, 17:104, 150. 1862. 12. Warder, 1867:440. *fig.* 13. Barry, 1883:338. 14. Van Deman, *U. S. Pom. Rpt.*, 1888:570. 15. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 16. Bailey, *Am. Hort.*, 1892:242. 17. Burrill and McCluer, *Ill. Sta. Bul.*, 45:309, 323. 1896. 18. Powell, *Del. Sta. Bul.*, 38:18. 1898. 19. Macoun, *Can. Dept. Agr. Rpt.*, 1901:96. 20. Beach, *Western N. Y. Hort. Soc. Rpt.*, 1901:76. 21. Budd-Hansen, 1903:105. 22. Bruner, *N. C. Sta. Bul.*, 182:21. 1903. 23. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 24. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:45. 1903. 25. Beach and Clark, *N. Y. Sta. Bul.*, 248:126. 1904.

SYNONYMS. *Everbearing* (20). *Grantham* (17). *JEFFERIES* (10, 12, 14, 17, 20). *JEFFERIS* (3, 4, 7, 13, 15, 16, 18, 19, 21, 22, 23, 24, 25). *JEFFRIES* (1, 2, 5, 6, 8, 9, 11).

Fruit of medium size, yellow, blushed and splashed with red; flesh tender, mild subacid, delicious. It begins to ripen in September and continues in season till early winter. Commercial limit October (25). It is an excellent variety for the home orchard but not for commercial planting because it ripens unevenly, is apt to be deficient in size and is not especially attractive in color. The tree is a moderately vigorous grower, hardy, healthy, comes into bearing moderately early and is a reliable cropper yielding full crops biennially.

Historical. Originated with Isaac Jefferies, Newlin township, Chester county, Pa. It was named after the originator by the Committee of the Pennsylvania Horticultural Society which awarded this variety the premium for the best seedling apple exhibited in 1848 (10).

TREE.

Tree medium size, moderately vigorous. *Form* upright to roundish, open. *Twigs* short, straight, slender; internodes long. *Bark* brown mingled with olive-green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels*



JEFFERIS

scattering, small, oblong, not raised. *Buds* small, plump, obtuse, free, pubescent.

FRUIT.

Fruit small to medium, very uniform in size and shape. *Form* roundish oblate often inclined to conic, regular or obscurely ribbed. *Stem* medium length, thick to moderately slender. *Cavity* acute to acuminate, medium in depth to deep, medium to broad, symmetrical, russeted but slightly if at all. *Calyx* small to medium, closed or partly open; lobes short, rather broad, acute. *Basin* moderately shallow to rather deep, moderately wide, somewhat abrupt, smooth or nearly so, symmetrical.

Skin thin, tough, greenish-yellow or pale yellow more or less blushed and mottled with moderately dull red overlaid with narrow splashes and stripes of carmine. *Dots* small, scattering, inconspicuous, submerged or russet.

Calyx tube narrow, conical to funnel-shape. *Stamens* marginal to median.

Core small, axile, or nearly so; cells slightly open; core lines somewhat clasping or meeting. *Carpels* elliptical to somewhat obovate, emarginate, sometimes tufted. *Seeds* numerous, medium to rather large, wide, long, flat, very irregular, obtuse.

Flesh yellowish-white, firm, fine, crisp, tender, very juicy, mild subacid, very good.

Season September to January.

JEFFERSON COUNTY.

REFERENCES. 1. *Horticulturist*, 10:254. 1855. fig. 2. Downing, 1857:156. 3. Warder, 1867:723. 4. *Am. Pom. Soc. Cat.*, 1873. 5. Thomas, 1875:201. 6. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292.

SYNONYMS. None.

This variety originated in Jefferson county, N. Y., hence its name (2). The tree is of medium size, moderately vigorous; form somewhat drooping; twigs rather slender. It comes into bearing young and is a reliable bearer producing some fruit nearly every year, alternating heavy with lighter crops. The fruit is yellow shaded and splashed with red, not very bright in color, in season during October and November. It is particularly suitable for dessert, the flesh being tender, rather firm, crisp, of good flavor and excellent quality but it is not regarded as a good market variety for there is apt to be a rather large amount of small, imperfect or otherwise unmarketable fruit and when the fruit does not color properly, as happens in many cases, it is of poor flavor. It was listed by the American Pomological Society in 1873. It has been sparingly disseminated in various parts of the country but is as yet little known. So far as we can learn it is not being planted in New York.

JENNETTING or JUNEATING.

This name has been applied by some to White Juneating. For a description of this variety together with Hogg's account of the derivation of the name the reader is referred to White Juneating, page 240.

JERSEY SWEET.

REFERENCES. 1. Downing, 1845:87. 2. *N. Y. Agr. Soc. Trans.*, 1846:190. 3. Cole, 1849:110. 4. Thomas, 1849:145. *fig.* 5. Barry, 1851:284. 6. Emmons, *Nat. Hist. N. Y.*, 3:25. 1851. 7. Waring, 1851:26. 8. Elliott, 1854:86. *fig.* 9. Hooper, 1857:48. 10. *Ib.*, 1857:107, 111. 11. *Am. Pom. Soc. Cat.*, 1862. 12. *Horticulturist*, 17:104, 150. 1862. 13. Warder, 1867:395. 14. Fitz, 1872:153. 15. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 16. Bailey, *Am. Hort.*, 1892:242. 17. Hoskins, *Rural N. Y.*, 53:278. 1894. 18. *Mich. Sta. Bul.*, 118:60. 1895. 19. *Ib.*, 143:200. 1897. 20. *Ib.*, 205:45. 1903. 21. Budd-Hansen, 1903:105. 22. Beach and Clark, *N. Y. Sta. Bul.*, 248:126. 1904.


SYNONYMS. *American* (13). JERSEY SWEET (5, 7, 10, 12, 13, 16, 18, 21). JERSEY SWEETING (1, 2, 3, 4, 6, 8, 9, 14, 17).

An early autumn apple of medium size. It does not always color well but under favorable conditions it is highly colored, rich in flavor, tender and excellent in quality for either dessert or culinary uses. It is one of the best of the sweet apples of its season for planting for home use in New York but it has proved unsatisfactory as a commercial sort because it ripens at a time when there is little demand for fruit of this kind, is not a good keeper, is apt to be scabby and does not always color well. The tree is hardy, moderately long-lived, comes into bearing young and bears nearly every year, yielding moderate to good or sometimes heavy crops. The fruit comes in season late in August or early in September and ripens in succession during a period of several weeks; often some portion may be kept till early winter, but its commercial limit in ordinary storage is September or early October (22).

Historical. Origin unknown. Elliott calls it an American variety (8). It is pretty well known in different parts of New York state, but is now rarely found except in old orchards. It is commonly listed by nurserymen (16) but is now seldom or never planted except occasionally for home use.

TREE.

Tree rather large, moderately vigorous to vigorous; branches long, moderately stout, filled with spurs. *Form* upright to roundish, open. *Twigs* moderately long, straight, slender; internodes long. *Bark* brown, lightly streaked with scarf-skin; pubescent near tips. *Lenticels* scattering, medium to small, oblong, not raised. *Buds* medium size, plump, obtuse, appressed, pubescent.



JERSEY SWEET

FRUIT.

Fruit medium size. *Form* roundish ovate inclined to conic or to oblate conic; sides unequal. *Stem* long to medium length, rather slender. *Cavity* acute, usually rather deep, varying to shallow, medium in width, occasionally lipped, sometimes slightly russeted. *Calyx* small, closed; lobes medium to long, narrow, acute to acuminate. *Basin* rather small, moderately shallow to rather deep, narrow to medium in width, somewhat abrupt, ribbed and wrinkled.

Skin thin, tender, at first greenish-yellow but becoming clear yellow washed and mottled with brownish-red and overlaid with narrow stripes of bright carmine. *Dots* inconspicuous, greenish, submerged.

Calyx tube narrow, conical to funnel-form, often with fleshy pistil point projecting into the base. *Stamens* median.

Core medium size or above, axile or nearly so; cells symmetrical, usually closed; core lines clasping the funnel cylinder. *Carpels* elongated ovate, tufted. *Seeds* medium to large, variable in length and width, acute to acuminate.

Flesh yellowish, moderately firm, fine, crisp, tender, juicy, sweet, good to very good.

Season September to December.

JUDSON.

REFERENCES. 1. *Am. Pom. Soc. Cat.*, 1899:17. 2. *Ib.*, *Rpt.*, 1901:40. 3. Hansen, S. D. *Sta. Bul.*, 76:62. 1902. fig. 4. Budd-Hansen, 1903:108. fig. SYNONYMS. JUDSON (1, 2, 3, 4). *Thompson's Seedling No.* 29 (3, 4).

Fruit large, green or yellowish, more or less covered with red, not especially attractive in appearance and only fair to good in quality. Season October to December. Not valuable enough to be worthy of trial in New York except perhaps in those districts where superior hardiness is particularly desirable.

Historical. Originated in Grundy county, Iowa, by J. S. B. Thompson.

TREE.

Tree moderately vigorous with short, moderately stout, somewhat drooping branches. *Form* open, roundish to spreading. *Twigs* above medium to short, somewhat curved, medium to stout, rather pubescent with large terminal buds; internodes medium or below. *Bark* brown or reddish-brown tinged with olive-green, heavily coated with gray scarf-skin; pubescent. *Lenticels* quite numerous, rather conspicuous, medium or below, round or irregularly elongated, not raised. *Buds* prominent, large to medium, broad, plump, obtuse, free or nearly so, pubescent.

FRUIT.

Fruit large or very large, fairly uniform in size but not in shape. *Form* roundish conical or a little inclined to oblong, indistinctly ribbed, irregular;

sides often unequal. *Stem* short to medium, thick. *Cavity* very acuminate, deep, moderately narrow to rather broad, somewhat furrowed, irregularly russeted, frequently compressed. *Calyx* large, usually somewhat open. *Basin* moderately deep to deep, medium in width, very abrupt, furrowed and wrinkled.

Skin smooth, rather tough, clear bright yellow washed with red which sometimes deepens to a pinkish blush, striped and splashed with carmine and mottled over the base with dull scarf-skin. *Dots* scattering, small to large, pale gray, conspicuous. *Prevailing effect* greenish-yellow; not particularly attractive.

Calyx tube long, wide, funnel-shape. *Stamens* variable but usually median.

Core rather small, usually abaxile; cells sometimes unsymmetrical, wide open; core lines clasping. *Carpels* broadly ovate to elliptical, emarginate, sometimes tufted. *Seeds* rather dark brown, small to medium, rather numerous, very short, very plump, obtuse.

Flesh nearly white, firm, rather coarse, crisp, juicy, brisk subacid, fair to good.

Season October to December.

JULY.

REFERENCES. 1. *Rural N. Y.*, 1861 (cited by 15). 2. Hovey, *Mag. Hort.*, 29:112. 1863. 3. Hanford, *Horticulturist*, 19:273. 1864. fig. 4. Warder, 1867:719. 5. Downing, 1869:181. fig. 6. Thomas, 1875:499. 7. *Ill. Hort. Soc. Rpt.*, 1876:252. 8. *Ia. Hort. Soc. Rpt.*, 1879:471. 9. Hoskins, *Rural N. Y.*, 47:646. 1888. 10. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:292. 11. *Amer. Gard.*, 12:570. 1891. figs. 12. Bailey, *Am. Hort.*, 1892:239. 13. Burrill and McCluer, *Ill. Sta. Bul.*, 45:321. 1896. 14. Budd-Hansen, 1903:108. 15. Ragan, *U. S. B. P. I. Bul.*, 56:160. 1905.


SYNONYMS. *August* (4, 15, of Cassel, Germany, 3). *FOURTH OF JULY* (2, 4, 6, 8, 9, 10, 11, 12, 13). *Fourth of July* (15, ? 5). *JULY* (1, 3, 7, 15). *JULY, Fourth of* (14). *McAdow's June* (4, 15). *Siberian August* (4, 15, of Germany 2). *Stewart's Nonpareil* (15, ? 4). *TETOFSKI* (5). *Tetofsky*, erroneously (4, 15).

This fruit closely resembles Tetofsky and some have considered the two varieties identical (4) but they are quite distinct in tree. It is not recommended for planting in New York because it is not equal to standard varieties of its season.

Historical. Hovey states that "The Fourth of July apple, in Germany called the Siberian August apple, was sent from the Russian province Lief-land, in the year 1807, to the celebrated pomologist, Dr. Diel, and is celebrated, like all our summer apples which originated in Russia, for its great productiveness and hardiness" (2). It was introduced into Columbus, O., from Cassel, Germany (3). It has been disseminated to a considerable extent in various parts of this country and is still listed by a considerable number of nurserymen (12).

TREE.

Tree vigorous. *Form* upright, roundish and rather dense. *Twigs* short, straight, stout with large terminal buds; internodes medium. *Bark* dull



JUDSON

brown tinged with green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, small, oblong, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit below medium to above, pretty uniform in shape but not in size. *Form* usually roundish conical, irregularly ribbed; sides often unequal. *Stem* medium to long, moderately slender, often bracted. *Cavity* acuminate to acute, moderately deep, medium to narrow, slightly furrowed, thinly russeted. *Calyx* medium to large, usually closed; lobes medium in length, moderately narrow, acuminate. *Basin* rather shallow to medium in depth, narrow, somewhat abrupt, furrowed and wrinkled.

Skin thin, tough, smooth, rather glossy, pale yellow washed and mottled with red striped and splashed with carmine and overspread with whitish bloom. *Dots* small, numerous, submerged, inconspicuous, light, areolar.

Calyx tube variable in length, funnel-shape. *Stamens* median to marginal. *Core* medium or below, axile; cells closed; core lines slightly clasping or meeting. *Carpels* roundish ovate or elongated ovate. *Seeds* very dark dull brown, medium size, moderately wide, short, plump, obtuse.

Flesh yellowish, a little coarse, crisp, tender, moderately juicy, sprightly subacid, fair to good.

Season last of July to September.

KAIGHN.

REFERENCES. 1. Coxé, 1817:128. fig. 2. Downing, *Horticulturist*, 1:341. 1847. 3. Cole, 1849:125. 4. Thomas, 1849:173. 5. Emmons, *Nat. Hist. N. Y.*, 3:60. 1851. 6. Elliott, 1854:141. fig. 7. Hooper, 1857:49. 8. Downing, 1857:158. 9. Warder, 1867:681. fig. 10. Downing, 1876:55 app. 11. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 12. Bailey, *An. Hort.*, 1892:242.

SYNONYMS. KAIGHN (11). KAIGHN'S SPITZENBURG (1). KAIGHN'S SPITZENBURG (3, 6, 9). KAIGHN'S SPITZENBURGH (2). KAIGHN'S SPITZENBURG (7, 12). KAIGHN'S SPITZENBURGH (4, 5, 8). KAIGHN'S SPITZENBURG (10). *Lady Finger* (erroneously 6, 7). *Long John* (6, 7). *Long Pearmain* (6, 7). *Ohio Wine* (6). *Red Pearmain* (6, 7, 8). *Red Phoenix* (6). *Red Pippin* (6). *Red Spitzenberg* (6). *Red Spitzenberg* (8). *Red Winter Pearmain* (6). *Russam* (6). *Scarlet Pearmain*, erroneously (6). Downing states that Kaighn's Spitzenberg and Long Red Pearmain, for many years considered identical, are in reality distinct varieties in both tree and fruit. He gives a long list of synonyms for Long Red Pearmain including all of the synonyms cited above and adds, "The true Kaighn's Spitzenberg, so far as I know, has no synonyms" (10).

This is an old variety of New Jersey origin which has been disseminated through various parts of the West even to the Pacific Coast. The fruit is showy and the tree productive. The tree makes a spreading, straggling growth (1, 11). According to Coxé (1) the fruit bears "a faint resemblance to the Esopus Spitzenberg but is more pointed toward the crown; the color is a lively but pale red, faintly streaked and full of white spots; the

skin is smooth, the stem long and deeply planted, the crown very hollow—the flesh finely flavored, yellow, juicy and tender." It is now practically obsolete in New York.

KALKIDON.

REFERENCES. 1. Budd, *Ia. Agr. Coll. Bul.*, 1885:25. 2. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-7:76. 3. Gibb, *Am. Pom. Soc. Rpt.*, 1887:56. 4. Lyon, *U. S. Pom. Bul.*, 2:41. 1888. 5. Beach, *N. Y. Sta. An. Rpt.*, 11:593. 1892. 6 *Ib.*, 13:589. 1894. 7. *Ib.*, 14:261. 1895. 8. Beach, Paddock and Close, *Ib.*, 15:272, 281. 1896. 9. Beach and Clark, *N. Y. Sta. Bul.*, 248:127. 1904. 10. Ragan, *U. S. B. P. I. Bul.*, 56:161. 1905.

SYNONYMS. KALKIDON (3, 4, 9, incorrectly 10). KALKIDONSKOE (1). KALKIDOUSKOE (5, 6, 7, 8). KALKIDOVSKOE (2). *Khalkidonskoe* (4, 9). *Khalkidouskoe* (3, 10). No. 510 (10). No. 94 (1, 2, 3, 10).

A Russian variety which was received from Ellwanger and Barry, Rochester, N. Y., in 1884 for testing at this Station. It is an apple of moderately attractive appearance and fair to good quality, in season in September and October. The tree comes into bearing moderately young and is a reliable biennial cropper. It is not recommended for planting in New York because it is inferior to standard varieties of its season.

FRUIT.

Fruit large to medium, fairly uniform in size and shape. *Form* oblate conical to ovate, regular or faintly ribbed. *Stem* medium to very short, moderately thick. *Cavity* acute to almost acuminate, moderately shallow to deep, medium in width, furrowed and compressed, russeted and with outspreading greenish-russet rays. *Calyx* medium to large, closed or partly open; lobes medium in length, broad, acute. *Basin* shallow to medium in depth, narrow to medium in width, rather abrupt, slightly furrowed.

Skin thick, rather tough, smooth, greenish or pale yellow, largely washed and mottled with dull red, splashed and striped with carmine. *Dots* variable in size, numerous, inconspicuous, submerged.

Calyx tube rather long, moderately wide, conical or funnel-shape. *Stamens* median to marginal.

Core small, axile or sometimes abaxile; cells symmetrical, closed or open; core lines clasping. *Carpels* ovate to elliptical, slightly emarginate, slightly tufted. *Seeds* medium to large, wide, plump, acute to obtuse, dark brown.

Flesh tinged with yellowish-green, moderately fine-grained, tender, rather juicy, mild subacid, fair to good.

Season September to midwinter (9).

KARABOVKA.

REFERENCES. 1. Gibb, *Montreal Hort. Soc. Rpt.*, 1883:74. 2. Budd, *Ia. Agr. Coll. Bul.*, 1885:7. 3. Gibb, *Ia. Hort. Soc. Rpt.*, 1885:274. 4. *N. Y. Sta. An. Rpt.*, 13:582. 1894. 5. Ragan, *U. S. B. P. I. Bul.*, 56:161. 1905.

SYNONYMS. *Kajabowka* (5). KARABOFF (2, 3, 4). *Karaboff* (1, 5).

KARABOVKA (5). *Karabovka* (3, 4). KARABOWKA (1). *Karabowka* (2, 4). *No. 21 M* (3, 4). *No. 205* (2, 3, 4, 5). *Shro. to Ia. No. 21* (5).

A Russian variety received from Ellwanger and Barry, Rochester, N. Y., in 1884 for testing at this Station. As grown here the tree does not come into bearing very early but when mature yields full crops biennially. The fruit is medium to rather small, not specially attractive in appearance, fair to possibly good in quality being inferior to standard sorts of its season. Season late August and September. It does not agree with the description of *Karabovka* given by Budd (2).

TREE.

Tree moderately vigorous with moderately long, stout, curved branches. *Form* rather open, flat, spreading. *Twigs* long, curved, stout, with large terminal buds; internodes medium. *Bark* dark brown, heavily streaked with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous, small to medium, round, slightly raised. *Buds* large, prominent, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit small to medium, uniform. *Form* oblate, regular, symmetrical. *Stem* medium in length, moderately thick to rather slender. *Cavity* acute to nearly acuminate, moderately deep to deep, narrow to medium in width, occasionally lipped, usually russeted. *Calyx* large, open or nearly closed; lobes short, rather broad, acute. *Basin* medium in depth, wide, furrowed or wrinkled and with mammiform protuberances.

Skin thin, tender, smooth, rather dull pale greenish-yellow, with scattering narrow stripes of dull dark red, or when highly colored it is largely striped, splashed and shaded with red, and overspread with pinkish bloom. *Dots* rather numerous, small, light, obscure, submerged.

Calyx tube large, medium in length, rather wide, broadly conical to funnel-shape. *Stamens* median to marginal.

Core medium size, axile; cells closed; core lines clasping. *Carpels* elliptical, emarginate. *Seeds* very dark dull brown, medium size, wide, rather short, flat, plump, obtuse.

Flesh nearly white, moderately fine, tender, moderately juicy, mild subacid with peculiar flavor, fair to good in quality.

Season late August and September.

KENTISH FILLBASKET.

REFERENCES. 1. Kenrick, 1832:95. 2. Downing, 1845:114. 3. Thomas, 1849:168. 4. Emmons, *Nat. Hist. N. Y.*, 3:62. 1851. 5. Elliott, 1854:172. 6. Hooper, 1857:49. 7. Warder, 1867:723. 8. Hogg, 1884:120. 9. Green, *Country Gentleman*, 1885:840. 10 *Ib.*, *Can. Hort.*, 11:8. 1888. 11. Bailey, *Am. Hort.*, 1892:242. 12. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 4:3. 1897. figs. 13. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354.

SYNONYMS. KENT FILLBASKET (13). KENTISH FILLBASKET (3). KENTISH FILLBASKET (1. 2. 4. 5. 6. 7. 8. 9. 10. 11. 12). *Lady de Grey's* (2, 5, 8).

Potter's Large (8). *Potter's Large Grey Seedling* (5). *Potter's Large Seedling* (2).

A very large, handsome, late autumn apple, desirable for cooking but not for dessert use. Tree a strong grower and a fair cropper. Not recommended for planting in New York.

According to Hogg (8) the Kentish Fillbasket of Miller, Forsyth and Rogers is a different variety being evidently the Kentish Codlin or common old English Codlin, a lemon-yellow apple which is in season from August to October.

The Kentish Fillbasket of Buel¹ appears to be the same as that of Forsyth.²

Historical. This is an old English variety. It has been sparingly cultivated in portions of New York state for many years and has been grown to some extent also in Ontario (12).

FRUIT.

Fruit very large. *Form* oblate or roundish, ribbed broadly and obscurely if at all, irregular, fairly uniform. *Stem* not exerted, short, medium in thickness. *Cavity* acute to somewhat acuminate, deep, broad, symmetrical or somewhat furrowed, green or more often with outspreading russet. *Calyx* small to rather large, closed or partly open; lobes broad, obtuse to acute. *Basin* pretty abrupt, moderately deep to deep, medium in width to wide, sometimes a little furrowed or wrinkled.

Skin thin, tough, smooth, somewhat waxy, pale yellow with thin brownish blush often deepening to red, somewhat mottled and splashed with carmine. *Dots* small usually not conspicuous, dark brown or grayish or submerged and whitish. *Prevailing effect* yellow somewhat striped with red.

Calyx tube wide, conical. *Stamens* basal to nearly median.

Core abaxile, medium to large; cells often unsymmetrical and open, sometimes closed; core lines nearly meeting. *Carpels* broadly ovate, mucronate, not emarginate, somewhat tufted. *Seeds* medium or below, plump, acute.

Flesh whitish, firm, moderately coarse, crisp, rather tender, juicy, brisk subacid, good.


Season October to December.

KESWICK.

REFERENCES. 1. Forsyth, 1824:132. 2. *London Hort. Soc. Cat.*, 1831:No. 225. 3. Kenrick, 1832:89. 4. Floy-Lindley, 1833:23. 5. Downing, 1845:87. 6. Thomas, 1849:156. 7. Emmons, *Nat. Hist. N. Y.* 3:37. 1851. 8. Barry, 1851:280. 9. Elliott, 1854:141. 10. Barry, *Horticulturist*, 10:87. 1855. 11. Gregg, 1857:37. 12. Hooper, 1857:25, 49, 107, 111. 13. *Am. Pom. Soc. Cat.*, 1860. 14. Mead, *Horticulturist*, 17:150. 1862. 15. Warder, 1867:288. fig. 16. Fitz, 1872:160. 17. Hogg, 1884:122. 18. Wickson, 1889:243. 19. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 20. Bailey, *Am. Hort.*, 1892:242. 21. Taft, *Mich. Sta. Bul.*, 105:108. 1894. 22. Lyon, *ib.*, 118:60. 1895. 23 *ib.*, 143:200. 1897. 24. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 25. Dickens and

¹ Buel, *N. Y. Bd. Agr. Mem.*, 1826:477.

² Forsyth, 1803:50.



KESWICK

Greene, *Kan. Sta. Bul.*, 106:53. 1902. 26. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 27. Budd-Hansen, 1903:110. 28. Beach and Clark, *N. Y. Sta. Bul.*, 248:128. 1904.

SYNONYMS. CODLIN, KESWICK (1, 2, 24). KESWICK (19, 21, 22, 23, 26, 27, 28). KESWICK CODLIN (3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 24, 25). *Keswick Codlin* (27, 28). KESWICK CODLING (15). *No. 225* (2).

This variety is particularly esteemed on account of its excellence for culinary use. It is not adapted for storage, its commercial limit in ordinary storage being September and early October. It comes into season late in August or early in September and ripens continuously during a period of several weeks. It does not stand heat well before going into storage and goes down quickly (28). The fruit is of good medium size to rather large, greenish-yellow, sometimes with faint blush; flesh brisk subacid. It is grown chiefly for home use and only to a very limited extent for local market. The tree is a good grower, hardy, healthy, long-lived, comes into bearing quite young and yields good to very good crops almost annually.

This is distinct from the Codling or English Codling described by Coxe, Thacher, Forsyth and others.

Historical. Hogg (17) states that: "This excellent apple was first discovered growing among a quantity of rubbish behind a wall at Gleaston Castle, near Ulverstone, and was first brought into notice by one John Sander, a nurseryman at Keswick, who, having propagated it, sent it out under the name of Keswick Codlin.

"In the Memoirs of the Caledonian Horticultural Society, 1813, Sir John Sinclair says: 'The Keswick Codlin tree has never failed to bear a crop since it was planted in the episcopal garden at Rose Castle, Carlisle, twenty years ago.'"

It has long been known in this country and very old trees of it are found in some orchards but it is nowhere extensively cultivated being grown chiefly for home use. It is quite commonly listed by nurserymen (20).

TREE.

Tree medium to large, moderately vigorous. *Form* upright spreading to roundish. *Twigs* moderately long, curved, stout; internodes short. *Bark* dull brown, heavily coated with gray scarf-skin; pubescent. *Lenticels* numerous, rather conspicuous, medium to small, round, not raised. *Buds* medium to large, prominent, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit above medium to nearly large, not very uniform. *Form* roundish conic or inclined to oblong conic, rather broad at the base, distinctly ribbed;

sides frequently a little unequal. *Stem* medium to short, slender to rather thick. *Cavity* variably acute, medium in depth to shallow, rather narrow to moderately broad, more or less russeted. *Calyx* medium size, closed; lobes long, medium in width, nearly acuminate. *Basin* shallow, moderately narrow, furrowed or angular, often with fleshy protuberances alternating with the calyx lobes.

Skin thin, tough, smooth, waxy, pale greenish or yellow, sometimes with a faint blush and often with a suture line extending out from the cavity. *Dots* submerged, inconspicuous or russet.

Calyx tube medium in length, rather wide, bluntly cone-shape. *Stamens* median.

Core variable, large, abaxile; cells wide open; core lines meeting. *Carpels* variable, roundish ovate. *Seeds* very light brown, very small, medium in width, short, very plump, acute.

Flesh nearly white, fine, tender, very juicy, brisk subacid, good for culinary use, too acid for dessert unless very ripe.

Season August and September.

KIRKBRIDGE.

REFERENCES. 1. Downing, 1857:160. 2. *Mag. Hort.*, 24:108. 1858. 3. Warder, 1867:671. 4. *Am. Pom. Soc. Cat.*, 1871:8. 5. Downing, 1872:10 index, app. 6. Thomas, 1875:195. 7. Downing, 1881:11 index, app. 8. *Ib.*, 1881:12 index, app. 9. Budd-Hansen, 1903:111.

SYNONYMS. *Bohannon* (2). *Conic June* (7). KIRKBRIDGE (2, 3, 7, 8, 9). KIRKBRIDGE WHITE (1, 4, 5, 6). *White June* (5). *Yellow Flat* (8). *Yellow June* (6, erroneously 3, of some 5).

Fruit small to medium, oblong conic, pale yellow or whitish; flesh white, fine, tender, juicy, pleasant subacid, good to very good. The tree is a moderate grower, comes into bearing young and is productive. Season August and September. Not recommended for planting in New York.

Historical. In 1867 Warder (3) wrote concerning Kirkbridge White: "This fruit has been pretty extensively cultivated in some parts of the Western states and sometimes it is mistaken for the Yellow June." It is said to be of American origin. It is but little known in New York.

LADY FINGER.

REFERENCE. 1. Downing, 1869:245.

SYNONYMS. None.

Under this name Downing (1) describes an apple which he received from Maryland, the fruit of which is of medium size, yellowish, nearly overspread with deep crimson; flesh white, tender, pleasant subacid, good to very good; season August.

Other varieties have been known under the name Lady Finger several of which have already been mentioned in the discussion of the winter apples. See Vol. I, page 183.



LANDSBERG

LANDSBERG.

REFERENCES. 1. Berghuis, 1868:No. 8. *col. pl.* 2. Downing, 1872:20 app. 3. Leroy, 1873:701. *fig.* 4. Lauche, 1:No. 38. 1882. *col. pl.* 5. Hogg, 1884:128. 6. Budd, *Ia. Agr. Coll. Bul.*, 1885:34, 41. 7. Lyon, *U. S. Pom. Bul.*, 2:42. 1888. 8. Gaucher, 1894:No. 17. *col. pl.* 9. Beach and Paddock, *N. Y. Sta. An. Rpt.*, 14:253, 261. 1895. 10. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:47. 1903. 11. Beach and Clark, *N. Y. Sta. Bul.*, 248:129. 1904.

SYNONYMS. LANDSBERG (7, 11). LANDSBERGER REINETTE (2, 4, 5, 6, 8, 9). *Landsberger Reinette* (11). LANDSBURG (6). *Landsburger Reinette* (6, 7). LANSBERGER REINETTE (10). REINETTE DE LANDSBERG (3). *Reinette de Landsberg* (2, 8). REINETTE LANDSBERGER (1).

An attractive late autumn and early winter apple of good size and pleasant subacid flavor. The tree is thrifty, comes into bearing rather early and is reliably productive yielding good crops biennially. The fruit has a clear, pale waxen yellow or greenish skin which readily shows bruises. It is a good dessert apple but less desirable for culinary uses because when it is cooked it lacks character in texture, color and flavor. It is easily excelled for any purpose by standard sorts of its season and is not recommended for planting in New York.

Historical. Raised from seed about 1840 by Mr. Burkhardt, justice of the peace in Landsberg, Germany (2, 4). Imported from Silesia, Germany, in 1883 by Professor Budd for the Iowa State College (6).

TREE.

Tree vigorous. *Form* roundish or spreading, rather open. *Twigs* short to medium, nearly straight, rather stout; internodes medium. *Bark* olive-green tinged with brownish-red, mottled with scarf-skin; pubescent. *Lenticels* quite numerous, small, round, not raised. *Buds* large to below medium, broad, plump, generally obtuse, free, quite pubescent.

FRUIT.

Fruit medium to large, pretty uniform in shape and size. *Form* roundish conic to roundish oblate, obscurely angular, pretty regular; sides sometimes unequal. *Stem* short to medium. *Cavity* acute to acuminate, deep, rather wide, sometimes obscurely furrowed, usually smooth but sometimes russeted. *Calyx* segments long, acute, reflexed, sometimes closed. *Basin* medium in width and depth, often somewhat furrowed, wrinkled.

Skin thin, tough, smooth or slightly roughened by russet dots, waxen yellow or pale green, sometimes with attractive crimson blush. *Dots* numerous, submerged and whitish, sometimes russet. *Prevailing color* pale yellow, rather attractive.

Calyx tube large, wide, cone-shape. *Stamens* median.

Core abaxile, medium; cells usually symmetrical and wide open; core lines slightly clasping. *Carpels* broadly ovate, much concave, narrow toward the apex, nearly smooth. *Seeds* numerous, medium in size, broad, obtuse, medium brown, smooth or nearly so.

Flesh nearly white, very tender, crisp, rather fine-grained, mild subacid, agreeable in flavor, good to very good for dessert.

Season mid-October to January (8, 9) Some of the fruit keeps apparently sound till March or later but after January it loses in quality.

LATE STRAWBERRY.

REFERENCES. 1. Thomas, *Cultivator*, 5:246. 1848. 2. Thomas, 1849:150. *fig.* 3. Cole, 1849:111. 4. Waring, 1851:21. 5. Emmons, *Nat. Hist. N. Y.*, 3:27. 1851. *fig.* 6. Barry, 1851:282. 7. Elliott, 1854:65. 8. Downing, 1857:163. 9. Hooper, 1857:54. 10. Gregg, 1857:41. 11. *Am. Pom. Soc. Cat.*, 1862. 12. Warder, 1867:540. 13. Downing, 1869:250. *fig.* 14. Todd, 1871:154. *fig.* 15. *Ja. Hort. Soc. Rpt.*, 1879:538. 16. Roach, *Montreal Hort. Soc. Rpt.*, 1886-7:27. 17. Wickson, 1889:245. 18. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 19. Bailey, *Am. Hort.*, 1892:243. 20. Burrill and McCluer, *Ill. Sta. Bul.*, 45:329. 1896. 21. Badd-Hansen, 1903:113. *fig.*


SYNONYMS. AUTUMN STRAWBERRY (1, 4, 6, 7, 10, 14, 15). *Autumn Strawberry* (2, 5, 8, 12, 13, 17, 21). FALL STRAWBERRY (3). LATE STRAWBERRY (2, 5, 8, 9, 11, 12, 13, 16, 17, 18, 19, 20, 21). *Late Strawberry* (1, 3, 6, 7). *Strawberry* (2, 7).

Different varieties have been called Late Strawberry but the name is now almost exclusively applied to the variety described below. This is an attractive apple, pale yellow overspread or striped and splashed with light and dark red, very good in quality especially for dessert use. Many esteem it one of the best dessert apples of its season. It comes into use in September and ripens in succession during a period of several weeks continuing in season till December. While this habit of successive ripening makes the variety more desirable for the home orchard it renders it less valuable for commercial purposes, since several pickings are required to secure the crop in prime condition. The fruit is hardly as large as is desirable for a good market variety but its attractive appearance and excellent quality render it suitable for local and fancy trade. The tree is medium to rather large, vigorous; form upright spreading to roundish. It is hardy, healthy, long-lived and a regular cropper-yielding moderate to heavy crops biennially or nearly annually.

Historical. Late Strawberry originated at Aurora, Cayuga county, N. Y. (13). In 1848 Thomas described it as a new and newly introduced apple (1).

FRUIT.

Fruit below medium to above, uniform in size and fairly uniform in shape. *Form* roundish to slightly oblong conic, sometimes quite strongly ribbed,



LATE STRAWBERRY

rather irregular. *Stem* long, rather slender, often curved. *Cavity* acuminate, deep, usually broad, furrowed, sometimes with thin radiating streaks of light russet mingled with carmine. *Calyx* large, open or partly open; lobes often separated at base, rather short, acute, erect or reflexed. *Basin* deep, moderately narrow to rather wide, abrupt, furrowed and wrinkled.

Skin attractive pale yellow often almost entirely overspread with bright pinkish-red, dotted and streaked with purplish-carmine. *Dots* small, not very numerous, inconspicuous, light colored. *Prevailing effect* bright striped red.

Calyx tube rather wide, conical to slightly funnel-shape. *Stamens* basal.

Core rather small, nearly axile to somewhat abaxile; cells closed or somewhat open; core lines meeting or slightly clasping. *Carpels* obovate. *Seeds* rather large, flat, obtuse.

Flesh yellowish-white, fine, crisp, tender, juicy, somewhat sprightly aromatic, subacid, very good.

Season September to December.

LATHAM.

REFERENCE. 1. Downing, 1869:251.

SYNONYMS. None.

This is a variety which we have not seen; so far as we can learn it is no longer cultivated. According to Downing it originated in Sag Harbor, Suffolk county, N. Y.; tree very productive; fruit medium size, yellow mostly covered with light and dark red; flesh white, juicy, mild subacid, good; season November and December (1).

LEAD.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 8:40. 1881-82. 2. *Ia. Hort. Soc. Rpt.*, 1882:78. 3. *Montreal Hort. Soc. Rpt.*, 1883:102. 4. *Ia. Agr. Coll. Bul.*, 1883:28. 5. *Ib.*, 1885:11. 6. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-7:71. 7. *Ia. Agr. Coll. Bul.*, 1890:23. 8. Budd, *Ia. Sta. Bul.*, 19:540. 1892. 9. Bailey, *Am. Hort.*, 1892:243. 10. Munson, *Me. Sta. Rpt.*, 1896:74, 79. 11. Stinson, *Ark. Sta. Bul.*, 43:105. 1896. 12. Thomas, 1897:291. fig. 13. Troop, *Ind. Sta. Rpt.*, 1899:80. 14. Hansen, *S. D. Sta. Bul.*, 76:64. 1902. 15. Munson, *Me. Sta. An. Rpt.*, 18:84. 1902.

SYNONYMS. LEAD (1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15). *Lead Apple* (3). No. 3 M (2, 4, 7, 8, 10, 14, 15). No. 277 (2, 4, 5, 10). *Strinsovka* (1). SWINEZ (3). *Swinsowska* (5).

As noted below two distinct Russian varieties have been disseminated under the name Lead. Some of the references cited above refer to one of these varieties and some to the other.

A Russian variety was received from the Iowa Agricultural College in 1890 for testing at this Station, the fruit of which is pale greenish-yellow with a blushed cheek and carmine splashes, medium size or above; flesh tender, rather juicy, subacid, fair quality; season late August and September. The tree does not come into bearing very young. It is an annual cropper but only moderately productive. It is not worthy of further testing for this region.

This appears to be the same variety as that described by Hansen under the name Lead with the synonym No. 362 (14). This he says is not the true Lead. He describes the true Lead with synonym No. 3 M as a Russian variety, large, heavy, roundish, greenish-yellow with dull blush; flesh greenish-white, sharp subacid, good in quality; season early winter.

LINCOLN PIPPIN.

REFERENCE. 1. Downing, 1881:93 app. fig.

SYNONYMS. None.

Fruit medium to large, yellow with no blush, subacid, excellent for either dessert or culinary uses; season November and December. Under favorable conditions some portion of the fruit may be kept through the winter. The tree is large, spreading, somewhat open, moderately vigorous with rather short, stout twigs, hardy, long-lived. It does not come into bearing very young but when mature is a pretty reliable annual bearer, ripening the crop evenly. The fruit is fair, averages pretty uniform in size and shape and is reliable and satisfactory in color and quality. Downing describes it as "medium to large, roundish oblate, slightly conical, slightly angular, sides sometimes a little unequal; skin pale greenish-yellow, moderately sprinkled with grayish dots; stalk short to long, slender; cavity large, deep, calyx small, closed; basin small or medium, slightly corrugated; flesh half fine, pale whitish-yellow, tender, juicy, subacid, slightly aromatic; very good; core rather large. October, December" (1).

Historical. So far as we can learn this variety is cultivated only in the vicinity of Syracuse. Downing states that it is an old variety, said to have been brought to Syracuse from Connecticut; the original name having been lost it was named Lincoln after Reuben Lincoln who brought it into notice (1).

LINDENWALD.

REFERENCES. 1. Downing, 1869:254. 2. Burrill and McCluer, *Ill. Sta. Bul.*, 45:330. 1896.


SYNONYMS. None.

A variety which originated with J. G. Sickles, Stuyvesant, Columbia county, N. Y. Downing describes the fruit as of medium size, yellow with light shades of red; flesh crisp, juicy, pleasant subacid, good to very good; season September (1).

We have received no report of this being grown outside of the locality of its origin.

LONGFIELD.

REFERENCES. 1. Webster, *Montreal Hort. Soc. Rpt.*, 7:52. 1881. 2. *Ib.*, 8:71. 1881-82. 3. Budd, *Id. Hort. Soc. Rpt.*, 1882:77. 4. Gibb, *Ib.*, 1883:425. 5. Webster, *Am. Pom. Soc. Rpt.*, 1883:113. 6. Budd, *Id. Agr. Coll. Bul.*, 1883:28. 7. Gibb, *Montreal Hort. Soc. Rpt.*, 1883:66. fig. 8. Budd, *Id. Agr. Coll. Bul.*, 1885:5. 9. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:15. 10. Schroeder, *Ib.*, 1886-87:74. 11. Craig, *Ib.*, 1886-87:103. 12. Hoskins, *Rural*



LONGFIELD

N. Y. 47:558. 1888. 13. *Am. Pom. Soc. Cat.*, 1889:8. 14. Gibb, *Can. Hort.*, 12:27. 1889. 15. *Ib.*, 13:207, 216, 301. 1890. 16. Budd, *Ia. Agr. Coll. Bul.*, 1890:19. 17. *Ia. Sta. Bul.*, 19:536. 1892. 18. *Can. Hort.*, 15:29. 1892. 19. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57, 58. 20. Bailey, *An. Hort.*, 1892:243. 21. *Amer. Gard.*, 14:50, 177, 305. 1893. 22. *Can. Hort.*, 16:204, 360. 1893. 23. Craig, *Ont. Fr. Gr. Assn. An. Rpt.*, 26:16. 1894. 24. Beach, *N. Y. Sta. An. Rpt.*, 13:582. 1894. 25. Troop, *Ind. Sta. Bul.*, 53:123. 1894. 26. Craig *Can. Dept. Agr. Rpt.*, 1894:126. figs. 27. Freeborn, *Nat. Nurseryman*, 2:133. 1894. 28. Van Deman, *Rural N. Y.*, 55:848. 1896. 29. Munson, *Me. Sta. Rpt.*, 1896:75. 30. Stinson, *Ark. Sta. Bul.*, 43:105. 1896. 31. Buckman, *Rural N. Y.*, 56:39. 1897. 32. Waugh, *Pt. Sta. Bul.*, 61:31. 1897. 33. Thomas, 1897:291. fig. 34. *Am. Gard.*, 19:650, 682. 1898. 35. *Rural N. Y.*, 57:736, 819. 1898. 36. Troop, *Ind. Sta. Rpt.*, 1899:80. 37. Hansen, *S. D. Sta. Bul.*, 76:67. 1902. fig. 38. Munson, *Me. Sta. An. Rpt.*, 18:84, 85, 87, 95. 1902. 39. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:47. 1903. 40. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 41. Budd-Hansen, 1903:116. 42. Beach and Clark, *N. Y. Sta. Bul.*, 248:129. 1904.

SYNONYMS. *English Pippin* (26). 587 (26). 57 M (16). 56 M (2, 3, 6, 10). *Good Peasant* (9). *LANGERFELDSKOE* (1, 4, 5, 7, 13). *Langerfeldskoe* (8). *LONGFIELD* (2, 3, 6, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28. 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42). *Longfield* (4, 7). *Longfield's Apple* (1, 5). *No. 161* (2, 3, 6, 8, 16, 24, 29, 37).

The fruit of Longfield is usually below medium size but decidedly attractive in appearance for a yellow apple, being clear waxen yellow, lightly blushed with bright red. Its flesh is white, crisp, fine, very tender and of pleasant quality. It may well be classed among the fancy dessert apples; it is good also for culinary uses. In marketing this fruit it is necessary to handle it with great care because ordinarily its texture is so very tender and its color so delicate that it shows bruises very readily. It is not well adapted for holding outside of cold storage. In ordinary storage its commercial limit at Geneva is late September or early October (42) and in cold storage it may be kept till December (42); but as grown further north it may be kept through the winter (26). The tree is a moderate grower, very hardy and very productive; in fact it bears such heavy crops that the fruit is liable to be deficient in size. On account of the hardiness and productiveness of the tree and the beauty and good quality of the fruit Longfield is recommended for planting for home use and for local and special markets.

Historical. Longfield was first imported from Russia by the United States Department of Agriculture in 1870; later it was imported from various European sources for the Iowa Agricultural College by Professor Budd.

It is now frequently listed by nurserymen (20) and is being planted to a limited extent in various parts of the state, but it has not yet come to be commonly known among New York fruit growers.

TREE.

Tree medium in size with short, moderately stout, crooked branches and drooping laterals filled with small spurs. *Form* roundish or spreading, dense, rather low. *Twigs* medium in length, curved, moderately stout; terminal buds large; internodes short. *Bark* dark brown, lightly streaked with scarf-skin; pubescent. *Lenticels* scattering, medium size, oval, not raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium to small, usually below medium; uniform in size and shape. *Form* roundish conic, slightly ribbed. *Stem* medium to short, rather slender. *Cavity* acuminate to acute, medium to rather deep, narrow, quite symmetrical, usually slightly russeted. *Calyx* small, leafy, closed or partly open; lobes long, rather narrow, acute to acuminate. *Basin* small, shallow to medium in depth, narrow, somewhat abrupt, slightly furrowed and wrinkled.

Skin thin, tender, smooth, glossy, pale waxen yellow or whitish usually with a lively pinkish blush but not striped. *Dots* few, small, inconspicuous, whitish, usually submerged. *Prevailing effect* attractive bright pale yellow partly blushed with lively red.

Calyx tube narrow and elongated, often extending to the core. *Stamens* basal to median.

Core medium to above, axile or nearly so; cells symmetrical, not uniformly developed, closed or partly open; core lines clasping the funnel cylinder. *Carpels* roundish, slightly emarginate. *Seeds* rather large, moderately dark brown, long, acute approaching acuminate.

Flesh nearly white, fine, crisp, very tender, juicy, subacid, sprightly, aromatic, good to very good.

Season September to October or later.

LONG ISLAND PEARMAIN.

REFERENCES. 1. COXE, 1817:144. fig. 2. WARDER, 1867:682. 3. DOWNING, 1869:255.

SYNONYMS. *Autumn Pearmain* (2). *Hollow Crown Pearmain* (3). *LONG ISLAND PEARMAN* (1, 2, 3). *Winter Pearmain* (2).

This is an old variety which COXE (1) thus describes: "A handsome large apple, of an oblong form, about the size of a Priestly—the stem is short, not deeply planted; the crown large and hollow; the skin streaked with large blotches of red on a rich yellow ground, with faint russet spots—the flesh is tender, coarse and pleasant, partaking of that dryness characteristic of all the varieties of the pearmain—it ripens in October and keeps till March." DOWNING (3) states that it has sometimes been confused with *Winter Pearmain* which is an entirely different fruit. So far as we can discover, *Long Island Pearmain* is now obsolete in New York.

LONG RED PEARMAIN.

REFERENCE. 1. Downing, 1876:55 app.

SYNONYMS. *English Pearmain* incorrectly (1). *Hudson Red Streak* (1). *Kentucky Bellflower* (1). *Kentucky Gillflower* (1). *Kaighn's Spitzenburg* incorrectly (1). *Lady Finger* incorrectly (1). *Long John* (1). *Long Pearmain* (1). LONG RED PEARMAIN (1). *Mudhole* (1). *Park* (not of Kansas) (1). *Pearmain* (1). *Pound Royal* incorrectly (1). *Red Bellflower* incorrectly (1). *Red Pearmain* (1). *Red Phoenix* (1). *Red Pippin* (1). *Red Spitzenburg* (1). *Red Winter Pearmain* incorrectly (1). *Russam* (1). *Scarlet Pearmain* (1). *Sheepnose* of some (1). *Striped Pearmain* (1). *Wabash Bellflower* (1). *Winter Pearmain* (1).

This variety was for many years considered identical with Kaighn. See page 113. In 1876 Downing (1) stated that it was distinct from Kaighn in tree and fruit and gave the list of synonyms for it which is cited above.

The fruit is medium to large, oblong approaching conic, yellowish, shaded and striped with red; flesh coarse, pleasant subacid, good; season late fall and early winter.

LONG STEM.

REFERENCES. 1. Cole, 1849:106. 2. Hooper, 1857:55. 3. Warder, 1867:725. 4. Downing, 1869:256. 5. Ragan, *U. S. B. P. I. Bul.*, 56:183. 1905.

SYNONYMS. None.

Several varieties of the apple have been known in cultivation under the name Long Stem. One of these has already been described (Volume I, page 196) as Long Stem of Pennsylvania; others are noticed below. The references cited above do not all refer to the same variety.

LONG STEM OF MASSACHUSETTS. The following is Cole's description of this variety (1): "Large medial; flattish-round; pale yellow, brown full in the sun; dark specks and patches; stem extremely long, slender, in a broad, deep cavity; calyx large, rather open, in a broad, shallow basin; flesh white, rather tender, juicy, of a rich, mild, delicious, sprightly, aromatic flavor. First-rate for the dessert or cooking. Sept. 1st, to the last of Oct. Good and constant bearer. Origin, East Bridgewater, Ms."

LONG STEM OF CONNECTICUT. Downing (4) describes this variety as having young shoots of dull reddish-brown, somewhat downy, with prominent, flattened buds. Fruit medium, roundish oblate, yellow; flesh fine-grained, sweet, rich, good. Season September to January.

LONGWORTH.

REFERENCES. 1. Downing, 1881:94 app. 2. Beach and Clark, *N. Y. Sta. Bul.*, 248:129. 1904.

SYNONYMS. LONGWORTH (2). LONGWORTH RED WINTER (1).

Fruit of fairly good color but not very large; inferior in quality to standard varieties of its season. It is variable in season (2); some years it keeps fairly well till midwinter, but generally speaking, as grown at this Station,

its commercial limit in ordinary storage is November. The tree comes into bearing rather young and is a reliable cropper, yielding moderate to rather heavy crops biennially or sometimes annually. Not recommended for planting in New York.

Historical. Originated at Dubuque, Ia. (1). It was received for testing here in 1889 from Benjamin Buckman, Farmingdale, Ill.

LORD SUFFIELD.

REFERENCES. 1. Downing, 1869:257. 2. *Ib.*, *Tilt. Jour. Hort.*, 7:303. 1870. 3. Hogg, 1884:136. 4. Goff, *Rural N. Y.*, 46:685. 1887. figs. 5. *Can. Hort.*, 15:347. 1892. 6. Bailey, *Am. Hort.*, 1892:243. 7. Gaucher, 1894:No. 18. col. pl. 8. Dempsey, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. 9. *Jour. Roy. Hort. Soc.*, 1898:356.

SYNONYMS. *Lady Suffield* (7). *Lady Sutherland* (7). *Livesley's Imperial* (7). LORD SUFFIELD (1, 2, 3, 4, 5, 6, 7, 8, 9).

This variety of the Keswick Codlin group is considered by some one of the best apples of the group. The fruit is suitable for cooking as early as the middle of July and remains in use till September. It is large, roundish, varying from oblate to somewhat oblong; green, marked with clear light russet flecks and dots; calyx closed; basin ridged, shallow; stem short; cavity moderately wide, shallow; core large, abaxile; cells open; flesh white, rather fine, moderately juicy, subacid, good for culinary use. The tree is very productive, which perhaps accounts in part for its being short-lived. As grown at this Station it has blighted so badly as to make it an undesirable variety for the orchard.

LOU.

REFERENCES. 1. *Am. Pom. Soc. Rpt.*, 1885:28. 2. *Mo. Hort. Soc. Rpt.*, 1886:233. 3. *Mich. Sta. Bul.*, 118:60. 1895. 4. Beach, Paddock and Close, *N. Y. Sta. An. Rpt.*, 15:272. 1896. 5. *Mich. Sta. Bul.*, 143:200. 1897.

SYNONYMS. None.

A striped red apple of fairly good quality in season in early August. The tree is a good grower, comes into bearing rather young and is a reliable biennial cropper. It was originated from seed of Oldenburg by Peter M. Gideon, Excelsior, Minn., from whom it was received in 1888 for testing at this Station. It is not valuable for planting in New York.

LOUISE.

REFERENCES. 1. Woolverton, *Am. Pom. Soc. Rpt.*, 1889:155. 2. *Can. Hort.*, 15:19. 28. 1892. 3. Bailey, *Am. Hort.*, 1892:247. 4. *Can. Hort.*, 16:401. 1893. 5. Beach, *N. Y. Sta. An. Rpt.*, 12:601. 1893. 6. Hoskins, *Rural N. Y.*, 53:278. 1894. 7. Smith, *Can. Hort.*, 18:349. 1895. 8. Lyon, *Mich. Sta. Bul.*, 143:200. 1897. 9. *Am. Pom. Soc. Cat.*, 1899:18. 10. Waugh, *Vt. Sta. Bul.*, 83:91. 1900. 11. Woolverton, *Can. Hort.*, 23:46. 1900. fig. 12. *Ib.*, *Ont. Fr. Stas. An. Rpt.*, 7:8. 1900. figs. 13. Hansen, *S. D. Sta. Bul.*, 76:69. 1902. 14. Budd-Hansen, 1903:117. fig. 15. Farrand, *Mich. Sta. Bul.*, 205:45. 1903.

SYNONYMS. LOUISE (8, 10, 13, 15). LOUISE, *Princess* (9, 14). PRINCESS LOUISE (1, 2, 3, 4, 5, 6, 11, 12). *Princess Louise* (7, 10, 13). WOOLVERTON (7). *Woolverton* (4, 11, 12).

Louise is an apple of the Fameuse group. It is larger than Fameuse but not as large as McIntosh and has less red color and is less attractive than either of these apples. It is very desirable for dessert use but for culinary purposes it is decidedly inferior to other varieties of its season. It is of a clear pale yellow color with lively blush and delicate bloom. It shows bruises readily and requires very careful handling. The tree is hardy, healthy, comes into bearing rather young and yields fair to moderately good crops almost annually. It is doubtful whether it will ever prove a desirable commercial variety.

Historical. Originated with L. Woolverton, Grimsby, Ontario, who states that it was a chance seedling that sprung up among a half dozen old Fameuse trees.¹ It was first exhibited before the Ontario Fruit Growers' Association in 1879 under the name of "Woolverton." It has been disseminated to a limited extent only in New York. The McIntosh has sometimes been mistakenly disseminated for this variety.

TREE.

Tree medium size, moderately vigorous to vigorous with long, slender branches and willowy laterals. *Form* upright spreading to roundish, rather dense. *Twigs* moderately long; straight or somewhat irregularly curved, rather geniculate, slender; internodes medium. *Bark* reddish-brown with some pale olive-green, lightly mottled with scarf-skin; somewhat pubescent toward the tips. *Lenticels* scattering, small to medium, roundish to oblong. *Buds* small to medium, deep-set, plump, acute, free or nearly so, somewhat pubescent.

FRUIT.

Fruit of good medium size; fairly uniform in size and shape. *Form* usually roundish, sometimes roundish oblate, often somewhat elliptical or obscurely angular; sides often unequal. *Stem* red, medium to rather long, usually slender. *Cavity* obtuse to acute or sometimes approaching acuminate, shallow to medium in depth, moderately broad, frequently furrowed. *Calyx* medium size, closed or sometimes slightly open. *Basin* usually shallow to medium in depth, narrow to moderately wide, rather obtuse, lightly furrowed and wrinkled.

Skin thin, rather tough, clear pale yellow or greenish, with lively red or pinkish blush, striped obscurely if at all, overspread with thin bloom. *Dots* inconspicuous, pale, usually submerged.

Calyx tube short, wide, conical to funnel-form. *Stamens* median to basal.

¹ L. Woolverton, Letter, 1894.

Core medium to rather large, somewhat abaxile; cells symmetrical, partly open; core lines meeting. *Carpels* elongated ovate, sometimes emarginate. *Seeds* medium or below, moderately long, acute to acuminate.

Flesh whitish, not very firm, fine, crisp, very tender, very juicy, mild subacid, aromatic, with some of the perfume and texture of McIntosh, very good for dessert.

Season October to February or later.

LOWELL.

REFERENCES. 1. *Cultivator*, 5:246. 1848. 2. Cole, 1849:109. 3. Thomas, 1849:157. 4. Waring, 1851:27. 5. Barry, 1851:285. 6. Elliott, 1854:88. fig. 7. Hooper, 1857:55. 8. Gregg, 1857:46. fig. 9. Downing, 1857:166. 10. *Am. Pom. Soc. Cat.*, 1862. 11. Warder, 1867:576. 12. Fitz, 1872:145. 13. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 14. Bailey, *Am. Hort.*, 1892:243. 15. *Mich. Sta. Bul.*, 105:108. 1894. 16. Lyon, *Ib.*, 118:60. 1895. 17. Burrill and McCluer, *Ill. Sta. Bul.*, 45:330. 1896. 18. *N. C. Bd. Agr. Bul.*, 1900:7. 19. *Kan. Sta. Bul.*, 106:53. 1902. 20. *Mo. Fr. Sta. Bul.*, 3:27. 1902. 21. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:48. 1903. 22. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 23. Budd-Hansen, 1903:117. 24. Beach and Clark, *N. Y. Sta. Bul.*, 248:130. 1904.


SYNONYMS. *Greasy Pippin* (6, 9, 16, 22, 24, of some 23). *LOWELL* (2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24). *Lowell* (1). *LOWELL* (Pippin) (14). *Orange* (1, 2, 3, 5, 7, 8, 9, of some 6). *Pound Royal* (2, 7, erroneously 6). *Queen Anne* (1, 2, 6, 7, 9, 11, of Northern Ohio 3, of Ohio 5). *Tallow* (2, 7). *TALLOW APPLE* (1). *Tallow Apple* (3, 6, 9, of some 23). *Tallow Pippin* (3, 5, 6, 11, 15, 24).

Fruit rather large; clear yellow with waxy surface. *Flesh* a little coarse, very juicy, sprightly subacid and desirable for either dessert or culinary uses. It ripens in succession through a period of several weeks and is apt to drop as it ripens. Its season extends from late August to October and under favorable conditions a portion of the fruit may be kept till winter (24). The tree is a good grower, hardy, long-lived and a reliable cropper giving good crops biennially and sometimes annually. This fruit is grown for home use and to a limited extent for local market.

Historical. Lowell is an old variety of American origin (6) which is quite generally known in many parts of the state, especially in Western New York. It is still commonly listed by nurserymen, but is being planted but little and its cultivation appears to be on the decline.

TREE.

Tree rather large, vigorous. *Form* upright spreading, rather open. *Twigs* long, curved, moderately stout; internodes medium. *Bark* dark brown,



Vol. 11 - 12

LOWELL

mottled with scarf-skin; heavily pubescent. *Lenticels* numerous, small to medium size, not raised. *Buds* prominent, medium to large, broad, plump, obtuse, pubescent.

FRUIT.

Fruit large, fairly uniform in size but somewhat variable in shape. *Form* roundish oblong inclined to conic, unsymmetrical, irregular. *Stem* medium to rather long, moderately thick, deflected to one side. *Cavity* acute or approaching acuminate, rather shallow to moderately deep, medium to rather broad, sometimes furrowed, occasionally compressed, often lightly russeted. *Calyx* medium to large, closed or partly open. *Basin* rather shallow to moderately deep, medium in width, somewhat abrupt, often slightly furrowed and wrinkled.

Skin thin, tender, smooth or with occasional russet dots and flecks, waxy, at first green but eventually becoming rich yellow. *Dots* very numerous, small, inconspicuous, brown or russet or submerged.

Calyx tube usually rather long, rather wide, conical approaching funnel-form. *Stamens* median or nearly so.

Core rather large, axile to somewhat abaxile; cells closed; core lines meeting or somewhat clasping. *Carpels* roundish to pointed obovate, emarginate. *Seeds* dark brown, not uniform in shape, medium size, obtuse to somewhat acute.

Flesh greenish or tinged with yellow, rather firm, medium to rather fine-grained, crisp, tender, very juicy, sprightly subacid, good to very good.

Season August to October.

LOWLAND RASPBERRY.

REFERENCES. 1. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:438. 2. *Ib.*, *Montreal Hort. Soc. Rpt.*, 1883:94. 3. *Ib.*, *Am. Pom. Soc. Rpt.*, 1887:46. 4. Thomas, 1897:254. fig. 5. *U. S. Pom. Bul.*, 8:18. 1899. 6. Hansen, *S. D. Sta. Bul.*, 76:69. 1902. fig. 7. Budd-Hansen. 1903:117. fig.

SYNONYMS. HIMBEERAPFEL LIEVLANDER (1, 2). *Himbecerapfel* Lievlander (3). *Himbecerapfel* Lievlander (3). *Lievland Raspberry* (7). LIVELAND RASPBERRY (4). LIVLAND RASPBERRY (3). LOWLAND RASPBERRY (6). LOWLAND *Raspberry* (5, 7). *Lowland Raspberry* (1, 2, 3). No. 340 (1, 2, 6, 7).

A Russian apple which, according to Hansen (6, 7), is medium to large, clear waxen-white, striped, shaded and marbled with light crimson; flesh white, often stained with red, fine, very tender, pleasant mild subacid, almost sweet; season August; as early as Yellow Transparent.

We do not know that this variety has been tested in New York.

LUBSK QUEEN.

REFERENCES. 1. Tuttle, *Montreal Hort. Soc. Rpt.*, 8:136. 1881-82. 2. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:442. 3. Budd, *Ia. Agr. Coll. Bul.*, 1885:17. 4. *Am. Pom. Soc. Cat.*, 1889:8. 5. Budd, *Ia. Agr. Coll. Bul.*, 1890:19. 6. *Ia. Sta. Bul.*, 19:136. 1892. 7. *Can. Hort.*, 16:361. 1893. 8. Freeborn, *Nat. Nursery-*

man, 2:132. 1894. 9. Thomas, 1897:291. *fig.* 10. Hansen, *S. D. Sta. Bul.*, 76:70. 1902. *fig.* 11. Budd-Hansen, 1903:119. 12. Ragan, *U. S. B. P. I. Bul.*, 56:185. 1905.

SYNONYMS. LUBSK QUEEN (1, 2, 4, 5, 6, 7, 8, 9, 10, 11). *Lubsk Queen* (12). LUBSK REINETTE (3, 12). *Lubsk Reinette* (10, 11). *No. 444* (3, 5, 6, 8, 10, 11, 12). *Reinette Liubski* (2, 12). *Renet Liubskui* (3, 12).

A Russian variety which has attracted attention because of the beauty of the fruit and superior hardiness and productiveness of the tree. In other respects it does not rank high. It is fairly good in quality. Season August and September.

Historical. Imported from Russia by the United States Department of Agriculture in 1870 under No. 444. It was entered on the list of the American Pomological Society in 1889 and dropped from the list at the following meeting in 1891 (4). It has been planted but little in New York, and so far as we have been able to learn its cultivation in this state is not increasing.

TREE.

Tree moderately vigorous with long, slender, curved branches. *Form* upright spreading or roundish, rather open. *Twigs* short, curved, slender; internodes medium. *Bark* dull brown, roughly mottled with scarf-skin; slightly pubescent near tips. *Lenticels* scattering, medium size, round, not raised. *Buds* medium size, plump, obtuse, free, not pubescent.

FRUIT (10).

Fruit medium to large. *Form* regular, nearly round, truncated. *Stem* medium to long. *Cavity* acute, rather small, slightly russeted. *Calyx* closed; segments long, pointed. *Basin* shallow, wide, regular, with generally five fine corrugations around the eye.

Skin very smooth, polished and waxlike, a brilliant white, more or less covered with solid light rosy red, with delicate white bloom; a self-colored apple, but sometimes with short red splashes on lighter ground. *Dots* white, minute, numerous.

Calyx tube long, funnel-shape. *Stamens* marginal.

Core closed; cells ovate, slit. *Seeds* nine, plump.

Flesh snow white, firm, juicy, fine-grained, subacid, good.

Season August, September.

LYSCOM.

REFERENCES. 1. *N. E. Farmer*, 1830 (cited by 17). 2. Kenrick, 1832:36. 3. *Mag. Hort.*, 1:364. 395. 1835. 4. Hovey, *Ib.*, 4:48. 1838. 5. Manning, 1838:51. 6. *Ib.*, *Mag. Hort.*, 7:46. 1841. 7. Downing, 1845:89. 8. Cole, 1849:111. 9. Thomas, 1849:151. 10. Emmons, *Nat. Hist. N. Y.*, 3:32. 1851. 11. Elliott, 1854:145. 12. *Mag. Hort.*, 20:241. 1854. 13. Hooper, 1857:56. 14. *Am. Pom. Soc. Cal.*, 1862. 15. Warder, 1867:605. 16. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 17. Ragan, *U. S. B. P. I. Bul.*, 56:186. 1905.

SYNONYMS. LYSKOM (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17). *Matthew's Stripe* (3, 8, 17). *Osgood's Favourite* (3, 7, 8, 9, 17).

A very large apple somewhat resembling Twenty Ounce in general appearance but less attractive; flesh tender, mild, not high in flavor but acceptable for either dessert or culinary uses. The tree is a moderate grower and yields moderate to good crops. It is in season from late September or October to November or December. Not recommended for planting in New York because it is not superior to other varieties of its season.

Historical. Origin Southborough, Worcester county, Mass. (2). It is an old variety which has been quite widely disseminated. It is now nearly obsolete in New York.

FRUIT.

Fruit large or very large. *Form* roundish varying from a little oblate to slightly oblong conic, often broadly ribbed especially toward the apex. *Stem* deep set, short, moderately thick. *Cavity* acuminate, very deep, broad, symmetrical, russeted and with outspreading russet rays. *Calyx* medium to large, usually closed, pubescent. *Basin* often oblique, deep, moderately narrow to rather wide with broad, deep furrows.

Skin thick, tough, rather dull green or somewhat yellowish, striped, splashed and somewhat blushed with red.

Calyx tube wide, elongated cone-shape. *Stamens* median to basal.

Core rather large, axile; cells closed; core lines meeting or slightly clasping. *Carpels* ovate, slightly emarginate, somewhat tufted. *Seeds* short, often nearly round, plump, obtuse.

Flesh somewhat tinged with yellow, rather fine, tender, juicy, subacid becoming very mild subacid, good.

Season late September or October to November or December.

MABIE.

A red sweet apple of very good general appearance. It belongs in the same group as Victoria Sweet. Season late fall and early winter. A desirable variety of its class.

Historical. This is a local variety which has been grown to a limited extent in southern Rockland county. The following account of its origin is given by M. L. Bell, Sparkill, N. Y.:¹ "Mabie has been grown in Southern Rockland county, N. Y., and the adjacent portion of New Jersey for about forty-five years. About fifty years ago the original tree stood in an old stone fence row on the farm of Wm. Mabie. It was propagated in a local nursery and disseminated through the surrounding region where it is generally highly esteemed by those who have tested it."

FRUIT.

Fruit above medium to nearly large. *Form* roundish or somewhat inclined to oblong, a little irregular, unsymmetrical, not very uniform. *Stem* medium

¹ Letter, 1904.

to, rather long, moderately thick. *Cavity* large, acute to acuminate, moderately deep, wide, often obscurely furrowed, sometimes completely covered with russet which extends over the base of the fruit in broken rays. *Calyx* medium to rather large, closed or partly open. *Basin* medium size or below, pubescent, saucer-shape, medium in width and depth, a little obtuse to rather abrupt, smooth or slightly furrowed.

Skin very thick, tough, smooth or slightly roughened with russet dots, bright yellow mottled or deeply blushed with bright red, in highly colored specimens rather deep, dark red, obscurely striped with carmine and marked over the base with whitish scarf-skin. *Dots* often very small, gray, mingled with others that are large, whitish and areolar with russet center.

Calyx tube short, moderately wide, conical to funnel-form. *Stamens* median.

Core medium to rather large, axile to somewhat abaxile; cells closed or somewhat open; core lines clasping. *Carpels* roundish to broadly obovate, slightly tufted. *Seeds* dark brown, medium to small, plump, broadly acute.

Flesh slightly tinged with yellow, firm, medium grained, somewhat crisp, tender, juicy, sweet, good.

Season late fall and early winter.

McCARTY.

This appears to be a distinct strain or sport of Pumpkin Sweet. It is discussed under Pumpkin Sweet, page 173.

MAC DONOUGH.

REFERENCE. 1. Waugh, *Vt. Sta. An. Rpt.*, 14:299. 1901.

SYNONYMS. None.

A medium sized yellow apple of mild flavor, in season in August and September. We are unacquainted with this variety. Waugh (1) gave the following account of it in 1901: "A local variety, said to have originated on Cumberland Head, Clinton county, N. Y., opposite Grand Isle. A fairly good apple in many ways, but not common and not likely ever to become popular.

"Fruit roundish oblate, size medium, cavity medium deep and broad, waxy, slightly russeted, stem medium straight, basin shallow, corrugated, calyx nearly closed, color greenish-yellow, fine yellow when ripe, dots many, light greenish, bloom waxy, skin smooth, flesh greenish-white, mealy, core medium, closed, flavor neutral, nearly sweet, quality fair to good, season August-September. Tree hardy, rough dark bark, irregular in form, productive."

McINTOSH.

REFERENCES. 1. Downing, 1876:55 app. fig. 2. *Montreal Hort. Soc. Rpt.*, 5:17. 1879. 3. *Am. Pom. Soc. Cat.*, 1883:12. 4. Thomas, 1885:517. 5. Hoskins, *Mich. Hort. Soc. Rpt.*, 1886:220. 6. *Ib.*, *Rural N. Y.*, 47:558. 1888. 7. Budd, *Can. Hort.*, 13:24. 1890. 8. *Ib.*, 15:19, 92, 124. 1892. 9. Bailey,



McINTOSH

An. Hort., 1892:244. 10. *Can. Hort.*, 16:369. 1893. *col. pl.* 11. Hoskins, *Am. Gard.*, 15:288. 1894. 12. Craig, *Can. Dept. Agr. Rpt.*, 1894:125. 13. Hoskins, *Rural N. Y.*, 53:279. 1894. 14. *Ib.*, 54:238. 1895. 15. *Ib.*, 55:656. 1896. 16. *Ib.*, 55:738, 819. 1896. 17. *Ib.*, 56:839. 1897. 18. *Can. Hort.*, 22:506. 1899. 19. Whitney, *Ont. Fr. Gr. Assn. An. Rpt.*, 31:12. 1899. 20. *Ib.*, *Can. Hort.*, 23:24. 1900. *fig. of tree.* 21. *Ib.*, 23:45. 1900. *figs.* 22. *Vt. Sta. Bul.*, 83:91. 1900. *figs.* 23. Taylor, *U. S. Dept. Agr. Yr. Bk.*, 1901:383. *col. pl.* 24. Waugh, *Vt. Sta. An. Rpt.*, 14:299. 1901. 25. *Ib.*, *Rural N. Y.*, 61:781. 1902. *figs.* 26. *Kan. Sta. Bul.*, 106:54. 1902. 27. Hansen, *S. D. Sta. Bul.*, 76:72. 1902. 28. Budd-Hansen, 1903:119. 29. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 30. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:48. 1903. 31. Beach and Clark, *N. Y. Sta. Bul.*, 248:130. 1904. 32. Scriber, *Can. Hort.*, 28:277. 1905.

SYNONYMS. MACINTOSH RED (18). MCINTOSH (14, 16, 17, 19, 21, 25, 29). MCINTOSH RED (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 20, 26, 32). *McIntosh Red* (22, 23, 24, 27, 28, 30, 31).

This variety belongs in the Fameuse group. It is adapted to a wider range of localities than is the Fameuse. The fruit is very attractive in appearance, of bright deep red color and good size. The flesh is very tender, perfumed and delicious. It is desirable for local markets and special trade but because of its lack of firmness it is less suitable for general handling. As grown at this Station it begins to ripen in late September or early October. In Western New York it cannot be expected to keep much later than October in ordinary storage without considerable loss but in cold storage it may be held until December or January (31). When grown in more northern or elevated regions it is often held in good condition till midwinter or later. It is susceptible to scab but this may readily be controlled with proper treatment.¹ The crop ripens unevenly and a considerable portion of the fruit is liable to drop before it is ready to pick. On this account it is best to make two or three pickings. In some localities the tree is said to be a somewhat slow grower and not satisfactorily productive, but more often it is found to be a rather strong grower, hardy and healthy. It comes into bearing rather young and is a reliable cropper yielding good crops biennially and sometimes annually. It has not been sufficiently tested to demonstrate fully its value for commercial purposes but it is regarded by many as one of the most promising varieties of its class for general cultivation in New York.

¹ *N. Y. Sta. An. Rpt.*, 18:399-418. 1899. *Ib.*, 22:321-386. 1903.

Historical. Originated as a chance seedling on the McIntosh homestead, Matilda township, Dundas county, Ontario, where Allan McIntosh began the propagation of this variety in the nursery about 1870 (20). It has been widely disseminated. It is now commonly propagated by nurserymen and its cultivation is on the increase in New York.

TREE.

Tree vigorous with numerous, small, slender laterals. *Form* roundish or spreading. *Twigs* above medium to short, straight or nearly so, rather slender; internodes long to below medium. *Bark* bright reddish-brown, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* quite numerous, small, oval or elongated, raised. *Buds* deeply set in bark, medium to below, plump, obtuse to acute, free, slightly pubescent.

FRUIT.

Fruit above medium, sometimes large, pretty uniform in shape and size. *Form* roundish to somewhat oblate, regular or faintly ribbed, obscurely angular. *Stem* short, stout or moderately slender, usually not exserted, often with irregular protuberances. *Cavity* large, acuminate or somewhat acute, wide, medium in depth, somewhat broadly furrowed, often partly russeted. *Calyx* small, closed or partly open; lobes short to long, narrow, acute. *Basin* pubescent, rather small, medium in depth, narrow, abrupt, smooth or obscurely furrowed.

Skin thin, moderately tender, smooth, readily separating from the flesh, clear whitish-yellow or greenish washed and deeply blushed with bright red and striped with carmine; highly colored specimens become dark, almost purplish-red with the carmine stripes obscure or obliterated, overspread with thin, lilac bloom. Often the effect of the deep red is heightened by lively contrast with one or more spots of the clear pale yellow ground color where some twig or leaf pressed closely against the growing fruit. *Dots* whitish or yellow, usually very small.

Calyx tube short, conical or funnel-shape with broad limb. *Stamens* median to basal.



Core medium size, usually abaxile; cells usually wide open; core lines nearly meeting. *Carpels* roundish to elliptical, narrowing toward base and apex, smooth, much concave. *Seeds* medium brown, rather large, acute.

Flesh white or slightly tinged with yellow, sometimes veined with red, firm, fine, crisp, tender, very juicy, characteristically and agreeably aromatic, perfumed, sprightly, subacid, becoming mild and nearly sweet when very ripe, very good to best for dessert.

Season October to December or later.

McLELLAN.

REFERENCES. 1. Leavenworth, *Horticulturist*, 2:26. 1847. fig. 2. Thomas, 1849:169. 3. Cole, 1849:125. 4. Emmons, *Nat. Hist. N. Y.*, 3:67. 1851. 5. Hovey, *Mag. Hort.*, 20:508. 1854. fig. 6. Elliott, 1854:147. 7. Downing, 1857:87. 8. Hooper, 1857:57. 9. Warder, 1867:726. 10. *Am. Pom. Soc.*



Mc LELLAN

Cat., 1871:8. 11. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 12. Taylor, *Am. Pom. Soc. Rpt.*, 1895:193. 13. Burrill and McCluer, *Ill. Sta. Bul.*, 45:309, 330. 1896. 14. Farrand, *Mich. Sta. Bul.*, 205:42. 1903. 15. Budd-Hansen, 1903:120. *fig.*

SYNONYMS. *Martin* (1, 2, 6, 7). *McClellan* (1, 5). *McLellan* (8, 9). *McClellan* (2, 3, 4, 6, 7, 10, 11, 12, 13, 14, 15).

A very choice dessert apple, handsome, fragrant, tender and excellent in quality. It comes into season a little later than Maiden Blush and may keep till January or February. When properly colored it is well adapted for fancy market and fruit-stand trade but when the color remains greenish, as it often does, the flavor is inferior. The fruit shows bruises readily and must be handled with extra care. It drops easily from the tree and on this account should be picked as soon as colored. In some cases it may pay to make two pickings. The tree comes into bearing young and is a reliable biennial bearer yielding good crops. It is only a moderately vigorous grower and probably it would be an advantage to topwork it upon some more vigorous and longer-lived stock, such as Northern Spy or Baldwin. It is not recommended for extensive commercial planting but in some cases it may be grown to a limited extent with profit.

Historical. The original tree was planted in a seedling orchard in Woodstock, Conn., about 1780 (1, 5). It has been sparingly disseminated and is known locally in various parts of New York state, but it is now seldom offered by nurserymen (12) and is little propagated.

TREE.

Tree medium in size, moderately vigorous with long and moderately stout branches. *Form* erect, roundish, open. *Twigs* medium to long, erect, stout, generally somewhat curved; internodes short to medium. *Bark* dull, very dark reddish-brown, sometimes tinged with green and overlaid with rather heavy scarf-skin; rather heavily pubescent. *Lenticels* quite numerous, not very conspicuous, medium to small, roundish, sometimes a little raised. *Buds* above medium to large, rather deeply set in the bark, broad, flat, obtuse to somewhat acute, free, very pubescent.

FRUIT.

Fruit above medium to large; uniform in size and shape. *Form* roundish oblate to roundish conic, rather broad and flat at the base, symmetrical or nearly so, regular to elliptical. *Stem* short to medium, slender. *Cavity* acute to acuminate, rather wide, deep, symmetrical, sometimes a little fur-

rowed, smooth. *Calyx* small to medium, partly open or closed; lobes long, acuminate. *Basin* medium to rather small, abrupt, moderately shallow to deep, rather narrow to moderately wide, nearly round or sometimes angular, sometimes distinctly furrowed and wrinkled.

Skin tough, waxen, pale yellow or greenish, blushed and mottled with bright light red splashed and striped with bright carmine. Highly colored specimens are almost entirely red and very attractive. *Dots* numerous but mostly inconspicuous, whitish or russet.

Calyx tube funnel-form with wide limb or sometimes elongated. *Stamens* basal to median.

Core below medium to small, axile or nearly so; cells not uniformly developed, usually pretty symmetrical, closed or partly open; core lines clasping the funnel cylinder. *Carpels* smooth, roundish to elliptical narrowing toward the base and apex, slightly emarginate. *Seeds* rather small, obtuse, smooth, dark.

Flesh tinged with yellow, moderately firm, very tender, moderately fine-grained, juicy, moderately crisp, almost sweet, excellent in flavor, very good for dessert.


Season October to January or February or sometimes to March.

McMAHON.

REFERENCES. 1. *Am. Pom. Soc. Rpt.*, 1885:27. 2. *Mich. Hort. Soc. Rpt.*, 1886:221. 3. *Am. Pom. Soc. Rpt.*, 1887:93. 4. *Rural N. Y.*, 46:751. 1887. 5. *Can. Hort.*, 11:220. 1888. fig. 6. *Am. Pom. Soc. Cat.*, 1889:10. 7. *Am. Gard.*, 11:243. 1890. 8. *Can. Hort.*, 13:174. 216. 1890. 9. Van Deman, *U. S. Pom. Rpt.*, 1890:413. 10. *Can. Hort.*, 14:339. 1891. 11. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57. 59. 12. Bailey, *Am. Hort.*, 1892:244. 13. *Can. Hort.*, 15:393. 1892. 14. *Ib.*, 16:77. 134. 1893. col. pl. 15. Craig, *Ib.*, 16:137. 1893. fig. 16. *Ib.*, *Ont. Fr. Gr. Assn. An. Rpt.*, 26:16. 1894. 17. *Ib.*, *Can. Dept. Agr. Rpt.*, 1894:125. 18. *Ib.*, 1895:93. fig. 19. Beach, *N. Y. Sta. An. Rpt.*, 15:273. 1896. figs. 20. Thomas, 1897:625. 21. Waugh, *Vt. Sta. An. Rpt.*, 14:300. 1901. 22. Hansen, *S. D. Sta. Bul.*, 76:72. 1902. fig. 23. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:48. 1903. 24. Farrand, *Mich. Sta. Bul.*, 205:42. 1903. 25. Budd-Hansen, 1903:121. fig. 26. Beach and Clark, *N. Y. Sta. Bul.*, 248:130. 1904. 27. Ragan, *U. S. B. P. I. Bul.*, 56:188. 1905.

SYNONYMS. McMAHAN (14, 20). McMahan (27). McMAHAN WHITE (6, 7, 15, 16, 17, 18). McMAHON (11, 19, 21, 22, 23, 25, 26, 27). McMAHON WHITE (1, 2, 3, 4, 5, 8, 9, 10, 12, 13, 24). McMahan White (19, 21, 22, 23, 25, 26, 27).

A large apple of the Apert group, pale yellow or almost white, often with a delicate pink blush. The flesh is juicy, brisk subacid, fair to good in quality, excellent for culinary use. As grown at this Station it ripens unevenly and does not appear well adapted for storage (26). Season October to January or February. The



LEE



Mc LELLAN



McMAHON

tree comes into bearing rather young and yields good crops biennially. In 1895 Craig wrote concerning McMahon: "Attention is again drawn to some of its merits as an apple of value for regions where Northern Spy, Ribston Pippin and Rhode Island Greening cannot be grown profitably on account of their inability to withstand the severity of the climate. It has proved so far a remarkably vigorous and healthy grower, making probably more well matured wood growth than any other variety in the orchard, and is free from many of the defects characteristic to varieties unadapted to this climate. It has borne moderately heavy crops for the last two years. The fruit is of the largest size, smooth and handsome, though lacking in color as grown in this vicinity and somewhat soft in texture" (18). It is less desirable than standard kinds of its season for growing commercially in New York because it is only moderately attractive in general appearance, its color is such that it readily shows bruises and it does not rank high either in flavor or quality.

Historical. This variety originated about 1860 in Richland county, Wis., and is supposed to be a seedling of the Alexander. It is frequently listed by nurserymen, but it has thus far been but little planted in New York.

TREE.

Tree medium size, vigorous. *Form* rather spreading. *Twigs* medium size, slender, straight or nearly so; internodes above medium. *Bark* dull dark reddish-brown; slightly pubescent. *Lenticels* numerous, below medium or sometimes medium, elongated, somewhat raised, dull. *Buds* medium or below, rather acute, thin, often appressed, slightly pubescent.

FRUIT.

Fruit large or very large. *Form* roundish inclined to conic, faintly ribbed. *Stem* medium in length, moderately thick. *Cavity* remarkably acuminate, very deep, broad, compressed, slightly russeted and with outspreading rays. *Calyx* rather small, slightly open; lobes separated at base, short. *Basin* moderately deep, rather narrow to moderately wide, abrupt, compressed, slightly furrowed and wrinkled.

Skin pale greenish-yellow or nearly white with irregular stripes and patches of whitish scarf-skin extending out from the cavity over the base, and often having the cheek overspread with a thin blush which sometimes is faintly splashed and striped with carmine. *Dots* few, inconspicuous, small, greenish or russet.

Calyx tube rather long, wide, broadly conical inclined to funnel-form or cylindrical. *Stamens* median to basal.

Core medium to small, slightly abaxile to axile; cells pretty symmetrical, closed or slightly open; core lines nearly meeting or clasping. *Carpels* roundish or inclined to elliptical, slightly emarginate. *Seeds* rather dark brown, medium or below, moderately plump, obtuse to acute.

Flesh nearly white, nearly fine, tender, juicy, sprightly subacid, fair to good.

Season October to January or February.

MAGOG.

REFERENCES. 1. Downing, 1876:56 app. 2. Thomas, 1885:516. 3. *Can. Hort.*, 13:174. 1890. 4. Bailey, *Am. Hort.*, 1892:243. 5. Hoskins, *Am. Gard.*, 15:256. 1894. 6. Beach, *N. Y. Sta. An. Rpt.*, 15:273. 1896. 7. *Am. Pom. Soc. Cat.*, 1899:18. 8. Powell and Fulton, *U. S. B. P. l. Bul.*, 48:48. 1903. 9. Budd-Hansen, 1903:122. 10. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 11. Beach and Clark, *N. Y. Sta. Bul.*, 248:130. 1904.


SYNONYMS. MAGOG (4, 8, 10). MAGOG *Red Streak* (7, 9). MAGOG RED STREAK (1, 2, 3, 5, 6). *Magog Red Streak* (4, 11).

Fruit of good size, rather attractive appearance and good to very good quality, especially for culinary uses. Dr. Hoskins, who propagated it for several years, said in 1894: "It is a fair but uneven keeper, and might be well described as not quite valuable enough to retain, yet hardly deserving to be cast aside." It is reported as being very hardy in Northern New York, a good grower and a good bearer. As fruited at this Station the tree is a moderately good grower, comes into bearing rather young and yields moderate to good crops nearly annually. The keeping quality of the fruit varies in different seasons but its commercial limit in ordinary storage appears to be October. It remains in season to January or possibly later (11). It is not recommended for planting except perhaps for home use in localities where its superior hardiness gives it an advantage over ordinary varieties of its season.

Historical. Originated by Wm. Warren, Newport, Vt. (1). It has been planted to a considerable extent in portions of Northern New England, Northern New York and the parts of Canada adjoining. It is still propagated in a few nurseries (4).

TREE.

Tree moderately vigorous. *Form* upright spreading or roundish. *Twigs* short to medium, straight or nearly so, rather slender; internodes short to medium. *Bark* dark brown or brownish-red, lightly streaked with scarf-skin, pubescent. *Lenticels* scattering, small, oval, not raised. *Buds* small to above medium, plump, obtuse, free, slightly pubescent.



MAGOG

FRUIT.

Fruit medium to large, averaging above medium, rather uniform in size but variable in shape. *Form* roundish to oblong, inclined to conic or somewhat ovate, regular or faintly ribbed; sides often unequal. *Stem* medium to short, moderately thick to rather slender. *Cavity* acute or approaching acuminate, medium in depth, medium to rather narrow, usually smooth, occasionally lipped, often irregularly russeted. *Calyx* medium to small, closed; lobes medium length, rather narrow, acute to acuminate. *Basin* usually medium in width and depth, sometimes rather abrupt, coarsely wrinkled.

Skin thin, tough, smooth, waxy, pale greenish or yellow, lightly washed and mottled with thin brownish-red, sparingly striped and splashed with deeper red. *Dots* numerous, light, submerged, areolar, brown and russet. *Prevailing effect* yellow.

Calyx tube medium in width, long, conical to funnel-shape with long cylinder. *Stamens* nearly marginal.

Core large to medium, abaxile; cells open; core lines clasping the funnel cylinder. *Carpels* long ovate to broadly obcordate, sometimes tufted. *Seeds* light brown, small to medium, rather wide, short, very plump, obtuse to broadly acute.

Flesh tinged with yellow, rather firm, medium to rather fine-grained, tender, very juicy, sprightly, pleasant subacid, aromatic, good.

Season October to January or later.

MAIDEN BLUSH.

REFERENCES. 1. Cox, 1817:106. fig. 2. Buel, N. Y. Bd. Agr. Mem., 1826:476. 3. Wilson, 1828:136. 4. Fessenden, 1828:131. 5. London Hort. Soc. Cat., 1831:No. 704. 6. Kenrick, 1832:27. 7. Thacher, 1822:130. 8. Mag. Hort., 1:363, 396. 1835. 9. Downing, 1845:90. fig. 10. Thomas, 1849:157. fig. 11. Cole, 1849:113. fig. 12. Emmons, Nat. Hist. N. Y., 3:40. 1851. col. pl. No. 30 and fig. 13. Barry, 1851:285. 14. Hovey, Mag. Hort., 18:544. 1852. fig. 15. Elliott, 1854:469 app. 16. Hooper, 1857:57. 17. Am. Pom. Soc. Rpt., 1860:240. 18. Warder, 1867:412. fig. 19. Fitz, 1872:143, 164. 20. Hogg, 1884:138. 21. Rural N. Y., 47:713. 1888. 22. Wickson, 1889:244. 23. Lyon, Mich. Hort. Soc. Rpt., 1890:294. 24. Bailey, An. Hort., 1892:243. 25. Can. Hort., 15:65. 1892. 26. Ark. Sta. An. Rpt., 6:55. 1893. 27. *Ib.*, 7:44. 1894. 28. Hoskins, Rural N. Y., 53:279. 1894. 29. *Ib.*, 55:7. 1896. 30. Ill. Sta. Bul., 45:331. 1896. 31. Waugh, Vt. Sta. An. Rpt., 14:299. 1901. 32. Va. Sta. Bul., 130:124. 1901. 33. Am. Pom. Soc. Cat., 1901:18. 34. Powell and Fulton, U. S. B. P. I. Bul., 48:48. 1903. 35. Budd-Hansen, 1903:122. fig. 36. Beach and Clark, N. Y. Sta. Bul., 248:130. 1904.

SYNONYMS. *Lady Blush* (36). MAIDEN BLUSH (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36). *Red Check* (11).

This is a beautiful apple of pale lemon-yellow color with crimson check. The flesh is white, sprightly, not superior in flavor but

good in quality. It is valued especially for market and culinary uses. It makes very white evaporated stock. As grown in Western New York it is in season from September to November or December. Later than this although the fruit may appear sound it is deficient in quality. In cold storage its commercial limit appears to be about December 15th. It does not stand heat well before going into storage. It varies greatly in the time of maturing in different seasons. The earlier it matures the less satisfactory is it as a keeper (36). It is recognized as a standard market variety and usually sells above the average prices for varieties of its class. In many localities it has proven a very satisfactory variety for the commercial orchard, because the tree is a fine grower, hardy, pretty long-lived, comes into bearing rather young and is a reliable cropper, yielding good to heavy crops biennially or almost annually. In many cases the fruit does not mature uniformly and there is considerable loss from drops unless more than one picking is made in gathering the crop. Usually the fruit is pretty uniform in size but on unthrifty, old trees or under unfavorable circumstances a considerable amount of the fruit may be too small for market. It is sometimes badly injured by scab but this may readily be prevented by proper treatment.

Historical. Coxé described this variety in 1817 as very popular in the Philadelphia market and the best variety of its season for evaporating. He stated that it was named by Samuel Allinson, of Burlington, N. J., who first brought it to notice. In the American Pomological Society's Catalogue of fruits it is reported as either "wholly successful or successful" in nearly all the important apple-growing districts of the United States (33). It has long been well and favorably known in New York and it is still being planted both for commercial purposes and for home use.

TREE.

Tree medium size, moderately vigorous to vigorous. *Form* spreading, open. *Twigs* long, curved, slender; internodes short. *Bark* brown or reddish-brown, lightly mottled with scarf-skin; slightly pubescent. *Lenticels* quite numerous, small, round, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium or above, sometimes large, uniform in size and shape. *Form* oblate a little inclined to conic, regular, symmetrical. *Stem* short to



MAIDEN BLUSH

medium, rather slender. *Cavity* rather large, acute to obtuse, medium to wide, moderately deep to shallow, usually symmetrical, sometimes russeted. *Calyx* medium size, closed; lobes separated at base, medium length, moderately broad, acute. *Basin* moderately shallow, medium to wide, obtuse, regular, smooth or slightly furrowed, symmetrical.

Skin thin, tough, smooth, pale waxen yellow with crimson blush. *Dots* numerous, whitish and submerged or areolar.

Calyx tube small, narrow, conical to funnel-shape. *Stamens* median to marginal.

Core medium size, axile or somewhat abaxile, broadly elliptical; cells closed or slightly open; core lines meeting or slightly clasping. *Carpels* very broadly ovate. *Seeds* medium brown, moderately wide, moderately long, plump, acute.

Flesh white or with slight yellow tinge, fine, moderately crisp, tender, very juicy, subacid, good in quality especially for culinary uses.

Season September to November or December.

MAIDEN FAVORITE.

REFERENCES. 1. Downing, 1857:167. 2. Warder, 1867:725. 3. Thomas, 1875:505.

SYNONYMS. *Maiden's Apple* (1). MAIDEN FAVORITE (1, 2, 3).

A late fall and early winter variety which originated in Stuyvesant, Columbia county, N. Y. According to Downing it is a desirable amateur variety of delicacy and beauty, medium size or below, whitish or pale waxen yellow sometimes mottled with crimson. Flesh tender, crisp, very delicate, vinous, sweet, good to very good. Tree a rather slow grower with upright, slender branches, and a good bearer. We are not acquainted with this variety. So far as we have been able to learn it is not cultivated outside of the vicinity of its origin and is no longer propagated.

MARGARET.

REFERENCES. 1. Forsyth, 1803:50. 2. *Am. Gard. Cal.*, 1806:584. 3. Forsyth, 1824:114. 4. *Pom. Mag.*, 1:No.46. 1828. col. pl. 5. *London Hort. Soc. Cat.*, 1831:No. 708. 6. Kenrick, 1832:26. 7. Floy-Lindley, 1833:7. 8. Manning, 1838:46. 9. Downing, 1845:73. fig. 10. Cole, 1849:98. 11. Thomas, 1849:137. fig. 12. Emmons, *Nat. Hist. N. Y.*, 3:11. 1851. 13. Elliott, 1854:145. 14. Hooper, 1857:31. 15. Warder, 1867:717. 16. Fitz, 1872:160. 17. Downing, 1872:10 index, app. 18. Leroy, 1873:454. fig. 19. Barry, 1883:332. 20. Hogg, 1884:141. 21. Bailey, *Am. Hort.*, 1892:238. 22. *Jour. Roy. Hort. Soc.*, 1898:354. 23. Waugh, *Vt. Sta. An. Rpt.*, 14:299. 1901. 24. Budd-Hansen, 1903:124.

SYNONYMS. *D'Eve* (18). *Duverson's June* (17). *Early June* of South (13). *EARLY MARGARET* (16). *Early Margaret* (4, 20). *Early Red* (24). *EARLY RED JUNEATING* (6). *Early Red Juneating* (7, 9, 11, 12, 13, 20). *EARLY RED MARGARET* (4, 7, 8, 9, 10, 11, 12, 14, 15, 17, 19, 21). *Early Red Margaret* (6, 13, 18, 20, 23). *Early Striped Juneating* (4, 5, 6, 7, 20). *Eve Apple* (7, of Ireland 4, 5, 6, 9, 12, 13, 20). *Herr's June* (17). *June* of

some in Ohio (13). *Lammas* (3, 20). *Magdalene* (20). *MARGARET* (1, 2, 3, 13, 20, 22, 23, 24). *Margaret* (4, 5, 6, 7, 9, 12, 18). *Margaret, Early* (5). *MARGARET, EARLY RED* (5). *Margaretha Apfel* (4, 7, 9, 12, 13). *Marget Apple* (20). *MARGUERITE* (18). *Marguerite* (5, 20). *Maudlin* (18, 20). *Red Joaneting* (22). *Red June of South* (13). *Red Juneating* (4, 5, 6, 7, 9, 10, 11, 12, 13, 20). *Red Juneting* (3). *Reinette Quarrendon* (18). *Rother Jacobs* (18). *Rother Jacobs Apfel* (4, 7). *Striped Juneating* (5, 9, 11, 12, 13, 18, 20). *Striped Quarendon* (17). *Striped Quarrenden* (5, 18, 20). *Summer Traveller* (5, 17, 20). *Virginia June* (17).

An early summer apple, small to nearly medium, roundish ovate, yellowish striped and marbled with dull red. Flesh pleasant subacid, good; season late July and early August. It is a dessert apple suitable for the home orchard only. The tree is of medium size, moderately vigorous, upright, hardy, long-lived, comes into bearing fairly young and is a pretty reliable biennial cropper.

Historical. This is a very old English variety. Hogg (20) states that "It is without doubt the Margaret of Rea, Worlidge, Ray, and all our early pomologists except Miller." It has long been cultivated in this country, but only to a very limited extent. It is now rarely propagated by our nurserymen and seldom planted.

MILLER.

REFERENCES. 1. Downing, 1857:172. 2. *Horticulturist*, 13:530. 1858. 3. Warder, 1867:726. 4. Downing, 1869:275. 5. Leroy, 1873:469. fig. 6. Thomas, 1875:506. 7. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294.

SYNONYMS. MILLER (1, 3, 4, 5, 6, 7). MILLER SEEDLING (2).

This apple was brought to notice by James O. Miller, Montgomery, Orange county, N. Y. (1). Downing (4) describes the fruit as rather large, roundish oblate inclined to conic, yellow shaded, striped and splashed with light red. Flesh yellowish, crisp, pleasant subacid, good to very good. Season October and November (3). Lyon refers to it as a promising fruit for market and general purposes (7). We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

MILLIGEN.

REFERENCES. 1. *N. Y. Sta. An. Rpt.*, 8:348. 1889. 2. *U. S. B. P. I. Bul.*, 48:49. 1903. 3. Beach and Clark, *N. Y. Sta. Bul.*, 248:132. 1904.

SYNONYMS. None.

Fruit of good size, rather attractive in general appearance, yellow striped with red, good to very good. It comes into season in October and some portion of the fruit may be kept in good condition till spring. In common storage there is apt to be a rather high rate of loss during late autumn, so that its commercial limit is October or early November, but it may be held in cold storage till midwinter (3). The tree is a vigorous grower, almost an annual bearer and usually produces good crops. It does not excel standard varieties of its season for any purpose. Not recommended for planting in New York.

Historical. This variety was originated by Mrs. Milligen, near Claysville, Washington county, Pa. It was received for testing at this Station from J. R. and R. A. Murdock, Pittsburg, Pa., in 1888.

MOSHER.

REFERENCES. 1. Downing, 1881:97 app. 2. Waugh, *Vt. Sta. An. Rpt.*, 14:300. 1901.

SYNONYMS. MOSHER (2). MOSHER SWEET (1). *Mosher Sweet* (2).

A good variety for the home orchard but not attractive enough for market (1). Originated in the orchard of Ephraim Mosher, Washington, N. Y., many years since. So far as we can learn this variety is not now being propagated by nurserymen and its cultivation appears to be declining. The tree is large, rather vigorous, upright spreading or roundish, a good grower, comes into bearing rather young and yields pretty good crops annually or nearly annually. Fruit medium, oblate conic, ribbed pale yellow or greenish; flesh white, moderately juicy, sweet, aromatic, good; season September and October.

MOTHER.

REFERENCES. 1. *Mag. Hort.*, 10:210. 1844. 2. Thomas, *Cultivator*, 5:306. 1848. fig. 3. *Horticulturist*, 3:581. 1848. 4. Cole, 1849:115. fig. 5. Hovey, *Mag. Hort.*, 15:65. 1849. fig. 6. Thomas, 1849:169. fig. 7. Waring, 1851:27. 8. Emmons, *Nat. Hist. N. Y.*, 3:58. 1851. 9. *Am. Pom. Soc. Cat.*, 1852. 10. *Mag. Hort.*, 20:241. 1854. 11. Elliott, 1854:147. 12. Downing, 1857:87. 13. Hooper, 1857:62. 14. Gregg, 1857:55. 15. *Mag. Hort.*, 28:281. 1862. 16. Lothrop, *Mag. Hort.*, 32:363. 1866. 17. Warder, 1867:663. fig. 18. Fitz, 1872:141. 19. Barry, 1883:350. 20. Hogg, 1884:7. 21. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 22. Bailey, *An. Hort.*, 1892:245. 23. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:356. 24. Alwood, *Va. Sta. Bul.*, 130:124. 1901. 25. Dickens and Greene, *Kan. Sta. Bul.*, 106:54. 1902. 26. *Rural N. Y.*, 61:829. 1902. figs. 27. Budd-Hansen, 1903:132. figs. 28. *Rural N. Y.*, 62:7, 380. 1903. 29. Bruner, *N. C. Sta. Bul.*, 182:21. 1903. 30. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:50. 1903. 31. Beach and Clark, *N. Y. Sta. Bul.*, 248:133. 1904.

SYNONYMS. AMERICAN MOTHER (20). *Gardener's Apple* (20). *Mother Apple* (20). MOTHER (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31). MOTHER OF AMERICA (23). *Queen Anne* (12, 20).

A beautiful red apple of good size, with tender, rich, aromatic flesh of best dessert quality. It is less desirable for culinary uses, being somewhat lacking in acidity. It resembles Esopus *Spitzenburg* to a marked degree but ripens earlier and is not so good a keeper. In ordinary storage it does not keep well and November

is its safe commercial limit, but in cold storage it may be held till March or later (31). The tree is below medium size and but a moderate grower. In many localities it is rather tender and liable to scald or canker on the trunk and larger branches. On this account it is desirable to topwork it upon Northern Spy or some other hardy, vigorous stock. It does not come into bearing very young and commonly is a biennial cropper yielding moderate to good crops. It cannot be recommended for general commercial planting but it is desirable for the home orchard.

Historical. Thomas described it in 1848 as "a new, handsome late autumn and early winter apple, of the highest quality," and stated that it originated at Bolton, Worcester county, Mass. (2). It is still listed by some nurserymen (22) but it is nowhere being extensively planted. It is but seldom found in cultivation in New York.

TREE.

Tree below medium size, moderately vigorous or a rather slow grower. *Form* upright spreading to roundish, rather open. *Twigs* long, curved, moderately stout; internodes long. *Bark* brown, mingled with olive-green, lightly mottled with scarf-skin; slightly pubescent near tips. *Lenticels* numerous, rather conspicuous, medium size, oval, raised. *Buds* rather prominent, medium size, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit below medium to above, sometimes large, pretty uniform in size and shape. *Form* roundish or roundish conic to oblong ovate, obscurely and broadly ribbed. *Stem* long to medium, moderately slender to rather thick. *Cavity* acute or approaching acuminate, rather shallow to moderately deep, rather narrow to moderately broad, often russeted, sometimes furrowed or compressed or lipped. *Calyx* small, closed or nearly so; lobes medium, narrow, acute. *Basin* shallow, narrow, a little abrupt, somewhat furrowed and wrinkled.


Skin thin, smooth, golden yellow nearly covered with bright deep red, marbled and striped with carmine. *Dots* rather small, inconspicuous, yellowish or pale and submerged.

Calyx tube rather long, funnel-form with wide limb and narrow cylinder. *Stamens* marginal.

Core medium to rather small, abaxile; cells nearly symmetrical, open or partly open; core lines clasping. *Carpels* broadly ovate to roundish, emarginate, mucronate. *Seeds* rather dark, medium or below, plump, acute to acuminate.

Flesh fine, tender, juicy, very mild subacid, aromatic, very good to best in flavor and quality.

Season late September to January; it is in its prime in November.



MOTHER

MOUNTAIN SWEET.

REFERENCES. 1. Warder, 1867:388. 2. Downing, 1869:282.

SYNONYMS. MOUNTAIN SWEET (1, 2). *Mountaineer* (1, 2).

Fruit of medium size, fairly good appearance and moderate to good quality, not superior to other varieties of its season and not recommended for planting in this state.

Historical. Origin Pennsylvania (1). It is but little known in New York and its cultivation in this state is not being extended.

TREE.

Tree medium or below, not a strong grower, with short, stout, curved branches. *Form* spreading or roundish, rather dense. *Twigs* short, straight, moderately stout; internodes short. *Bark* clear brown mingled with green, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, small, oblong, slightly raised. *Buds* medium size, plump, acute, free, slightly pubescent.

FRUIT.

Fruit variable, usually medium or above. *Form* roundish or roundish oblate, pretty regular, somewhat angular; sides unequal. *Stem* short to rather long, rather slender. *Cavity* nearly acuminate to somewhat obtuse, medium in depth, rather broad, russeted and with outspreading russet rays. *Calyx* small to rather large, closed or open; lobes separated at base, medium to long, rather narrow, acuminate. *Basin* moderately shallow to shallow, narrow to moderately wide, wavy.

Skin thick, rather tough, somewhat rough, clear pale yellow overlaid with faint pinkish blush and scattering stripes of deeper red. *Dots* numerous, inconspicuous, small, russet. *Prevailing effect* striped.

Calyx tube medium in length, rather wide, conical to somewhat funnel-form. *Stamens* median to basal.

Core medium size, abaxile; cells open; core lines meeting or slightly clasping. *Carpels* broadly ovate to elliptical, emarginate. *Seeds* very small, rather wide, plump, acute, rather light brown.

Flesh white, moderately fine, very tender, juicy, sweet, good.

Season September to December.

MOUSE.

REFERENCES. 1. Downing, 1845:117. 2. Thomas, 1849:182. 3. Emmons, *Nat. Hist. N. Y.*, 3:82. 1851. 4. Elliott, 1854:147. 5. Hooper, 1857:62. 6. Warder, 1867:727.

SYNONYMS. *Moose* (1, 2, 3, 4, 5). *Mouse* (1, 2, 3, 4, 5, 6).

An old variety which originated in Ulster county, N. Y., where it was formerly popular (1). According to Downing the fruit is large, roundish conical, pale greenish-yellow with brownish blush; flesh very white, fine-grained, delicate, mild subacid; good; season October to November.

This variety is unknown to us and we have received no reports concerning it from any of our correspondents.

MUNSON.

REFERENCES. 1. *Genesee Farmer*, 10:288. 1849. 2. Barry, 1851:285. 3. Humrickhouse, *Mag. Hort.*, 19:163. 1853. 4. Elliott, 1854:148. 5. Downing, 1857:174. 6. Gregg, 1857:44. 7. *Horticulturist*, 17:104, 150. 1862. 8. *Am. Pom. Soc. Cat.*, 1862. 9. Warder, 1867:388. fig. 10. Thomas, 1875:198. 11. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 12. Bailey, *Am. Hort.*, 1892:245. 13. Hoskins, *Rural N. Y.*, 53:278. 1894. 14. Burrill and McCluer, *Ill. Sta. Bul.*, 45:333. 1896. 15. Munson, *Me. Sta. Rpt.*, 1896:71. 16. *Ib.*, 18:89. 1902. 17. Budd-Hansen, 1903:133. fig. 18. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 19. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:50. 1903. 20. Beach and Clark, *N. Y. Sta. Bul.*, 248:133. 1904.

SYNONYMS. *Meachem Sweet* (5). *MUNSON* (17, 18, 19, 20). *MUNSON SWEET* (2, 3, 5, 8, 9, 10, 11, 12, 13, 14, 15, 16). *Munson Sweet* (17, 19, 20). *MUNSON SWEETING* (4, 6). *Munson Sweeting* (1). *NORTHERN SWEETING* (1). *Orange Sweet* (5, 9). *Ray Apple* (5).

In some parts of New York this variety is better known under either the names Meachem Sweet or Rag Apple than it is under the correct name of Munson.

The fruit commonly averages about medium size. It is attractive in appearance for a yellow apple, being pale yellow often somewhat blushed; skin characteristically tough separating readily from the tender flesh; season late September to December. It is a desirable variety for home use but is not very satisfactory for growing commercially because there is little demand for a sweet apple of its season except in certain local markets. The tree is a good grower, comes into bearing rather early and is a pretty reliable cropper commonly yielding good crops biennially.


Historical. Origin uncertain. Supposed to have originated in Massachusetts. It is still propagated by nurserymen (12) but it is not being planted in New York now as much as it was a generation ago.

TREE.

Tree large, moderately vigorous to vigorous. *Form* spreading or roundish, rather dense. *Twigs* short to rather long, curved, stout to moderately slender with large terminal buds; internodes short. *Bark* dark brown, heavily streaked with scarf-skin; pubescent near tips. *Lenticels* quite numerous, rather conspicuous, medium size, roundish, raised. *Buds* large, prominent, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit below medium to rather large, averaging about medium size. *Form* roundish oblate, often somewhat elliptical, ribbed. *Stem* rather short, mod-



MUNSON

erately thick. *Cavity* medium to rather large, acuminate, narrow to medium width, rather unsymmetrical, lightly russeted. *Calyx* medium to small, closed; lobes rather narrow, acute. *Basin* shallow to very shallow, narrow, obtuse or a little abrupt, furrowed, often unsymmetrical.

Skin rather thick and tough separating readily from the flesh, smooth, greenish-yellow often somewhat blushed.

Calyx tube funnel-shape with long cylinder. *Stamens* marginal to median.

Core medium to large, axile or sometimes abaxile; cells often unsymmetrical, closed or somewhat open; core lines clasping the cylinder. *Carpels* roundish to elliptical, emarginate, tufted. *Seeds* medium size, rather short, flat, obtuse, dark brown.

Flesh tinged with yellow, moderately fine-grained, tender, moderately juicy, sweet, good to very good.

Season late September to December.

NORTHERN SWEET.

REFERENCES. 1. *Cultivator*, 6:353. 1849. 2. Battey, *Horticulturist*, 4:316. 1850. fig. 3. Jeffreys, *Id.*, 4:413. 1850. 4. Barry, 1851:285. 5. Emmons, *Nat. Hist. N. Y.*, 3:142. 1851. col. pl. No. 50 and fig. 6. *Mag. Hort.*, 20:241. 1854. 7. Elliott, 1854:149. 8. Downing, 1857:177. 9. Gregg, 1857:45. 10. HOFFY, *N. A. Pom.*, 1860. col. pl. 11. *Am. Pom. Soc. Cat.*, 1862. 12. Warder, 1867:632. 13. Leroy, 1873:502. fig. 14. Thomas, 1875:508. 15. Lyon, *Mich. Hort. Soc. Rpt.*, 1881:316. 16. Bailey, *Am. Hort.*, 1892:245. 17. Waugh, *Vt. Sta. An. Rpt.*, 14:302. 1901. 18. Budd-Hansen, 1903:138.

SYNONYMS. *Golden Sweet* (8, 10, 13). NORTHERN GOLDEN SWEET (5). *Northern Golden Sweet* (2, 7, 8, 13). *Northern Golden Sweeting* (10). NORTHERN SWEET (1, 2, 3, 4, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18). NORTH-ERN SWEETING (9).

Fruit yellow or sometimes with a crimson cheek. Flesh whitish, fine, tender, juicy, sweet, very good; season midautumn. It is a good apple for the home orchard but not desirable for commercial purposes.

Historical. Supposed to be a native of Chittenden county, Vermont. Brought to notice by Jonathan Battey, Keeseville, Clinton county, N. Y., about 1849, who stated that it had then been cultivated in the vicinity of its origin for about fifty years (2). It has probably been grown more in the Champlain valley than in any other section of the state. It is still occasionally listed by nurserymen (16) but is now seldom planted.

OGDENSBURGH.

REFERENCE. 1. Downing, 1869:291.

SYNONYMS. None.

Originated with A. B. James, Ogdensburg, N. Y. According to Elliott's description cited by Downing (1) the fruit is medium size, whitish-yellow with brownish-blush; flesh tender, very mild subacid, very good; season November and December. We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

OHIO NONPAREIL.

REFERENCES. 1. *Mag. Hort.*, 14:114. 1848. 2. Emmons, *Nat. Hist. N. Y.*, 3:77. 1851. *fig.* 3. Elliott, 1854:148. *fig.* 4. *Mag. Hort.*, 22:85, 506. 1856. 5. Hooper, 1857:15. 6. Downing, C., *Mag. Hort.*, 27:59. 1861. 7. *Am. Pom. Soc. Cat.*, 1862. 8. Elliott, *Mag. Hort.*, 32:51. 1866. 9. Warder, 1867:447. *fig.* 10. Downing, 1869:29. *fig.* 11. *Am. Pom. Soc. Cat.*, 1869. 12. Fitz, 1872:171. 13. Thomas, 1875:203. 14. Barry, 1883:340. 15. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 16. Bailey, *Am. Hort.*, 1892:245. 17. Budd-Hansen, 1903:141. *fig.*

SYNONYMS. BELLFLOWER OF THE WEST (5). *Cattell Apple* (10, of Pennsylvania 6). MYERS NONPAREIL (3, 7, 8, 13). *Myers Nonpareil* (6, 9, 10, 12, 14). *Nonpareil* (6). OHIO NONPAREIL (1, 4, 6, 9, 10, 11, 12, 14, 15, 17). *Ohio Nonpareil* (5, 13). OHIO NONPAREIL (2, 16). *Red Bellflower* of some (5). *Rusty Core* (6, 10). *Wells* (5). *Western Beauty* (9, erroneously 10).

A fine fall apple of good color and good quality for either dessert or culinary purposes. The tree is a moderate grower and appears to be hardy and moderately long-lived. It does not come into bearing very young and is not always a reliable cropper. It is regarded as a good variety for home use and some recommend it for commercial planting. Season October and November.

Historical. Originated near Massillon, Ohio. Although this is an old variety having first been described in 1848 (1) we do not find that it has been much disseminated in New York. It is more popular in the Middle West and is still offered by nurserymen in that region (16).

TREE.

Tree medium size, moderately vigorous. *Form* rather spreading, not dense. *Twigs* medium length, curved, rather stout. *Bark* olive-green with some reddish-brown, thinly streaked and mottled with gray scarf-skin. *Lenticels* scattering, conspicuous, large, usually round, becoming laterally compressed. *Buds* medium to small, obtuse, appressed, pubescent.

FRUIT.

Fruit medium to large. *Form* roundish oblate, often obscurely ribbed. *Stem* medium to rather short, rather thick. *Cavity* rather large, acute, deep, pretty symmetrical, more or less russeted, the russet sometimes outspreading over the base. *Calyx* medium or above, closed or slightly open; lobes medium in length, narrow, acute. *Basin* rather small, medium in depth, narrow to moderately wide, somewhat abrupt, rather symmetrical.

Skin pale yellow to deep yellow almost entirely overspread with bright red, mottled and irregularly striped and splashed with carmine. *Dots* moderately numerous, small to medium, areolar with russet center or grayish.

Calyx tube moderately short, conical. *Stamens* basal to nearly median.

Core small, somewhat abaxile; cells symmetrical, not uniformly developed, nearly closed to somewhat open; core lines meeting or slightly clasping. *Carpels* roundish, rather flat, tufted. *Seeds* medium size, moderately long, plump, acute, tufted.

Flesh tinged with yellow, firm, moderately fine, crisp, tender, juicy, agreeable subacid, aromatic, good to very good.

Season October and November.

OHIO PIPPIN.

REFERENCES. 1. Warder, 1867:484. fig. 2. Downing, 1869:292. 3. *Am. Pom. Soc. Cat.*, 1881:12. 4. Thomas, 1885:223. 5. Taylor, *Am. Pom. Soc. Rpt.*, 1895:193. 6. Budd-Hansen, 1903:141. 7. Beach and Clark, *N. Y. Sta. Bul.*, 248:135. 1904.

SYNONYMS. *Buchanan* (1). *Ernst's Apple* (1). ERNST'S PIPPIN (4). *Ernst's Pippin* (2). OHIO PIPPIN (1, 2, 3, 5, 6, 7). *Ohio Pippin* (4). *Shannon* (1, 2, 3, 6).

Fruit of good medium size, quite attractive for a yellow apple, mild in flavor and of good quality. Season late September or October to January; October appears to be its commercial limit in this latitude (7). The tree attains good size, is rather vigorous and healthy, comes into bearing rather young and is a reliable cropper, being almost an annual bearer and often yielding full crops.

Historical. This variety is supposed to have originated in Dayton, Ohio (1, 2). It has been disseminated pretty widely in the Middle West but it is scarcely known among New York fruit growers.

TREE.

Tree rather vigorous. *Form* open, somewhat roundish or spreading and inclined to droop. *Twigs* short, straight, stout; internodes short. *Bark* dark dull brown, lightly streaked with scarf-skin, slightly pubescent near tips. *Lenticels* scattering, medium to below, oblong, not raised. *Buds* very deeply set in the bark, medium size, flat, obtuse, appressed, slightly pubescent.

FRUIT.

Fruit above medium to medium. *Form* oblate conic to roundish oblate, rather irregular in shape being often somewhat elliptical or obscurely ribbed. *Stem* short, slender, usually not exerted. *Cavity* rather large, acute or slightly acuminate, varying sometimes to rather obtuse, deep, broad, often furrowed, sometimes lipped, sometimes russeted and with outspreading russet rays. *Calyx* medium to large, open; lobes reflexed, rather broad, obtuse, separated at the base. *Basin* small and shallow to rather broad, deep and abrupt, sometimes compressed, wrinkled.

Skin smooth, somewhat glossy, attractive bright yellow often with a faint orange or pinkish blush. *Dots* whitish, submerged, sometimes russet or areolar with russet point.

Calyx tube short, rather wide above, cone-shape or approaching truncate funnel-form. *Stamens* basal or nearly so.

Core small, usually axile; cells symmetrical, closed or partly open; core lines clasping. *Carpels* broadly roundish, approaching elliptical, but slightly emarginate if at all, mucronate, slightly tufted. *Seeds* numerous, light colored, rather small, very plump, obtuse.

Flesh whitish or tinged with yellow, firm, fine, tender, crisp, moderately juicy, mild subacid becoming mildly sweet, good.

OKABENA.

REFERENCES. 1. *Am. Pom. Soc. Rpt.*, 1887:132. 2. *Ib.*, *Cat.*, 1899:19. 3. Macoun, *Can. Dept. Agr. Rpt.*, 1901:97. 4. Hansen, *S. D. Sta. Bul.*, 76:78. 1902. fig. 5. Dickens and Greene, *Kan. Sta. Bul.*, 106:54. 1902. 6. Munson, *Me. Sta. An. Rpt.*, 18:84. 1902. 7. Budd-Hansen, 1903:141. fig.

SYNONYMS. OKABENA (1, 2, 3, 4, 5, 7). OKOBENA (6).

An autumn apple not particularly attractive in appearance, nor more than moderately good in quality. Not recommended for planting in New York.

Historical. Originated in 1871 near Worthington, Minn., from seed of Oldenburg said to be fertilized by Wealthy (7). Received for testing at this Station in 1892, from the Jewel Nursery Company which introduced this variety. In 1899 it was given a place on the list of the American Pomological Society as a variety of value in the Upper Mississippi valley (2). So far as we can learn it has been grown in the East only in an experimental way.

OLDENBURG.

REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 341. 2. Kenrick, 1832:64. 3. Manning, 1838:52. 4. Ives, *Mag. Hort.*, 6:125. 1840. 5. Manning, *Ib.*, 7:44. 1841. 6. Downing, 1845:82. 7. Cole, 1849:102. 8. Thomas, 1849:147. 9. Hovey, *Mag. Hort.*, 16:495. 1850. fig. 10. Emmons, *Nat. Hist. N. Y.*, 3:34. 1851. 11. Barry, 1851:283. 12. Waring, 1851:28. 13. Elliott, 1854:131. 14. Hooper, 1857:30. 15. Gregg, 1857:42. 16. *Am. Pom. Soc. Cat.*, 1862. 17. Barry, *Horticulturist*, 22:148. 1867. 18. Warder, 1867:431. 19. Todd, 1871:186. fig. 20. Leroy, 1873:148. fig. 21. *Montreal Hort. Soc. Rpt.*, 1876:6. 22. *Ib.*, 6:97. 1880. 23. *Am. Pom. Soc. Cat.*, 1883:12. 24. Hogg, 1884:64. 25. *Can. Hort.* 11:221. 1888. 26. Hoskins, *Rural N. Y.*, 47:646. 1888. 27. Dunlap, *Ill. Hort. Soc. Rpt.*, 1889:23. 28. *Can. Hort.*, 12:75. 110. 1889. 29. *Montreal Hort. Soc. Rpt.*, 15:26. 1890. 30. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 31. Bailey, *Am. Hort.*, 1892:237. 32. *Am. Gard.*, 14:519. 1893. 33. *Can. Hort.*, 17:291. 1894. 34. *Rural N. Y.*, 53:28. 1894. 35. *Am. Gard.*, 17:519. 1896. 36. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:354. 37. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 6:8. 1899. figs. 38. Craig, *Cyc. of Hort.*, 1901:1404. 39. Van Deman, *Rural N. Y.*, 60:248. 1901. 40. Alwood, *Va. Sta. Bul.*, 130:121. 1901. 41. Waugh, *Vt.*



OLDENBURG - Reduced Size

Sta. An. Rpt., 14:302. 1901. 42. Hansen, *S. D. Sta. Bul.*, 76:46. 1902. 43. *Ib.*, 76:79. 1902. fig. 44. Dickens and Greene, *Kan. Sta. Bul.*, 106:52. 1902. 45. Bruner, *N. C. Sta. Bul.*, 182:21. 1903. 46. Farrand, *Mich. Sta. Bul.*, 205:45. 1903. 47. Budd-Hansen, 1903:141. fig. 48. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:51. 1903. 49. Beach and Clark, *N. Y. Sta. Bul.*, 248:136. 1904.

SYNONYMS. *Barowski* (20). *Barowski* (20). *Borovitsky* (20, 37, 38). *BOROWICKI* (20). *Borowitski* (24). *Borowitsky* (43). *Charlamowski* (24). *Charlamowski d'Automne* (20). *Charlamowskircher Nalleoid* (20). *Charlamowsky* (43). *DUCHESS* (27, 29, 32, 35, 37, 38, 43). *DUCHESS OF OLDENBURG* (3, 4, 11, 12, 16, 17, 19, 21, 22, 24, 25, 28, 31, 33, 36, 44). *Duchess of Oldenburg* (23, 41, 48, 49). *DUCHESS OF OLDENBURGH* (1, 2, 5, 9, 10, 14, 18). *Duchesse d'Oldenbourg* (20). *DUTCHESS* (13). *DUTCHESS OF OLDENBERG* (15). *DUTCHESS OF OLDENBURGH* (6, 7, 8). *Dutchess of Oldenburgh* (13). *New Brunswick* (34). *OLDENBURG* (23, 26, 39, 45, 48, 49). *Oldenburg* (35, 37). *OLDENBURGH* (30, 34). *OLDENBURG, Duchess of* (40, 42, 46, 47).

This Russian apple is known throughout the West either by the name *Duchess*, or by the full name *Duchess of Oldenburg*; the American Pomological Society has abbreviated the full name to *Oldenburg*, but this has not been generally accepted by Western fruit growers. In European nurseries it is propagated under the names of *Charlamowsky* and *Borowitsky*. It was early imported into the West, coming to this country by the way of England and it was the extreme hardiness of this variety in the early test winters that kept up the hopes of prairie orchardists in time of great discouragement and led to the importations of more varieties from Russia (42).

Oldenburg is one of the most valuable of the Russian apples thus far introduced into this country. It is of good size and attractive appearance. It is generally highly esteemed for home use on account of its excellent culinary qualities and with some fruit growers it has proved a very profitable variety for the commercial orchard. When properly grown and carefully handled it stands shipment pretty well and sells well for a variety of its season. In some few localities in Western New York it is grown in sufficient quantities so that it can be shipped in car lots to distant markets, but in very many places it is produced in greater quantities than the local markets can absorb and yet not in quantities large enough so that it can be economically shipped to distant markets. Since the fruit is quite perishable it does not stand heat well before shipment

and goes down rather quickly, particularly if the weather is unfavorable. When sent to distant markets it should be shipped under ice. The fruit ripens in succession so that several pickings are required in order to secure the crop in prime marketable condition. It is in season during late August and September, but it may be used for culinary purposes before it is fully ripe. The tree is highly valued because of its great hardiness. It is vigorous when young but with age it becomes a rather moderate or slow grower. It is only moderately long-lived, but is a reliable cropper, yielding good crops biennially, often with lighter crops alternating. Generally speaking, the trees require good cultivation, thorough fertilizing and careful spraying in order to secure the best commercial results. The fruit hangs fairly well to the tree till it is ripe. It is quite uniform in size and quality, with but a small percentage of loss from unmarketable fruit.

Historical. This is one of the four pioneers among Russian apples in America, the other three being Alexander, Tetofsky and Red Astrachan (38). These four varieties were imported by the Massachusetts Horticultural Society from the London, England, Horticultural Society about 1835. Oldenburg was brought to England from Russia about twenty years prior to that date. It was tested by Robert Manning, Superintendent of the Test Garden of the Massachusetts Horticultural Society at Salem who published the following description in 1838 (3): "A valuable and handsome apple said to be of Russian origin. The size is middling, form round and rather flat; skin of a beautiful yellow, striped with red; flavor very pleasant and good. It bears well and ripens in September and October." In 1850 Hovey wrote, "Mr. Manning, we believe, first proved the Duchess of Oldenburg and gave a brief account of it in his Book of Fruits. Since then it has been considerably disseminated, and though yet far from being common is to be found in many fine collections of fruit" (9). Later it became disseminated throughout the Middle West and Northwest where it proved to be much superior in hardiness to Baldwin, Rhode Island Greening, Northern Spy and other varieties which have been commonly cultivated in this state. Its ability to withstand severe climates encouraged the importation of other Russian sorts some of which have proved valuable in the northern portion of the apple belt. Oldenburg is commonly listed by nurserymen (31) and its planting both in home orchards and in commercial orchards is increasing in this state.

TREE.

Tree medium in size. *Form* at first upright spreading but eventually roundish. *Twigs* moderately long, curved, slender; internodes long. *Bark* dark brown, lightly mottled with scarf-skin; slightly pubescent. *Lenticels*

scattering, small to medium, oblong, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium to large, averaging above medium, uniform in size and shape. *Form* roundish oblate to oblate, regular, symmetrical. *Stem* short to sometimes medium, moderately slender. *Cavity* acute to acuminate, deep, broad, usually partly covered with greenish-russet. *Calyx* medium to rather large, usually closed; lobes rather broad, acute. *Basin* moderately deep to deep, wide, abrupt, smooth or with small mammiform protuberances.

Skin moderately thick, tender, smooth, pale greenish-yellow or pale yellow, almost covered with irregular splashes and stripes of bright red mottled and shaded with crimson. *Dots* scattering, small, light colored. *Prevailing effect* red striped; attractive.

Calyx tube moderately long, rather wide, funnel-shape with broad truncate cylinder or approaching urn-shape. *Stamens* median.

Core medium to rather large, axile; cells symmetrical, closed or slightly open; core lines clasping. *Carpels* broadly ovate, slightly emarginate. *Seeds* medium to rather large, wide, obtuse to acute, moderately plump, dark brown.

Flesh tinged with yellow, rather firm, moderately fine, crisp, tender, juicy, sprightly subacid, aromatic, good to very good for culinary purposes. It has too much acidity for a good dessert apple.

Season late August and September.

ORANGE.

REFERENCES. 1. M'Mahon, *Am. Gard. Cal.*, 1806:585. 2. Cox, 1817:139. 3. Emmons, *Nat. Hist. N. Y.*, 3:91. 1851. 4. *Horticulturist*, 8:247. 1853. 5. *Mag. Hort.*, 19:172. 1853. 6. Hooper, 1857:67. 7. Downing, 1857:178. 8. Warder, 1867:728. 9. Downing, 1869:294. 10. Bailey, *Am. Hort.*, 1892:245.

SYNONYMS. None.

Different varieties of the apple have been known under the name Orange. The Fall Orange has already been described on page 60.

Orange has sometimes been used as a synonym for Lowell which is described on page 128.

ORANGE OF NEW JERSEY (1, 2, 3, 6, 7, 8, 9, 10). Cox (2) gives the following description of this variety: "This is a fine table apple in the fall and early winter months; and is thought to be a good cider fruit; the size is small, the form oblong—the colour a greenish-yellow—the flesh yellow, rich, juicy and sprightly; the tree is of moderate size, the growth upright, and its fruitfulness great. It is much cultivated in several of the middle counties of New-Jersey as a highly estimable apple." Downing (7, 9) states that the tree is vigorous and moderately productive; the fruit pleasant subacid; very good; season September and October.

ORANGE OF PENNSYLVANIA. A variety which originated at Reading, Pa., was brought to notice under the name Orange by the *ad interim* report of the Fruit Committee of the Pennsylvania Horticultural Society in 1853 (4, 5). The fruit is described as medium size, roundish, nearly orange-yellow, sprightly,

good (4, 5, 8, 9). We have received no reports concerning this variety from any of our correspondents and so far as we know it is not in cultivation in New York.

ORANGE PIPPIN.

REFERENCES. 1. Forsyth, 1803:54. 2. *Ib.*, 1824:119. 3. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 4. *London Hort. Soc. Cat.*, 1831:No. 587. 5. Kenrick, 1832:81. 6. Lindley, 1833:80. 7. Warder, 1867:728. 8. *Am. Pom. Soc. Cat.*, 1869. 9. Downing, 1872:62 app. fig. 10. Leroy, 1873:457. fig. 11. *Am. Pom. Soc. Cat.*, 1873. 12. Hogg, 1884:116. 13. *Ib.*, 1884:164. 14. Thomas, 1885:519. 15. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 16. Bailey, *Am. Hort.*, 1892:245.

SYNONYMS. *Englese Orange Appel* (12). *ISLE OF WIGHT PIPPIN* (4, 12). *Isle of Wight Pippin* (5, 6). *Isle of Wight Orange* (4, 5, 6, 12). *MARIGOLD* (10). *Marigold* (5). *Marigold Creed's* (10). *Marigold Pippin* (2). *Marygold* (6). *ORANGE* (7). *ORANGE PIPPIN* (1, 2, 3, 5, 6, 8, 9, 11, 13, 14, 15, 16). *Orange Pippin* (4, 12). *Pomme d'Orange* (12).

The above references are not all to the same variety and are included only because in many cases it is uncertain which variety the writer had in mind.

At least two varieties have been known in this country under the name Orange Pippin but so far as we can learn neither of them is now considered of superior value by fruit growers and both are going out of cultivation. One, which has been called also the Isle of Wight Pippin, Isle of Wight Orange, Marygold and Marigold, was disseminated from the Isle of Wight where, as some have supposed, it was brought from Normandy (1, 3, 4, 5, 6, 10, 12). Fruit medium size, roundish, skin yellowish, golden gray russeted and highly colored with orange and red on the sunny side; flesh firm, crisp, pleasant acid, suitable for dessert; season October to January (6, 9). This was entered on the list of the American Pomological Society in 1869, dropped in 1871 and re-entered in 1873.

An Orange Pippin grown in some parts of New Jersey is described (8, 9, 11, 14, 15) as a profitable summer market apple of unknown origin. Tree vigorous, at first upright but eventually spreading; a reliable biennial cropper. Fruit medium or above, pale yellow with some orange red in the sun; flesh white, half fine, tender, pleasant subacid; season September and October.

ORANGE SWEET.

REFERENCES. 1. *Mag. Hort.*, 1:396. 1835. 2. Warder, 1867:566. 3. Downing, 1869:295. 4. Thomas, 1885:519.

SYNONYMS. *Orange Russet* (2). *ORANGE SWEET* (3, 4). *ORANGE SWEETING* (1, 2).

Several varieties are described under this name by Downing (3): one from Ohio, large, greenish-yellow; flesh whitish, tender, sweet, good; season October and November; one from Massachusetts, the fruit medium, oblate, greenish-yellow; the flesh yellowish-white, rather coarse, rich, sweet; season August and September; and one from Maine; fruit medium, roundish ovate,

bright yellow with blush; flesh yellowish, tender, sweet, rich; season September and October.

Warder (2) describes under the name Orange Sweeting or Russet "An eastern variety not much cultivated; fruit large, very round, regular, greenish-yellow, bronzy, orange russeted; flesh rather tough, fine-grained, juicy, good; season December."

The name Orange Sweet has also been used as a synonym for both Munson (p. 146) and Golden Sweet (p. 81).

OSTRAKOFF.

REFERENCES. 1. Gibb, *Montreal Hort. Soc. Rpt.*, 1883:106. fig. 2. *Ib.*, *Ia. Hort. Soc. Rpt.*, 1883:443. 3. Budd, *Ia. Agr. Coll. Bul.*, 1885:18. 4. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:71. 5. Craig, *Ib.*, 1886-87:103. 6. Budd, *Rural N. Y.*, 47:692. 1888. 7. *Ib.*, *Ia. Agr. Coll. Bull.*, 1890:23. 8. Bailey, *An. Hort.*, 1892:246. 9. Budd, *Ia. Sta. Bul.*, 19:540. 1892. 10. Taylor, *Me. Pom. Soc. Rpt.*, 1892:58. 11. *Can. Hort.*, 16:402. 1893. 12. Beach, *N. Y. Sta. An. Rpt.*, 13:583. 1894. 13. Munson, *Me. Sta. Rpt.*, 1896:75. 14. Thomas, 1897:648. 15. Munson, *Me. Sta. Rpt.*, 1902:84. 16. Hansen, *S. D. Sta. Bul.*, 76:80. 1902. fig. 17. Beach and Clark, *N. Y. Sta. Bul.*, 248:136. 1904.

SYNONYMS. *Astravaskoe* (12). OSTRAKOFF (7, 9, 10, 12, 13, 15, 16, 17). OSTRAKOFF GLASS (11). *Ostrakoff Glass* (12). OSTREKOFF (3, 5). *Ostre-koff's Glass* (1, 2). *Ostrekovskaya Steklanka* (3). OSTREKOWSKAJA STEK-LIANKA (1, 2). OSTROKOFF (6, 8, 14). OSTROKOFF'S GLASS (4). *No. 4 M* (4, 5, 6, 7, 9, 11, 13, 15, 16). *No. 472* (1, 3).

A Russian variety of good size, greenish-yellow, brisk subacid, fair to good quality. Its keeping qualities vary much in different seasons. As grown at this Station it is commonly in its prime from late September into November but a portion of the fruit may often be kept into the winter in very good condition. It is reported as a promising variety for portions of Northern New England and other regions where superior hardiness is a prime requisite. It is of no value where our common standard varieties succeed.

Historical. Described by Budd in 1885 under the name Ostrakoff's Glass and in 1890 under the name Ostrakoff (3, 7). It was received in 1884 for testing at this Station from Ellwanger and Barry, Rochester, N. Y., under the name Astravaskoe.

TREE.

Tree moderately vigorous. *Form* spreading or roundish, open. *Twigs* short, curved, stout with large terminal buds; internodes medium. *Bark* dark brown tinged with green, heavily streaked with scarf-skin; pubescent near tips. *Lenticels* quite numerous, medium size, round, raised, rather conspicuous. *Buds* prominent, large, long, plump, obtuse, free.

FRUIT.

Fruit medium or above, pretty uniform in size and shape. *Form* roundish to somewhat ovate or oblong, somewhat conical, a little angular. *Stem*

characteristically long, often curved, twisted or irregularly enlarged or inserted under a fleshy protuberance. *Cavity* acute or approaching acuminate, shallow to moderately deep, medium in width to narrow, partly covered with light greenish-russet, often lipped. *Calyx* closed or open, rather large, leafy; lobes rather broad, acute to obtuse. *Basin* characteristically irregular, medium in width and depth, abrupt, sharply ridged and wrinkled.

Skin moderately thin, rather tough, smooth, pale waxy-yellow sometimes with a faint reddish shade. *Dots* very numerous, small, submerged, often areolar.

Calyx tube medium to large, rather wide, conical to peculiarly funnel-form, with broad truncate cylinder. *Stamens* basal to median.

Core medium size, axile; cells closed or slightly open; core lines meeting. *Carpels* roundish to broadly ovate, emarginate. *Seeds* medium brown, large, wide, rather flat, obtuse to acute.

Flesh yellowish-white, firm, rather fine, juicy, brisk subacid, fair to good.

Season late fall and early winter.

PALOUSE.

REFERENCES. 1. Van Deman, *U. S. Pom. Rpt.*, 1891:390. 2. *Rural N. Y.*, 50:815. 1891. 3. Hexamer, *Am. Pom. Soc. Rpt.*, 1891:159. 4. Williams, *Gard. and For.*, 5:11. 1892. 5. Bailey, *Am. Hort.*, 1892:246. 6. Craig, *Can. Dept. Agr. Rpt.*, 1896:133. 7. Macoun, *ib.*, 1901:97.

SYNONYMS. None.


Palouse is a comparatively new variety which originated in Washington. In color, texture, flavor and aroma it is so much like Tompkins King that some suppose it to be a seedling of that variety. Its form, however, is quite different from that of Tompkins King, typical fruit being oblong, as described by Craig (6) and others (1, 2), rather than oblate or roundish oblate, as described by Hexamer (3) and Macoun (7). It is hardly as good a keeper as Tompkins King and would probably be classed as a late fall variety, being in season from October to early winter. It is, however, more productive than Tompkins King and is being much planted in commercial orchards in Washington where many regard it as one of the most valuable of the seedling varieties which have originated in that region. It appears to be worthy of testing in New York.

Historical. Palouse originated in Whitman county, Washington, from seed brought from Illinois in 1879. The original tree is located five miles east of Colfax¹. It was introduced about 1892 by George Ruedy, Colfax, Wash., whose

¹ Ruedy, Letter and Circular, 1904.



PARRY WHITE



attention was first called to the variety in 1889 when it took first premium as the best seedling apple exhibited at the Whitman County Fair. It is as yet but little grown in the East and so far as we can learn it has as yet been planted in New York to but a limited extent and for trial only.

FRUIT (2, 3).

Fruit large. *Form* oblong conic, ribbed or scalloped. *Stem* long, slender. *Cavity* remarkably deep. *Calyx* closed or partly open. *Basin* rather shallow to deep, distinctly furrowed, the furrows often extending to the cavity.

Skin bright yellow, blushed and more than half covered with crimson, splashed, blotched and dotted with darker red. *Prevailing effect* attractive red and yellow.

Core large.

Flesh yellowish, crisp, firm, tender, juicy, very aromatic, subacid, very good.

Season October and early winter.

PARRY WHITE.

REFERENCES. 1. Downing, 1872:25 app. 2. Bailey, *An. Hort.*, 1892:246. 3. Beach, *N. Y. Sta. An. Rpt.*, 14:263. 1895. 4. Beach and Clark, *N. Y. Sta. Bul.*, 248:137. 1904.

SYNONYMS. *Imperial White* (1). PARRY WHITE (1, 2, 3, 4). *Superior White* (1). *White Apple* (1).

Fruit pale yellow or whitish, waxen, of medium size; quite attractive in appearance for an apple of its class. The flesh is white, juicy, subacid, good in quality for either dessert or culinary use. It follows Yellow Transparent in season, beginning to ripen late in August or early in September and continuing in use into or through October (4). The tree comes into bearing young and is an annual cropper, yielding good to very heavy crops. In spite of its remarkable productiveness it makes a pretty good growth. It is worthy of trial where an apple of its type is desired.

Historical. Origin unknown. It is supposed to be a Pennsylvania apple. It is occasionally listed by nurserymen (2). It has not yet become known to any considerable extent in New York.

TREE.

Tree of medium size and moderately vigorous. *Form* at first upright spreading but after bearing full crops it becomes roundish and somewhat drooping; dense. *Twigs* short, curved, stout with large terminal buds; internodes short. *Bark* brown tinged with olive-green, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous.

rather conspicuous, medium size, round, not raised. *Buds* medium to large, plump, obtuse, free, pubescent.

FRUIT.

Fruit usually medium but varies according to the abundance of the crop from below medium to rather large; quite uniform in size and shape. *Form* nearly globular varying to somewhat oblate or slightly inclined to ovate, quite regular. *Stem* above medium to rather short, slender. *Cavity* slightly acuminate, medium to shallow, medium in width, smooth, symmetrical. *Calyx* small, closed; lobes medium in length, narrow, acute. *Basin* very shallow, moderately narrow to rather broad, obtuse, smooth or slightly wrinkled, symmetrical.

Skin moderately thick, rather tough, waxen yellowish-white or greenish. Occasionally blushed. *Dots* medium size to very small, pale or brown, numerous, depressed.

Calyx tube short, narrow, funnel-shape. *Stamens* marginal to median.

Core medium to small, abaxile; cells open; core lines meeting or slightly clasping. *Carpels* round, emarginate. *Seeds* medium or above, acute or inclined to obtuse, light brown.

Flesh white, quite firm, rather fine, tender, juicy, subacid, good.


Season very late August into or through October.

PATTEN.

REFERENCES. 1. Watrous, *Am. Pom. Soc. Rpt.*, 1889:124. 2. Patten, C. G., *Descriptive Circular*, 1891. 3. Taylor, *Me. Pom. Soc. Rpt.*, 1892:60. 4. *Me. Sta. Rpt.*, 1892:90. 5. Macoun, *Can. Hort.*, 22:396. 1899. 6. *Am. Pom. Soc. Cat.*, 1899:19. 7. Macoun, *Can. Dept. Agr. Rpt.*, 1901:98. 8. Munson, *Me. Sta. An. Rpt.*, 18:84. 1902. 9. Hansen, *S. D. Sta. Bul.*, 76:81. 1902. fig. 10. Budd-Hansen, 1903:145. fig. 11. Ragan, *U. S. B. P. I. Bul.*, 56:226. 1905.

SYNONYMS. *Duchess No. 3* (2, 9, 10). PATTEN (11). *Patten's Duchess No. 3* (11). PATTEN GREENING (1, 2, 3, 4, 5, 6, 7, 8, 9). *Patten Greening* (11). PATTEN GREENING (10).

A seedling of Oldenburg which, on account of its hardness, productiveness and the uniformly large size of its fruit is valuable in the northern portions of the apple-growing regions of the country (1, 5, 6, 8, 9, 10). It is grown as far north as the Red River valley of Minnesota and North Dakota and in other regions where the winters are correspondingly severe. It is attractive in color for a green apple, has a sprightly subacid flavor and good texture and is very good in quality for culinary use. The tree is a somewhat stronger grower than Oldenburg, with limbs strongly shouldered (2, 9, 10). As grown at this Station it comes into bearing moderately young and is an annual cropper, yielding moderate to full crops. It is worthy of trial in the colder regions of the state.



PATTEN

Historical. Originated by C. G. Patten, Charles City, Ia. (2), who states that it is a seedling of the Oldenburg from seed grown near Portage, Wis., and planted by him at Charles City, Ia., in the fall of 1866. It was first illustrated and described in the catalogue of the originator in the autumn of 1885 when stock of this variety was first offered for sale. In 1899 it was entered in the catalogue of the American Pomological Society as a valuable variety for the Upper Mississippi valley (6). It has as yet been but little disseminated in New York.

TREE.

Tree moderately vigorous with moderately long, somewhat stout, curved, drooping branches. *Form* spreading, dense, flat. *Twigs* rather short, somewhat curved, moderately stout or rather slender; internodes medium to long. *Bark* brownish-red with some olive-green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, small, roundish, sometimes raised. *Buds* medium size, plump, acute, free, slightly pubescent.

FRUIT.

Fruit medium to large, pretty uniform in size and shape. *Form* oblate or roundish oblate, sometimes inclined to conic, usually regular, pretty symmetrical. *Stem* very short to medium in length, moderately thick. *Cavity* acuminate, deep or sometimes medium in depth, rather broad or compressed, russeted and with outspreading russet rays. *Calyx* large or very large, closed or somewhat open. *Basin* deep to medium in depth, medium in width, often abrupt, usually smooth and symmetrical.

Skin moderately thin, tough, smooth, clear pale greenish-yellow, sometimes blushed and occasionally faintly striped. *Dots* small, numerous, pale and submerged or brownish.

Calyx tube conical to funnel-shape. *Stamens* median to basal.

Core below medium to small, axile or somewhat abaxile; cells sometimes unsymmetrical, closed or partly open; core lines clasping. *Carpels* roundish, irregular. *Seeds* dark brown, medium or above, plump, obtuse.

Flesh tinged with yellow, moderately firm, a little coarse, rather tender, juicy, sprightly subacid, good in quality especially for culinary purposes.

Season October to January.

PEACH (MONTREAL).

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 1876:7. 2. *Raynes, Ib.*, 1879:26. 3. *Budd, Ia. Hort. Soc. Rpt.*, 1880:524. 4. *Lyon, Mich. Hort. Soc. Rpt.*, 1881:178. 5. *Am. Pom. Soc. Cat.*, 1883:12. 6. *Thomas*, 1885:520. 7. *Montreal Hort. Soc. Rpt.*, 1885:99. 8. *Ib.*, 1886-87:92. 9. *Burnet, Can. Hort.*, 12:339. 1889. 10. *Hamilton, Montreal Hort. Soc. Rpt.*, 15:26. 1890. 11. *Bailey, An. Hort.*, 1892:246. 12. *Munson, Me. Sta. Rpt.*, 1893:133. 13. *Hoskins, Am. Gard.*, 15:272. 1894. 14. *Hansen, S. D. Sta. Bul.*, 76:82. 1902. 15. *Budd-Hansen*, 1903:147.

SYNONYMS. *Canada Peach* (14). *Montreal Peach* (3, 14). *PEACH* (3, 11, 14). *PEACH APPLE of Montreal* (13). *PEACH OF MONTREAL* (1, 4, 5, 6, 7, 8, 9). *PEACH of Montreal* (2, 10, 12, 15). *Pomme Peche* (1).

Fruit predominantly yellow with some red; in form it somewhat resembles Porter being oval or conical. It is salable in local markets but being easily bruised it does not stand transportation well (2, 3, 6, 13). The tree is hardy, thrifty and very productive.

According to Hoskins it is a very popular fall apple all through Northern New England and Eastern Canada. It has long been the leading market apple of its season, September, in Montreal and the surrounding territory on both sides of the line. Its season does not extend far into October except by extra care but it forms a very good successor to the Yellow Transparent which it resembles except for its blushed cheek (13).

This variety is but little known in New York. It cannot be expected to displace Oldenburg which is of the same season, more productive (10) and much better known.

Historical. Some writers state that this is a variety of French origin (1) but Hoskins considers this doubtful. He states: "Some of its characters would indicate it to be a Russian apple which has reached Canada via France, but this is only a conjecture" (13).

PEACH POND.

REFERENCES. 1. Downing, 1845:91. 2. Thomas, 1849:145. *fig.* 3. Emmons, *Nat. Hist. N. Y.*, 3:36. 1851. 4. Elliott, 1854:151. *fig.* 5. Hooper, 1857:68. 6. Warder, 1867:476. *fig.* 7. *Am. Pom. Soc. Cat.*, 1871:8. 8. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 9. Bailey, *Am. Hort.*, 1892:246. 10. Burrill and McCluer, *Ill. Sta. Bul.*, 45:335. 1896.

SYNONYMS. PEACH POND (3, 4, 5, 6, 7, 8, 9). PEACH-POND SWEET (1, 2). PEACH POUND SWEET (10).

A beautiful autumn sweet apple of excellent quality in season from September to November. The tree is vigorous and spreading. It originated in Dutchess county, N. Y. (1). It was entered in the Catalogue of the American Pomological Society in 1871 and dropped in 1899. Although it has long been in cultivation it has failed to establish itself as a commercial variety in New York and is comparatively little grown for home use.

FRUIT (1, 6).

Fruit medium to small. *Form* slightly conic, rather flat, angular and a little one-sided. *Stem* medium to long, slender, green, sometimes knobbed. *Cavity* acute, deep, regular, brown. *Calyx* small, closed. *Basin* narrow, regular, wrinkled.

Skin smooth, pale yellow lightly covered with mixed striped red and beautifully splashed with crimson.

Flesh yellowish, very mellow, fine-grained, moderately juicy, rich, sweet, agreeable, very good or almost best.

Season September to November.

PEARSALL.

REFERENCES. 1. Downing, 1869:300. 2. Thomas, 1875:508.

SYNONYMS. PEARSALL'S SWEET (1, 2).

This variety is supposed to have originated in Queens county, N. Y. According to Downing the fruit is a good keeper and valuable for baking. The tree upright spreading, quite productive. Fruit rather large, yellow, partly covered with light red. Flesh moderately juicy, sweet, good. Season November to January (1).

PEASE.

REFERENCES. 1. *Rural N. Y.*, 54:776. 1895. fig. 2. Heiges, *U. S. Pom. Rpt.*, 1895:36. 3. *Rural N. Y.*, 56:222. 1897. 4. *Am. Pom. Soc. Cat.*, 1899:19. 5. *Rural N. Y.*, 61:249. 1902. 6. Budd-Hansen, 1903:147. 7. J. W. Adams and Co., Springfield, Mass., *Cat.*

SYNONYMS. PEASE, *Waller* (4, 6). WALTER PEASE (1, 2, 3, 5, 7).

A pleasant-flavored apple of good size, attractive appearance and excellent dessert quality, but too mild in flavor to excel for culinary uses. The fruit being rather tender requires careful handling and on this account is better adapted for local than for distant markets. It is worthy of the attention of New York fruit growers where a dessert apple of this type is desired. The crop ripens unevenly. The earliest fruit comes in season the latter part of September or early in October, while a considerable portion of the later ripening fruit may remain sound till midwinter or later. Sometimes there is considerable loss from premature dropping of the fruit during September wind storms. The tree is a good grower, hardy, comes into bearing moderately young and is a pretty reliable cropper, alternating light with heavier crops.

Historical. Originated in the seedling orchard of Walter Pease, Somers, Conn., in the early part of the last century (3). It was at first propagated by the Shakers near the place of its origin and there came to be recognized locally as a valuable variety. Within recent years it has been propagated to a considerable extent by nurserymen and is being more widely disseminated.

TREE.

Tree rather large, moderately vigorous to vigorous. *Form* upright to roundish. *Twigs* moderately long, a little curved, stout; internodes medium to short. *Bark* clear brownish-red, heavily coated with scarf-skin; pubescent. *Lenticels* scattering, below medium to above medium, roundish, slightly

raised. *Buds* below medium to above, broad, plump, rather obtuse, free or nearly so, slightly pubescent.

FRUIT.

Fruit large or above medium, somewhat variable in size and shape. *Form* flattened at base, varying from oblate to roundish oblong and often inclined to conic; sides often unequal. *Stem* below medium to long, slender. *Cavity* obtuse to acute, moderately deep to deep, broad, sometimes lipped, often somewhat russeted. *Calyx* usually medium to large, closed or nearly so; lobes leafy. *Basin* moderately deep or sometimes shallow, rather narrow, abrupt, often compressed or furrowed, wrinkled.

Skin thin, tough, smooth, somewhat glossy, with pale green or yellowish ground color which in highly colored specimens is largely covered with bright red, striped with bright carmine and flecked with whitish scarf-skin. *Dois* numerous, often submerged, whitish or russet, sometimes areolar. *Prevailing effect* greenish-yellow more or less striped with red.

Calyx tube small, conical.

Core small to medium, axile; cells partly open or sometimes closed; core lines meeting. *Carpels* small, roundish, somewhat tufted. *Seeds* rather large, narrow, long, somewhat tufted, dark; often some are abortive.

Flesh whitish, slightly tinged with yellow, firm, rather fine, crisp, tender, juicy, aromatic, sprightly, mild pleasant subacid, good to very good for dessert.

Season October to midwinter.

PEASGOOD NONSUCH.

REFERENCES. 1. Downing, 1881:100 app. 2. Hogg, 1884:170. 3. Bailey, *Am. Hort.*, 1892:246. 4. *Can. Hort.*, 15:346. 1892. 5. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:356.

SYNONYMS. PEASGOOD'S NONSUCH (1, 2, 5). PEASGOOD NONSUCH (3, 4).

An English culinary apple, very large, green or yellowish, blushed and striped with red, excellent for culinary use. In season during September and October (1). In England it is regarded as one of the valuable varieties introduced in the last fifty years (2, 5). It has been but little tested in this country.


PERRY REDSTREAK.

REFERENCES. 1. *Horticulturist*, 24:157. 1869. 2. Downing, 1876:63 app. 3. Ragan, *U. S. B. P. I. Bul.*, 56:230. 1905.

SYNONYMS. PERRY REDSTREAK (3). PERRY RED STREAK (1, 2). *Perry Red Streak* (3).

A November apple of medium size and mild flavor. It originated at Lowville, N. Y., with Dr. David Perry (1). The tree is hardy, vigorous and a reliable cropper alternating heavy with lighter crops. The fruit is of medium size, yellow, shaded, striped and splashed with light and dark red; flesh whitish, fine, tender, juicy, slightly aromatic, mild subacid (2).

This variety is unknown to us and we have received no report concerning it from any of our correspondents.



PEASE *WALTER*

PETER.

REFERENCES. 1. *Iowa Hort. Soc. Rpt.*, 1886:180. 2. *N. Y. Sta. An. Rpt.*, 7:50, 90. 1888. 3. *Bailey, An. Hort.*, 1892:246. 4. *Beach, N. Y. Sta. An. Rpt.*, 12:601. 1893. 5. *Heiges, U. S. Pom. Rpt.*, 1895:31. 6. *Munson, Me. Sta. Rpt.*, 1896:75. 7. *Beach, W. N. Y. Hort. Soc. Rpt.*, 41:52. 1896. 8. *Am. Pom. Soc. Cat.*, 1899:19. 9. *Hansen, S. D. Sta. Bul.*, 76:84. 1902. 10. *Munson, Me. Sta. An. Rpt.*, 18:84, 95. 1902. 11. *Farrand, Mich. Sta. Bul.*, 205:46. 1903. 12. *Budd-Hansen, 1903:149.* 13. *Powell and Fulton, U. S. B. P. I. Bul.*, 48:52. 1903. 14. *Beach and Clark, N. Y. Sta. Bul.*, 248:137. 1904.

SYNONYMS. None.

This fruit resembles its parent Wealthy so closely that it is very difficult to distinguish between the two. It is similar to Wealthy in season as well as in the appearance and quality of the fruit. In ordinary storage its commercial season is September and October (14). In cold storage it extends to about January 1 (13). Sometimes a portion of the fruit may keep into or through the winter (14). Hansen states that even if really distinct the two varieties are now mixed to a considerable extent. As fruited at this Station the flesh of Peter is not so white as that of Wealthy but somewhat milder and better in quality (4). It has been found that the seeds of Peter are larger, broader, less pointed and a little darker than those of Wealthy. The tree is a moderate grower, comes into bearing young and yields full crops biennially. The fruit does not ripen uniformly and on this account there is apt to be some loss from drops unless more than one picking is made.

Historical. Originated from seed of Wealthy by Peter M. Gideon, Excelsior, Minn. (1), from whom this variety was received in 1888 for testing at this Station.

TREE.

Tree a fairly strong grower in the nursery; in the orchard it makes a moderately vigorous growth and eventually becomes rather large with moderately stout, somewhat drooping branches. *Form* upright spreading to roundish, open. *Twigs* short, straight, slender, with large terminal buds; internodes medium. *Bark* brown tinged with red, lightly streaked with scarf-skin; pubescent near tips. *Lenticels* quite numerous, medium size, oval, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium or above, uniform in size and shape. *Form* oblate or roundish oblate, a little inclined to conic, regular. *Stem* long to medium, sometimes short, slender. *Cavity* acuminate, deep, medium to rather broad, compressed, lightly russeted or nearly smooth. *Calyx* small, closed; lobes medium in length, rather broad, acute. *Basin* deep to moderately deep, narrow to medium in width, abrupt, gently furrowed, sometimes compressed.

Skin thin, moderately tough, nearly smooth, clear pale yellow washed and mottled with bright red conspicuously striped and splashed with deep carmine. Highly colored specimens are nearly covered with deep red. *Dots*

medium size, scattering, brown, mingled with some that are whitish and submerged. *Prevailing effect* red or striped red.

Calyx tube small, funnel-shape. *Stamens* median to marginal.

Core medium to below, usually axile; cells symmetrical, closed or partly open; core lines clasping. *Carpels* roundish, emarginate. *Seeds* above medium to large, moderately wide, long, flat, moderately acute.

Flesh slightly tinged with yellow, sometimes stained with red, firm, medium-grained, tender, juicy, with a pleasant, mild subacid, somewhat aromatic flavor, good to very good.

Season September and October or later (14).

PLUMB CIDER.

REFERENCES. 1. *Wis. Hort. Soc. Rpt.*, 1869 (cited by 2). 2. *Horticulturist*, 27:310. 1872. 3. *Id.*, 28:119. 1873. 4. *Am. Pom. Soc. Cat.*, 1873. 5. Williams, *Horticulturist*, 29:16. 1874. fig. 6. Downing, 1876:65 app. 7. Budd, *Id. Hort. Soc. Rpt.*, 1879:472. 8. Barry, 1883:352. 9. Thomas, 1885:520. 10. Bailey, *Am. Hort.*, 1892:246. 11. *Rural N. Y.*, 53:794. 1894. 12. Alwood, *Va. Sta. Bul.*, 130:124. 1901. 13. Hansen, *S. D. Sta. Bul.*, 76:35. 1902. fig. 14. Budd-Hansen, 1903:151. fig. 15. Beach and Clark, *N. Y. Sta. Bul.*, 248:138. 1904.

SYNONYMS. PLUMB CIDER (1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 14, 15). PLUM CIDER (10, 11).

This variety has been received with some favor in many parts of the country on account of its hardiness and productiveness. The fruit is of good medium size, fairly attractive in color, yellowish shaded and splashed with red. As grown in New York state it is inferior to standard varieties of its season.

Historical. Origin unknown. It was brought from Ohio in 1844 to Wisconsin by Mr. Plumb, where it proved to be a good grower, hardy and productive (5).

FRUIT (5, 13, 14).

Fruit above medium. *Form* round-ovate, slightly conic in some specimens. *Stem* stout, short. *Cavity* shallow, narrow. *Calyx* very small, closed. *Basin* very narrow and shallow, slightly plaited.

Skin yellowish shaded with pale red and somewhat striped with brighter red. *Dois* few, fine, gray.

Calyx tube long, very narrow, funnel-form. *Stamens* extremely marginal touching the segments, a marked characteristic.

Core little above medium; cells open; core lines clasping. *Carpels* cordate. *Seeds* pale brown, short, plump, pointed.

Flesh of a greenish cast, firm, fine, breaking, juicy, brisk subacid, good.


Season October to January.

POMONA.

REFERENCES. 1. Downing, 1869:135. 2. Leroy, 1873:248. fig. 3. Hogg, 1884:155. 4. Bailey, *Am. Hort.*, 1892:237. 5. Beach, *N. Y. Sta. An. Rpt.*, 13:579. 1894. 6. *Id.*, *Gard. and For.*, 8:428. 1895. 7. *Id.*, *Can. Hort.*, 20:



POMONA (Reduced Size)



183. 1897. 8. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:356. 9. Beach and Clark, *N. Y. Sta. Bul.*, 248:139. 1904.

SYNONYMS. COX'S POMONA (1, 2, 3, 4, 5, 6, 7, 8). *Cox's Pomona* (9). *Dean's Codlin* (2). POMONA (9).

Fruit frequently large and when highly colored rather attractive, being mottled and striped with crimson over a clear pale yellow ground color. The fruit is not very uniform in size, often unsymmetrical, tender, easily bruised and not a very good keeper. In ordinary storage its commercial limit is October (9). The tree is a good grower, comes into bearing young and is an annual cropper, or nearly so, but only moderately productive. It is not recommended for commercial planting in New York.

Historical. Originated by Mr. Cox, Colnbrook Lawn near Slough, Bucks, England, who also originated Cox Orange (3).

TREE.

Tree medium size, moderately vigorous with branches moderately stout, spreading and often drooping. *Form* roundish. *Twigs* straight, stocky, long; internodes medium. *Bark* rather bright reddish-brown mingled with olive-green, thinly overlaid with narrow streaks of gray scarf-skin. *Lenticels* rather numerous, conspicuous, medium to sometimes large, roundish or sometimes elongated. *Buds* medium to large, broad, obtuse, appressed, somewhat pubescent. *Leaves* rather large, broad, dark green; base of petioles red.

FRUIT.

Fruit medium to very large, fairly uniform in size but not in shape. *Form* oblate conic, very irregular, ribbed; sides usually unequal. *Stem* usually short, moderately thick. *Cavity* large, acuminate to acute, deep, broad, rather symmetrical or sometimes compressed, irregularly russeted. *Calyx* medium size, open or nearly closed; lobes separated at base, short, rather broad, obtuse to acute, reflexed. *Basin* deep, medium in width to rather wide, abrupt, smooth or ridged, slightly wrinkled.

Skin thin, rather tender, waxy, smooth, greenish-yellow or pale yellow partly blushed with crimson, and mottled and narrowly striped with carmine. *Dots* scattering, small, inconspicuous, usually whitish and submerged, sometimes gray or russet.

Calyx tube wide, conical, sometimes extending to the core. *Stamens* median to nearly marginal.

Core medium to small, somewhat abaxile; cells open or closed, symmetrical; core lines clasping. *Carpels* ovate to broadly roundish or elliptical, emarginate. *Seeds* medium to small, wide, short, very plump, flat, obtuse, dark brown.

Flesh whitish, not very firm, rather fine, crisp, tender, juicy, subacid, sprightly, good to very good for culinary use.

Season September and October.

PORTER.

REFERENCES. 1. Kentrick, 1832:27. 2. *Mag. Hort.*, 1:363. 1835. 3. Manning, 1838:51. 4. *Ib.*, *Mag. Hort.*, 7:48. 1841. 5. Downing, 1845:92. fig. 6. Floy-Lindley, 1846:411 app. 7. *Horticulturist*, 1:196, 256. 1846. 8. Hovey, *Mag. Hort.*, 14:116. 1848. fig. 9. Cole, 1849:107. fig. 10. Thomas, 1849:157. fig. 11. Waring, 1851:28. 12. Barry, 1851:285. 13. Emmons, *Nat. Hist. N. Y.*, 3:37. 1851. col. pl. No. 27 and fig. 14. Hovey, 1:43. 1851. col. pl. and fig. 15. *Horticulturist*, 7:217. 1852. 16. *Am. Pom. Soc. Cat.*, 1852. 17. Elliott, 1854:98. 18. Hooper, 1857:71. 19. Gregg, 1857:47. fig. 20. Warder, 1867:673. fig. 21. Fitz, 1872:153, 162. 22. Leroy, 1873:580. fig. 23. Downing, 1881:11, 12 index, app. 24. *Rural N. Y.*, 47:713. 1888. 25. Wickson, 1889:243. 26. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 27. Bailey, *Am. Hort.*, 1892:246. 28. Burrill and McCluer, *Ill. Sta. Bul.*, 45:336. 1896. 29. Waugh, *Vt. Sta. An. Rpt.*, 14:303. 1901. 30. Budd-Hansen, 1903:152. fig.

SYNONYMS. *Jennings* (23). PORTER (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30). *Yellow Summer Pearmain* (23).

This fruit is of very fine dessert quality. It is also highly esteemed for canning and other culinary uses, because when it is cooked it is not only excellent in quality but it retains its form remarkably well. The fruit is yellow, faintly marked with red, decidedly attractive for an apple of its class. It does not stand shipping very well because the skin is rather tender and readily shows marks of handling. Since it is quite variable in its season of ripening there is apt to be considerable loss from dropping unless more than one picking is made. It varies in size from large to small, with a considerable percentage of the crop undersized or otherwise unmarketable. The tree is vigorous, compact, hardy, comes into bearing early and is a pretty reliable biennial cropper. Fifty years ago it was the principal September apple in the Boston market (1, 3, 5, 9, 18, 19), and in spite of the fact that it is not red it continues to sell well in that market (29). It is also in good demand in many local markets. It is desirable for planting for home use or for some local markets, but generally it is not regarded as a profitable commercial variety by New York fruit growers.

Historical. Originated about 1800 with Rev. Samuel Porter, Sherburne, Mass. (1, 3, 5, 14), and up to about 1850 its cultivation was confined principally to the vicinity of its origin. It gradually became very widely known and has become well disseminated in many of the more important apple-growing regions of the country. Old trees of it are occasionally found in New York orchards but it is now seldom planted here.



PORTER

TREE.

Tree medium to large, vigorous. *Form* roundish or somewhat spreading. *Twigs* rather slender, very short-jointed, with prominent shoulders (14). *Buds* medium in size.

FRUIT.

Fruit small to large, usually rather large. *Form* oblong inclined to conic, rather truncate at base and with apex oblique and somewhat ribbed. *Stem* medium to rather short, moderately thick to slender, sometimes knobbed, curved. *Cavity* acute to acuminate, medium to deep, medium in width to rather narrow, symmetrical or compressed, sometimes lipped, usually faintly russeted. *Calyx* rather large, closed or partly open; lobes usually separated at base, short, rather narrow, broadly acute. *Basin* moderately deep to shallow, rather narrow, abrupt, broadly furrowed and wrinkled.

Skin moderately thin, smooth, rather glossy, clear bright yellow with a faint blush, usually rather obscurely striped with darker red marked with scattering red dots. *Dots* usually medium to very small, often submerged, green with whitish center, sometimes russet.

Calyx tube rather short, wide, broadly conical. *Stamens* median to basal.

Core medium to large, axile to somewhat abaxile; cells partly open to wide open; core lines meeting or slightly clasping. *Carpels* broadly ovate to elliptical, mucronate. *Seeds* below medium to rather large, plump, rounded, acute.

Flesh yellow, fine, crisp, tender, juicy, subacid, agreeably aromatic, sprightly, good to very good for either dessert or culinary uses.

Season. It begins to ripen in September and continues in use till November or later.

POUND SWEET.

REFERENCE. 1. Downing, 1869:311.

SYNONYMS. None.

This name has been applied to several varieties of large sweet apples. Downing mentions one which is large, roundish conic, greenish-yellow with slight red in sun; flesh yellowish, tender, moderately juicy, sweet; season September and October; and another which is large, roundish, red; flesh whitish, moderately juicy, aromatic, sweet (1). Others are described by other authors and some which are known locally by this name have perhaps never been described in any publication.

The variety most commonly known in Central and Western New York under the name Pound Sweet is large, globular, marbled with yellow and green and streaked with whitish scarf-skin. It is described under its correct name Pumpkin Sweet on page 171.

PRIMATE.

REFERENCES. 1. Cowles, *Mag. Hort.*, 16:450. 1850. *fig.* 2. *Ib.*, 17:506. 1851. 3. *Am. Pom. Soc. Cat.*, 1854. 4. *Mag. Hort.*, 20:241. 1854. 5. Elliott, 1854:153. *fig.* 6. *Mag. Hort.*, 21:62. 1855. 7. Gregg, 1857:38. 8. Downing, 1857:93. *fig.* 9. *Horticulturist*, 14:471. 1859. 10. *Mag. Hort.*,

27:69. 1861. 11. *Horticulturist*, 17:103, 150. 1862. 12. *Mag. Hort.*, 32:17. 1866. 13. Warder, 1867:643. fig. 14. Downing, 1869:313. 15. Fitz, 1872:161. 16. Thomas, 1875:196. 17. Downing, 1881:11, 12 index, app. 18. Barry, 1883:334. 19. *Rural N. Y.*, 47:649. 1888. 20. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 21. Blackwell, *Rural N. Y.*, 50:447. 1891. 22. Bailey, *Am. Hort.*, 1892:247. 23. Hoskins, *Rural N. Y.*, 53:248. 1894. 24. *Can. Hort.*, 17:252. 1894. 25. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 8:9. 1901. figs. 26. Budd-Hansen, 1903:153. fig. 27. *Rural N. Y.*, 62:740. 1903.

SYNONYMS. *Belle Rose* (17). *Early Baldwin* (17). *Early Tart Harvest* (14). *Harvest* (2). *Highland Pippin* (17). *July Apple* (14). *NORTH AMERICAN BEST* (9). *North American Best* (14). *Powers* (2, 14). *PRIMATE* (2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27). *ROUGH AND READY* (1). *Rough and Ready* (2, 5, 7, 8, 14). *Scott* (14). *Sour Harvest* (9, 14). *Sour Bough* (9, 14).

A dessert apple, pale yellow or whitish, often slightly blushed; in season in August and September. It is well known throughout the state, but not much grown except for home use. It is commonly considered the best apple of its season for the home orchard because the tree is a pretty good grower and a reliable cropper, and the fruit ripens in succession during a period of several weeks and is of fine flavor and excellent quality particularly for dessert use. Being less attractive than a red apple it is in demand in market only where its fine quality is known. The fruit ripens unevenly and it should be picked from time to time as it matures to prevent loss from the dropping of the fruit. In some localities the tree has proved somewhat tender, not very long-lived and rather susceptible to the attacks of the apple canker, but, generally speaking, as grown throughout Central and Western New York, particularly where it has been topworked upon good thrifty stock, the tree is a pretty good grower, moderately long-lived and reliably productive. Often it yields very heavy crops biennially with lighter crops, or none, on alternate years, but in some localities it is almost an annual bearer.

Historical. This variety was disseminated by traveling grafters in Central and Western New York as much as fifty years ago. In 1850, Charles P. Cowles of Syracuse in a communication to the *Magazine of Horticulture* stated: "As it is not known in this place, nor state, by the best judges, I safely think it is a seedling. I found a few trees in Onondaga county in a town of the same name which had been circulated by grafts but nothing further could be traced of its origin. * * * I propose to call it 'Rough



PRIMATE

and Ready' from the fact of its being first tested during that campaign. * * * Messrs. Downing, Barry and Thomas think it a new variety. * * * It is but little known as yet but where it is its qualities are much esteemed" (1). The following year Mr. A. Fahnestock, a nurseryman of Syracuse, showed that this variety had been disseminated in Ontario and Monroe counties as well as in Onondaga county and that it was generally known by the name of Primate. Recently John T. Roberts of Syracuse has taken the trouble to look up the history of this variety and is convinced that the original seedling tree grew in the town of Camillus, Onondaga county, and through his efforts a bronze tablet has been erected to mark the spot. The tablet bears the following inscription: "On this farm Calvin D. Bingham about 1840 produced the marvelous Primate apple, named by Charles P. Cowles. God's earth is full of love to man" (27). In 1854 it was listed by the American Pomological Society (3) as a valuable variety. It is listed by various nurserymen in the region from the Atlantic to the Middle West but apparently is but little known either in the Southwest or the Northwest (22).

TREE.

Tree below medium to rather large, usually moderately vigorous. *Form* upright spreading to roundish, rather dense. *Twigs* short, straight, stout with large terminal buds; internodes short. *Bark* dull brown mingled with green, heavily streaked with scarf-skin, pubescent and rather rough. *Lenticels* scattering, medium size, oblong, raised. *Buds* prominent, large, broad, plump, acute, pubescent.

FRUIT.

Fruit medium or above, sometimes large. *Form* roundish conic to oblate conic, often distinctly ribbed. *Stem* short to medium, thick. *Cavity* acute, deep, broad, distinctly furrowed. *Calyx* medium size, closed; lobes long, narrow. *Basin* moderately shallow to rather deep, medium in width to rather narrow, abrupt, furrowed and wrinkled.

Skin thin, moderately tender, smooth, light green, often changing to whitish, sometimes with faint yellowish tinge, often slightly blushed but not striped. *Dots* scattering, numerous, rather small, submerged or russet.

Calyx tube rather large, rather long, broadly conical to somewhat funnel-shape. *Stamens* median. *Core* medium to large, nearly axile to somewhat abaxile with hollow cylinder in the axis; cells symmetrical, open; core lines clasping. *Carpels* cordate. *Seeds* medium to rather large, moderately wide, plump, acute.

Flesh whitish, fine, crisp, very tender, juicy, subacid, aromatic, sprightly, very good to best.

Season August and September.

PROLIFIC SWEETING.

REFERENCES. 1. Webster, *Montreal Hort. Soc. Rpt.*, 1881:54. 2. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:439. 3. *Ib.*, *Am. Pom. Soc. Rpt.*, 1887:46. 4. Van Derman, *U. S. Pom. Rpt.*, 1888:571. fig. 5. Hoskins, *Rural N. Y.*, 47:646. 1888. 6. *N. Y. Sta. An. Rpt.*, 7:91. 1888. 7. Bailey, *Am. Hort.*, 1892:247. 8. Beach, *N. Y. Sta. An. Rpt.*, 14:254. 1895. 9. Munson, *Me. Sta. Rpt.*,

1896:71. 10. Waugh, *Vt. Sta. An. Rpt.*, 14:304. 1901. 11. Munson, *Me. Sta. Rpt.*, 1902:84, 86, 88. 12. Hansen, *S. D. Sta. Bul.*, 76:88. 1902. fig. 13. Budd-Hansen, 1903:154.

SYNONYMS. *No. 351* (3, 4, 12). *Plodowitka CUADKAJA* (1). *Plodowitka Cuadkaja* (3). *Plodowitka CAUDKAJA* (2). *PROLIFIC SWEET* (10). *Prolific Sweet* (1). *PROLIFIC SWEETING* (3, 4, 5, 6, 7, 8, 9, 11, 12, 13). *Prolific Sweeting* (2).

A Russian variety which in size and color resembles Yellow Transparent but in form is roundish oblate and somewhat irregular. It has proved to be a valuable sweet apple for autumn use in Northern New England (5, 11, 12, 13). Worthy of trial in Northern New York when a variety of this class is desired.

Historical. Imported from Russia by the United States Department of Agriculture in 1870. It was received for testing at this Station from Dr. T. H. Hoskins, Newport, Vt., in 1888 (6). It has thus far been but little disseminated in this state.

TREE.

Tree moderately vigorous. *Form* upright spreading to roundish, open. *Twigs* long, curved, stout; internodes short. *Bark* brown, tinged with green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, medium size, round, not raised. *Buds* medium size, broad, flat, obtuse, appressed, pubescent.

FRUIT.

Fruit medium or above. *Form* roundish oblate, somewhat irregular. *Stem* medium size. *Cavity* deep, acute to acuminate, ribbed, russeted and with some outspreading russet rays. *Calyx* closed. *Basin* shallow, wide, wrinkled.

Flesh white, crisp, fine-grained, mildly sweet, good.

Season late August, September and October.


PUMPKIN RUSSET.

REFERENCES. 1. Kenrick, 1832:37. 2. *Mag. Hort.*, 1:363. 1835. 3. Downing, 1845:93. 4. Thomas, 1849:146. 5. Cole, 1849:115. 6. Elliott, 1854:152. 7. Hooper, 1857:73. 8. Warder, 1867:566. 9. Downing, 1881:111 index, app. 10. Bailey, *An. Hort.*, 1892:247. 11. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:53. 1903. 12. Beach and Clark, *N. Y. Sta. Bul.*, 248:139. 1904.

SYNONYMS. *Flint Russet* (3, 6). *Kingsbury Russet* (9). *PUMPKIN RUSSET* (3, 4, 6, 7, 8, 9, 10, 11, 12). *Pumpkin Russet* (5). *PUMPKIN SWEET* (5). *Pumpkin Sweet* (3). *PUMPKIN SWEETING* (2). *PUMPKIN SWEETING of New England* (1). *Sweet Russet* (3, 4, 6). *York Russet* (6).

This is a very large, round, yellowish-russet apple, sweet, rich, very good for baking but of little value for other purposes. The tree is a vigorous, rapid grower, hardy, moderately long-lived and yields fair to good crops biennially or in some cases almost annually. It is not a profitable commercial variety and is now seldom cultivated even for home use.

Historical. Pumpkin Russet is an old New England variety. Kenrick (1) in 1832 described it under the name Pumpkin Sweeting of New Eng-



PUMPKIN RUSSET

land. In 1849 Cole (5) described it under the name Pumpkin Sweet giving Pumpkin Russet as a synonym. In 1845 it was described by Downing (3) under the name Pumpkin Russet which is the name now generally accepted for it by pomologists. It is still occasionally listed by nurserymen but is now seldom planted in New York.

TREE.

Tree large, vigorous or very vigorous, at first upright but eventually becoming roundish or spreading, open; branches long, stout, curved. *Twigs* short, curved, stout; internodes long. *Bark* dark reddish-brown tinged with olive-green, lightly streaked with scarf-skin, pubescent. *Lenticels* quite numerous, oblong, slightly raised. *Buds* large, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit large, fairly uniform in size and shape. *Form* oblate or somewhat inclined to conic, sometimes irregular, faintly ribbed, often compressed. *Stem* medium to short, moderately slender to thick. *Cavity* acute to acuminate, usually deep, moderately broad to rather narrow, nearly smooth, sometimes slightly furrowed. *Calyx* large, closed or slightly open; lobes long, narrow, acute to acuminate. *Basin* small to medium, rather shallow to moderately deep, medium in width, somewhat abrupt, slightly furrowed and wrinkled.

Skin thick, rather tough, greenish or yellowish sometimes with bronze blush on exposed cheek, more or less covered with russet patches or netted veins of russet. *Dots* large and small, scattering, usually russet, irregular.

Calyx tube short, wide, broadly conical to nearly urnshape. *Stamens* median to basal.

Core above medium to below, abaxile; cells usually open, symmetrical but not uniformly developed; core lines slightly clasping. *Carpels* broadly ovate, very slightly emarginate, sometimes tufted. *Seeds* moderately light brown, rather large, wide, flat, acute.

Flesh greenish-white or yellowish, firm, rather coarse, tender, juicy, sweet, good.

Season September and October or in cold storage extending to January 1 (12).

PUMPKIN SWEET.

REFERENCES. 1. N. E. Farmer, 1834 (cited by 20). 2. Downing, 1845:89. fig. 3. Emmons, *Nat. Hist. N. Y.*, 3:48. 1851. 4. *Mag. Hort.*, 20:241. 1854. 5. Elliott, 1854:152. 6. Hooper, 1857:56. 7. *Am. Pom. Soc. Cat.*, 1862. 8. *Horticulturist*, 17:150. 1862. 9. Warder, 1867:527. 10. Barry, 1883:339. 11. Thomas, 1885:216. 12. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 13. Bailey, *An. Hort.*, 1892:247. 14. Munson, *Me. Sta. Rpt.*, 1893:133. 15. Miller, *Rural N. Y.*, 53:278. 1894. 16. Taylor, *Am. Pom. Soc. Rpt.*, 1895:198. 17. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:53. 1903. 18. Budd-Hansen, 1903:155. fig. 19. Beach and Clark, *N. Y. Sta. Bul.*, 248:139. 1904. 20. Ragan, *U. S. B. P. I. Bul.*, 56:244. 1905.


SYNONYMS. *Lyman's Large Yellow* (20). LYMAN'S PUMPKIN SWEET (2, 3, 6, 9, 10, 11). *Lyman's Pumpkin Sweet* (5, 14, 17, 19, 20). *Pound*

Sweet (9, 11, 12, 14, 17, 18, 19). *PUMPKIN SWEET* (1, 4, 5, 7, 8, 12, 13, 14, 15, 16, 17, 18, 19, 20). *Pumpkin Sweet* (10, 11). *Pumpkin Sweeting* (20). *Rhode Island Sweet* (20). *Round Sweet* (20). *Sweet Lyman's Pumpkin* (20). *Vermont Pumpkin Sweet* (10, 20). *Vermont Sweet* (20). *Yankee Apple* (5, 20).

Fruit large to very large, marbled with light and dark green and streaked over the base with whitish scarf-skin; well colored specimens eventually become quite yellow and sometimes are faintly bronzed on the exposed cheek. It is never marked with red, nor is it russeted except about the cavity. So far as we know all other varieties which have been cultivated under the name Pumpkin Sweet are either russeted or marked with red.

This is the variety generally known in Central and Western New York as Pound Sweet, and it commonly appears under this name in market quotations. By many it is esteemed as one of the best sweet apples of its season for baking and for canning or stewing with quinces, but generally it is not valued for dessert because it is rather coarse and has a peculiar flavor. It often sells well in local or special markets, and there is a limited demand for it in the general trade. Its keeping qualities differ greatly in different localities and in different seasons. As grown in Western New York it comes in season early in October. The rate of loss in ordinary storage is usually high during the fall, and the season closes in December or early January, although in some years a considerable portion of the fruit may remain sound till midwinter or later (19).

The tree is a good strong grower, rather long-lived, fairly hardy and generally healthy, but it sometimes suffers from winter injury, sunscald and canker. It appears to thrive particularly well on well fertilized gravelly or sandy loam, with well drained subsoil. Under right conditions it is a pretty reliable cropper, yielding good crops biennially. The crop ripens somewhat unevenly and often there is a considerable loss from water-cored fruit and from windfalls, but on the other hand there is a small percentage of loss in undersized or deformed apples. In order to lessen the loss from windfalls it is well to plant this variety in a location that is sheltered from prevailing winds.



PUMPKIN SWEET

Historical. Originated in the orchard of S. Lyman, Manchester, Conn. (9). It has been distributed throughout this state for more than fifty years, but nowhere has it been largely planted. Generally speaking it is not grown so extensively now as it was formerly, but in a few localities its cultivation for commercial purposes is increasing somewhat.

TREE.

Tree medium to large, vigorous, with long, moderately stout branches. *Form* upright spreading or roundish, open. *Twigs* medium to short, straight or nearly so, stout to moderately slender; internodes short to medium. *Bark* reddish olive-green varying to brownish-red, uniformly mottled with thin scarf-skin, pubescent. *Lenticels* scattering, rather inconspicuous, small, roundish, not raised. *Buds* medium or below, plump, acute, free, slightly pubescent.

FRUIT.

Fruit large to very large, pretty uniform in size and shape. *Form* globular to roundish conic, sometimes irregularly elliptical or prominently ribbed. *Stem* very short, stout, often inserted under a lip or having itself a fleshy protuberance. *Cerise* acuminate, moderately shallow to rather deep, narrow to moderately wide, often somewhat furrowed or lipped, sometimes russeted and with outspreading russet. *Calyx* medium to rather large, open; lobes often separated at the base, short, broad, acute. *Basin* small to medium in size, moderately shallow to rather deep, narrow to moderately wide, abrupt to somewhat obtuse, often slightly furrowed or wrinkled.

Skin rather thin, tough, smooth, at first green but eventually clear yellow marbled with greenish-yellow. Stripes of whitish scarf-skin radiate from the cavity. Well colored specimens occasionally show a brownish-red blush but never a distinct red color. *Dots* conspicuous, whitish, often areolar with russet center.

Calyx tube rather wide, conical or elongated cone-shape or sometimes slightly funnel-form. *Stamens* median to basal.

Core medium to rather large, axile; cells symmetrical, closed or somewhat open, not uniformly developed; core lines clasping. *Carpels* thin, broadly roundish, but slightly emarginate if at all, often tufted. *Seeds* medium to rather small, wide, plump, acute, light brown, tufted.

Flesh tinged with yellow firm, medium in texture, crispness and juiciness, decidedly sweet with a peculiar flavor; good for culinary use and especially esteemed for baking.

Season October to January.

McCARTY is identical with Pumpkin Sweet in general characters and is sold as Pumpkin Sweet, but it appears to differ enough from the type in certain characteristics to entitle it to recognition as a distinct strain. As compared with typical Pumpkin Sweet the fruit of McCarty averages smaller, is not so yellow and keeps longer. B. J. Case, Sodus, N. Y., who grows McCarty commer-

cially, reports: "The tree does not produce any water-cored fruit except when the crop is light. It appears to be fully as productive as Pumpkin Sweet. In common storage it is not unusual to keep McCarty later than January. In quality it is not so good as Pumpkin Sweet. The origin of this type is unknown."

(I) QUINCE (*of Cole*).

REFERENCES. 1. Cole, 1849:99. *fig.* 2. Downing, 1857:130. 3. Hovey, *Mag. Hort.*, 27:71. 1861. *fig.* 4. *Am. Pom. Soc. Rpt.*, 1862. 5. Warder, 1867:645. *fig.* 6. Downing, 1869:319. 7. Lyon, *Mich. Sta. Bul.*, 143:201. 1897. 8. Budd-Hansen, 1903:156. *fig.* 9. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:53. 1903. 10. Ragan, *ib.*, 56:246. 1905.

SYNONYMS. COLE'S QUINCE (1, 2, 4). *Cole's Quince* (3, 5, 6, 10). QUINCE (3, 5, 6). QUINCE (Cole) (10). QUINCE Cole (7, 8, 9). *Turn Off Lane* (10).

This is a very good variety for home use and is considered by some New York fruit growers fairly profitable in commercial orchards. The fruit is large, angular, yellow with white aromatic subacid flesh of very good quality; in season in late summer and early autumn. Commercial limit in cold storage, November 1 (9). The tree is of good medium size, spreading, a good grower, very hardy, comes into bearing moderately young, and is a reliable cropper. It is not extensively cultivated in New York.

Historical. Originated at Cornish, Me., from whence it was disseminated about fifty years ago (1).

FRUIT.

Fruit. Cole gave the following description of this fruit in 1849 (1): "Fruit large to very large; flattish-conical; ribbed; bright yellow, seldom a brown cheek, stem short, in a deep cavity; calyx large, in a deep basin; flesh when first ripe, firm, juicy, pleasant acid, and first-rate for cooking. When very mellow, remarkably tender, of a mild, rich, high quince flavor and aroma. When in perfection we have never seen its superior. July to September."

(II) QUINCE (*of Cox*).

REFERENCES. 1. McMahon, *Am. Gard. Cal.*, 1806:585. 2. Cox, 1817:138. *fig.* 3. Thacher, 1822:132. 4. Cole, 1849:115. 5. Thomas, 1849:158. 6. Downing, 1857:181. 7. *Horticulturist*, 23:146. 1868. *fig.* 8. Downing, 1869:319. 9. Fitz, 1872:153. 10. Downing, 1872:10 index, app. 11. Ragan, *U. S. B. P. I. Bul.*, 56:246. 1905.

SYNONYMS. *Pear Apple* (8, 11). QUINCE (1, 2, 3, 5, 6, 8, 9, 10). *Quince Apple* (11). QUINCE (Cox) (11). QUINCE OF COX (4, 7). *Quince of Cox* (11). *Seneca Favorite* (10). *Seneca Spice* (11, 78).

A variety which is distinct from the Quince of Cole was described by Coxé in 1817. According to Downing, it is in appearance much like a large Yellow Newtown, and the young wood is of a different shade of color from that of the Quince of Cole, being dull reddish-brown instead of clear reddish-brown, with buds small, reddish and pointed, instead of short, abrupt and prominent. The following is the description of the fruit given by Coxé (2): "The size of the apple is large; the shape flat; the skin, when fully ripe, is yellow; the flesh rich, yellow and juicy—in appearance it somewhat resembles a large yellow Newtown Pippin. It came originally from the state of New York; ripens in November."

This variety is unknown to us, and we have received no report concerning it from any of our correspondents.

RAMSDELL SWEET.

REFERENCES. 1. *Mag. Hort.*, 4:47. 1838. 2. Manning and Ives, 1844:41. 3. Downing, 1845:137. 4. *Mag. Hort.*, 12:475. 1846. 5. Thomas, 1849:161. 6. Cole, 1849:118. 7. Emmons, *Nat. Hist. N. Y.*, 3:76. 1851. 8. Elliott, 1854:131. 9. Hooper, 1857:74. 10. *Am. Pom. Soc. Cat.*, 1862. 11. Warder, 1867:664. fig. 12. Downing, 1869:163. 13. Fitz, 1872:121, 145. 14. Leroy, 1873:605. fig. 15. Barry, 1883:352. 16. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 17. Bailey, *Am. Hort.*, 1892:247. 18. Burrill and McCluer, *Ill. Sta. Bul.*, 45:337. 1896. 19. Farrand, *Mich. Sta. Bul.*, 205:46. 1903. 20. Budd-Hansen, 1903:159. fig.

SYNONYMS. *Avery Sweet* (12, 14). *Avery Sweeting* (8). *ENGLISH SWEET* (12, 18). *English Sweet* (14, 15, 16, 20). *ENGLISH SWEETING* (4, 8). *Hurlbut* (14). *Hurlbut Sweet* (12). *RAMSDALE'S SWEETING* (9). *RAMSDELL* (14, 19). *RAMSDELL'S RED* (11). *Ramsdell Red Sweet* (18). *RAMSDELL'S RED SWEETING* (1). *RAMSDELL SWEET* (10, 15, 16, 17, 20). *Ramsdell Sweet* (3, 8, 12). *RAMSDELL'S SWEETING* (3, 5, 6, 7, 12, 13). *Ramsdell's Sweeting* (8, 14). *RAMSDELL'S RED PUMPKIN SWEET* (2). *Ramsdell's Red Pumpkin Sweet* (3, 5, 7, 8, 12). *Randall's Red Winter* (12, 14). *Red Pumpkin Sweet* (3, 8, 12). *Red Pumpkin Sweeting* (9). *Reindell's Large* (14).

This is an attractive red apple of good size and very good quality, in season from midautumn to midwinter. The tree is a pretty good grower and an early bearer, yielding some fruit annually or nearly annually, but in many cases it is not satisfactorily productive and the fruit is not very uniform in size and quality, so that there is a considerable loss from unmarketable fruit. Not recommended for commercial planting in New York.

Historical. This variety was first brought to notice by being exhibited before the Massachusetts Horticultural Society by the Rev. H. S. Ramsdell, Thompson, Conn., and was named Ramsdell's Red Sweeting in compliment to him (1). Downing described it in 1845 (3) under the name Ramsdell's Sweeting, but in the 1869 edition (12) it was described under the name English Sweet, the name previously recognized by Elliott (8) but upon what authority we have been unable to learn. In 1862 (10) it was entered on the catalogue of the American Pomological Society under the name Ramsdell Sweet which name has been retained in that catalogue up to the present time. It is also commonly catalogued by nurserymen under the name Ramsdell or Ramsdell Sweet (17).

TREE.

Tree medium size, vigorous or moderately vigorous. *Form* upright, open. *Twigs* long, curved, moderately stout; internodes medium. *Bark* brown, streaked with heavy scarf-skin; slightly pubescent near tips. *Lenticels* scattering, small to medium, oval, raised. *Buds* medium size, plump, obtuse, free, pubescent.

FRUIT.

Fruit above medium, sometimes nearly large, somewhat variable in size, fairly uniform in shape. *Form* oblong conic to roundish conic, often somewhat elliptical and faintly ribbed. *Stem* short to medium in length, moderately slender, often red. *Cavity* acuminate, deep, rather broad, quite symmetrical, often with some stellate russet. *Calyx* small to medium, closed or slightly open; lobes narrow, acute to acuminate. *Basin* rather small, sometimes oblique, medium to rather deep, narrow to medium in width, abrupt, faintly furrowed and wrinkled.

Skin thin, rather tough, smooth, yellow nearly overspread with attractive dark red, or entirely red with obscure splashes and stripes of carmine, overspread with blue bloom. *Dots* many, distinct, conspicuous, small to rather large, pale yellow or grayish, often submerged. *Prevailing effect* red, attractive.

Calyx tube rather large, long, cylindrical to narrow funnel-shape. *Stamens* median.

Core small to medium, axile to somewhat abaxile; cells symmetrical but not uniformly developed, closed or slightly open; core lines clasping. *Carfals* ovate to nearly roundish. *Seeds* below medium, rather narrow, plump, acute.

Flesh tinged with yellow, firm, fine, tender, juicy, very sweet, good to very good.

Season October to February.

RASPBERRY.

REFERENCES. 1. *Ia. Hort. Soc. Rpt.*, 1883:435. 2. *Montreal Hort. Soc. Rpt.*, 1883:90. 3. *Ia. Agr. Coll. Bul.*, 1885:11. 4. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886:77. 5. *U. S. Pom. Rpt.*, 1891:390. 6. Thomas, 1897:252. fig. 7. *Am. Pom. Soc. Cat.*, 1899:19. 8. Hansen, *S. D. Sta. Bul.*, 76:89. 1902. fig. 9. Budd-Hansen, 1903:159. fig.

SYNONYMS. MALINOWSKOE (1, 2). *Malinowskoe* (3). No. 288 (3, 5, 8). RASPBERRY (3, 4, 5, 6, 7, 8, 9). *Raspberry* (1, 2). *Red Check* (6).

A Russian apple, small, fine dark red, sprightly subacid; season July and August. Hansen states (8, 9) that it is exceedingly productive and a good substitute for Red June where that variety winter-kills.

RED AND GREEN SWEET.

REFERENCES. 1. Coxe, 1817:162. 2. Thacher, 1822:129. 3. Kenrick, 1832:38. 4. Thomas, 1849:135. 5. Elliott, 1854:176. 6. Downing, 1857:221. 7. Hooper, 1857:78. 8. Warder, 1867:729. 9. Downing, 1872:10 index, app.

SYNONYMS. *Large Red Sweeting* (5). LARGE RED AND GREEN SWEETING (2). PRINCE'S LARGE RED AND GREEN SWEETING (1). *Prince's Large Red and Green Sweeting* (3). *Prince's Red and Green Sweet* (5). *Red Bough* (5). RED AND GREEN SWEET (4, 5, 6, 7, 8, 9). RED AND GREEN SWEETING (3). *Saille Sweet* (9). *Virginia Sweet* (9).

This is a very large green apple, striped with red, rather attractive when well colored. It does not rank very high in flavor or quality but is suitable for baking. Season August and September. The fruit does not last long after it becomes ripe. The tree is medium to large, moderately vigorous to vigorous, long-lived and a reliable cropper yielding heavy crops annually. Although some find it a profitable apple to grow for local market it is not worthy of being recommended for general planting.

Historical. This is an old variety which was described by Coxe (1). It was formerly grown to a comparatively limited extent in some portions of New York and in adjoining states. Occasionally a tree of it is still found in some of the oldest orchards of the state but it is fast going out of cultivation.

TREE.

Tree large. *Form* upright spreading to roundish, open. *Twigs* short, straight, stout with large terminal buds; internodes short. *Bark* brown mingled with olive-green, heavily coated with gray scarf-skin; pubescent. *Lenticels* scattering, medium size, oval, not raised. *Buds* large, rather prominent, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit large to very large. *Form* oblong conic, rather strongly ribbed; sides unequal. *Stem* short to medium, moderately slender. *Cavity* nearly acuminate, deep, broad, furrowed, sometimes lipped, sometimes thinly russeted. *Calyx* closed or partly open; lobes moderately long, narrow, acute. *Basin* medium in depth to rather deep, medium in width to rather narrow, wrinkled.

Skin thin, tender, smooth, green changing to yellow, more or less blushed and partly overspread with pinkish-red irregularly striped and splashed with rather bright carmine. *Dots* conspicuous, numerous, large and scattering toward the cavity, small and very numerous toward the calyx.

Calyx tube long, funnel-shape.

Core rather large; cells open; core lines clasping. Carpels nearly roundish, tufted. Seeds rather small, plump, acute.

Flesh white, fine, very tender, moderately juicy, of pleasant sweet flavor and fair to good quality.

Season August and September.

RED ASTRACHAN.

REFERENCES. 1. *London Hort. Soc. Trans.*, 4:522 (cited by 2). 2. Forsyth, 1824:131. 3. *Pom. Mag.*, 1830:No. 123, col. pl. 4. *London Hort. Soc. Cat.*, 1831:No. 31. 5. Kenrick, 1832:90. 6. Floy-Lindley, 1833:4. 7. Kenrick, 1835:58. 8. *Mag. Hort.*, 1:391. 1835. 9. Manning, 1838:50. 10. *Ib.*, *Mag. Hort.*, 7:48. 1841. 11. Downing, 1845:75. fig. 12. *Horticulturist*, 1:146. 1846. 13. Hovey, *Mag. Hort.*, 14:15. 1848. fig. 14. Cole, 1849:98. fig. 15. Goodrich, 1849:49. 16. Thomas, 1849:143. 17. Waring, 1851:29. 18. Barry, 1851:279. 19. Emmons, *Nat. Hist. N. Y.*, 3:14. 1851. col. pl. No. 53. 20. Hovey, 1:35. 1851. col. pl. and fig. 21. *Mass. Hort. Soc. Trans.*, 1852:103. col. pl. 22. *Am. Pom. Soc. Cat.*, 1852. 23. *Horticulturist*, 7:437. 1852. 24. Elliott, 1854:103. fig. 25. *Horticulturist*, 10:443. 1855. fig. 26. Hooper, 1857:13, 77. 27. Gregg, 1857:38. fig. 28. *Mag. Hort.*, 30:162. 1864. 29. *Ib.*, 32:17, 51. 1866. 30. Warder, 1867:456. fig. 31. Downing, 1869:323. fig. 32. Todd, 1871:210. fig. 33. Fitz, 1872:143, 160, 172. 34. Leroy, 1873:82. fig. 35. Hatch, *Horticulturist*, 29:51. 1874. 36. *Montreal Hort. Soc. Rpt.*, 1876:8. 37. Hogg, 1884:185. 38. Wickson, 1889:243. 39. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 40. *Can. Hort.*, 14:261. 1891. 41. Bailey, *An. Hort.*, 1892:234, 247. 42. *Am. Gard.*, 15:404. 1894. fig. 43. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 2:10. 1895. fig. 44. Burrill and McCluer, *Ill. Sta. Bul.*, 45:337. 1896. 45. Waugh, *Vt. Sta. Bul.*, 61:31. 1897. 46. Bunyard, *Four. Roy. Hort. Soc.*, 1898:354. 47. Craig, *Cyc. of Am. Hort.*, 1901:1404. 48. Waugh, *Vt. Sta. An. Rpt.*, 14:305. 1901. 49. *Can. Hort.*, 25:305. 1902. figs. 50. Hansen, *S. D. Sta. Bul.*, 76:90. 1902. 51. Bruner, *N. C. Sta. Bul.*, 182:21. 1903. 52. Budd-Hansen, 1903:160. fig.

SYNONYMS. *Abe Lincoln* (31, 32, 34, 48). *AMERICAN RED* (4, 46). *Anglesea Pippin* (37). *ASTRACHAN ROUGE* (34). *ASTRACHAN* (23). *ASTRACHAN RED* (2, 18). *Astrachan Rouge* (31). *Astrakhan Rouge* (32). *Deterding's Early* (31, 32). *Hamper's American* (37). *RED ASTRACHAN* (6, 7, 8, 9, 10, 19). *RED ASTRACHAN* (1, 3, 5, 11, 12, 13, 14, 15, 16, 17, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 33, 35, 36, 37, 39, 40, 41, 42, 43, 44, 45, 47, 48, 49, 50, 51, 52). *Red Astrachan* (34). *RED ASTRACHAN* (32). *Rother Astrakhan* (31, 32). *Vermillon d'Ete* (31, 32, 34). *Waterloo* (37).

This is a very beautiful early summer apple of good medium size, yellow, largely covered with light and dark red, presenting a striped appearance, and overspread with bluish bloom. It is generally well known throughout the state, being valued particularly for home use. It is fit for culinary purposes before it becomes fully ripe, so that for home use it is in season from late July to



RED ASTRACHAN

September. When fully ripe and mellow it is desirable for dessert use. The tree is of medium size, a good grower, moderately long-lived, comes into bearing rather young and is a reliable cropper, yielding moderate to good crops biennially or sometimes annually. The fruit hangs to the tree pretty well till it is fully ripe, but as the crop matures unevenly there is apt to be considerable loss from the dropping of the fruit unless several pickings are made. It is not very uniform in size, and a considerable amount of it is small or otherwise unmarketable. It is very perishable, and on this account not well adapted for shipping to distant markets. It is very common in local markets, but often the supply so much exceeds the demand that prices are consequently low.

Historical. Hogg states that Red Astrachan was imported from Sweden into England in 1816 but Lindley (6) states that "This very beautiful apple was imported from Sweden, and first fruited by William Atkinson, Esq., of Grove End, Paddington, in 1816." It was one of the first of the Russian apples imported into America. It was received by the Massachusetts Horticultural Society from the London (Eng.) Horticultural Society about 1835 but this was not the first importation for fruit of Red Astrachan was included in one of the exhibitions of the Massachusetts Horticultural Society in 1835 (8). It has become generally disseminated throughout the apple-growing districts of the continent and is commonly listed from the Atlantic to the Pacific and from Canada to the Gulf states.

TREE.

Tree medium to rather large, moderately vigorous to vigorous. *Form* upright spreading to roundish, rather dense. *Twigs* medium to long, curved, stout to rather slender; internodes long. *Bark* clear brown, lightly streaked with scarf-skin, pubescent near the tips. *Lenticels* quite numerous, medium to small, oblong, slightly raised. *Buds* medium in size, plump, obtuse to acute, free, pubescent.

FRUIT.

Fruit medium to sometimes large, not very uniform in size or shape. *Form* roundish to roundish oblate, inclined to conical, somewhat ribbed; sides a little unequal. *Stem* medium, rather slender, bracted. *Cavity* acute to sometimes obtuse, medium to deep, moderately broad, often thinly russeted with greenish russet, usually symmetrical, sometimes lipped. *Calyx* medium to rather large, open or closed; lobes rather long, moderately broad. *Basin* shallow, narrow to medium in width, obtuse, wrinkled.

Skin rather thin, moderately tender, smooth, pale yellow or greenish often nearly or quite overspread with light and dark red splashed and irregularly striped with deep crimson or carmine and covered with rather heavy distinct bluish bloom. *Dots* numerous, small, whitish.

Calyx tube long, funnel-form. *Stamens* median to marginal.

Core small, somewhat abaxile; cells closed or somewhat open; core lines clasping or sometimes nearly meeting. *Carpels* broadly ovate or obovate, slightly tufted. *Seeds* small, moderately wide, plump, obtuse.

Flesh white often strongly tinged with red, rather fine, crisp, tender, juicy, brisk subacid, aromatic, sometimes slightly astringent, good to very good.

Season late July to September.

RED GRAVENSTEIN.

REFERENCES. 1. Leroy, 1873:339. *fig.* 2. *Can. Hort.*, 16:362. 1893. 3. Lucas, Ed., *Ill. Handb. der Obstk.*, 1893:250.

SYNONYMS. GRAVENSTEIN ROUGE (1). RED GRAVENSTEIN (2, 3). *Rother Gravensteiner* (1).

Bud sports of the Gravenstein have appeared at different times which bear red fruit. For a statement concerning such sports the reader is referred to Gravenstein, page 84.

RED HOOK.

This is a large, late summer or early autumn apple which is being grown commercially to a limited extent in the vicinity of Red Hook, Dutchess county, N. Y. W. S. Teator, of Upper Red Hook, who furnished us with the fruit from which the following description was made, states that the variety originated in his locality and has been known under the name of Red Ox or Striped Ox. As the name Striped Ox has been applied to other varieties he proposes the name Red Hook for this apple. The tree is large, low branching and an annual cropper yielding heavy crops. The fruit is firm, smooth, bright, attractive, quite free from the attacks of fungi and good for culinary use but when overripe it becomes dry and worthless. It stands shipment well and is one of the earliest fruits of that locality that can be shipped to Europe.

FRUIT.

Fruit large to very large. *Form* roundish to roundish conic, somewhat elliptical, symmetrical, sometimes faintly ribbed. *Stem* short, thick to rather slender. *Cavity* acuminate, deep, broad, gently furrowed, smooth or partly russeted. *Calyx* medium to rather small, partly closed; lobes broad, obtuse to acute. *Basin* rather small, moderately deep, narrow to medium in width, abrupt.



Skin tough, attractive pale yellow, rather thinly mottled and blushed with lively red, and sparingly striped and splashed with bright carmine. *Dots* rather small, numerous, whitish or with russet center.

Calyx tube elongated funnel-shape approaching conical. *Stamens* median.

Core medium or below, axile or nearly so; cells pretty symmetrical, closed; core lines clasping the cylinder. *Carpels* broadly roundish. *Seeds* rather dark brown, medium size, irregular, obtuse.

Flesh firm, coarse, tender, juicy, sprightly subacid, good.

Season late summer or early autumn.



RED JUNE



RED JUNE.

REFERENCES. 1. Humrickhouse, *Mag. Hort.*, 14:389. 1848. fig. 2. Phoenix, *Horticulturist*, 4:472. 1850. 3. Barry, 1851:280. 4. *Horticulturist*, 10:87. 1855. 5. *Am. Pom. Soc. Cat.*, 1856. 6. *Horticulturist*, 12:473. 1857. 7. Downing, 1857:127. 8. Gregg, 1857:35. 9. Hooper, 1857:23. 10. Hovey, *Mag. Hort.*, 27:68. 1861. fig. 11. Warder, 1867:666. fig. 12. Downing, 1869:118. fig. 13. *Ib.*, 1872:10 index, app. 14. Fitz, 1872:151. 15. Thomas, 1875:190. 16. Wickson, 1889:242. 17. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 18. *Am. Pom. Soc. Cat.*, 1891. 19. Bailey, *Am. Hort.*, 1892:236. 20. *Ib.*, 1892:247. 21. Clayton, *Ala. Sta. Bul.*, 47:5. 1893. 22. Hoskins, *Rural N. Y.*, 53:248. 1894. 23. Stinson, *Ark. Sta. Rpt.*, 1894:44. 24. Massey, *N. C. Sta. Bul.*, 149:316. 1898. 25. Budd-Hansen, 1903:162. fig. 26. Bruner, *N. C. Sta. Bul.*, 182:21. 1903. 27. Farrand, *Mich. Sta. Bul.*, 205:46. 1903.

SYNONYMS. *Blush June* (12, 15, 23). *Blushed June* (7, 10). CAROLINA JUNE (4, 5, 6). *Carolina Red* (14). CAROLINA RED JUNE (2, 3, 7, 8, 9, 10, 11, 12, 15, 16, 19, 22, 23, 24). *Georgia June* (12). *Jones Early Harvest* (13). JUNE (1). *Knight's Red June* (12). RED JUNE (13, 17, 18, 20, 21, 27). *Red June* (7, 10, 12, 14, 15, 23). RED JUNEATING (14). RED JUNE, *Carolina* (25, 26). *Susy Clark* (13). *Wilson's June* (12).

An attractive little apple, deep red over yellow, tender, brisk subacid, very good. The tree is a moderate grower, a pretty reliable bearer, and commonly yields good crops. Some find it profitable because it is handsome and takes well in some markets. The crop ripens so unevenly that it should have two or three pickings in order to secure the fruit in good condition. The variety is more popular South and West than it is in New York.

Historical. This is a southern apple which is supposed to have originated in North Carolina. It has long been known in cultivation and is commonly listed by nurserymen, but it has not been planted to any considerable extent in New York.

TREE.

Tree moderately vigorous with short, moderately stout, curved branches. *Form* at first upright but becoming spreading or roundish. *Twigs* very short, straight, slender; internodes medium size. *Bark* dull brown, lightly mottled with scarf-skin; slightly pubescent. *Lenticels* scattering, small, oblong, not raised. *Buds* small, plump, acute, free, slightly pubescent.

FRUIT.

Fruit small or below medium, uniform in size and shape. *Form* roundish ovate or a little inclined to oblong, rather regular; sides usually unequal. *Stem* variable, usually long, slender. *Cavity* small, acuminate to acute, shallow to medium in depth, narrow, slightly symmetrical, sometimes compressed, russeted but slightly if at all. *Calyx* medium to large, leafy, closed

or sometimes a little open; lobes long, rather narrow, acuminate. *Basin* small, shallow, narrow, rather obtuse, smooth or wavy.

Skin thin, tender, smooth, glossy, pale yellow or greenish, nearly over-spread with deep purplish-red approaching blackish-purple on the exposed cheek; some specimens are entirely red. *Dots* rather numerous, very small, inconspicuous, light.

Calyx tube short, moderately wide, conical to funnel-form. *Stamens* median to marginal.

Core large, axile to somewhat abaxile; cells symmetrical, open or sometimes closed; core lines slightly clasping or meeting. *Carpels* broadly ovate to elliptical. *Seeds* rather dark brown, numerous, small to medium, plump, acute.

Flesh white, fine, tender, juicy, brisk subacid, good to very good.

Season late July to early winter.

STRIPED RED JUNE. A variety has found its way into cultivation which appears in all respects identical with the Red June except that the fruit is striped. It is supposed to be either a seedling of the Red June or a sport of that variety (11, 12).

RED TRANSPARENT.

REFERENCES. 1. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:437. 2. Budd, *Ia. Agr. Coll. Bul.*, 1885:13. 3. Van Deman, *U. S. Pom. Rpt.*, 1888:572. fig. 4. Budd, *Ia. Agr. Coll. Bul.*, 1890:18. 5. Bailey, *An. Hort.*, 1892:248. 6. Beach, *N. Y. Sta. An. Rpt.*, 13:584. 1894. 7. Troop, *Ind. Sta. An. Rpt.*, 12:81. 1899. **SYNONYMS.** No. 333 (1, 2, 3, 4). RED TRANSPARENT (2, 3, 4, 5, 6, 7). *Red Transparent* (1). *Skrosnoi krasnoi* (2). SKWOSNOI KRASNOI (1).

A Russian variety of little value where Primate can be grown. Fruit medium size with pale skin nearly covered with red and overspread with delicate bloom. Basin irregularly wrinkled; calyx prominent, closed; flesh greenish-white, not very crisp; water-cores badly; season late July and early August (6).

RED WINE.

REFERENCES. 1. Webster, *Montreal Hort. Soc. Rpt.*, 7:54. 1881. 2. Gibb, *Ib.*, 9:95. 1883. 3. *Ib.*, *Ia. Hort. Soc. Rpt.*, 1883:439. 4. Budd, *Ia. Agr. Coll. Bul.*, 1885:14. 5. Bailey, *An. Hort.*, 1892:248. 6. Taylor, *Mc. Pom. Soc. Rpt.*, 1892:57. 7. Hansen, *S. D. Sta. Bul.*, 76:92. 1902. 8. Budd-Hansen, 1903:163.

SYNONYMS. No. 343 (2, 4, 7). RED WINE (4, 5, 6, 7, 8). *Red Wine* (1, 2, 3). *Rother Weinapfel* (7). *Rotherwein appel* (4). *Vinnoe Krasnoe Oscannee* (4). WEINAPFEL ROTHER (2). WEINAPFEL ROTHER (3). WEINAPFEL ROTHER (1).

A Russian apple of the Lowland Raspberry type (7, 8). Fruit medium, waxen-white almost completely covered with bright red. Flesh white, tender, subacid, good. Season August and September.

So far as we know this variety has not been tested in New York.

REED.

REFERENCE. 1. Downing, 1869:329.

SYNONYMS. None.

A local variety which according to Downing originated with George Reed, Leedsville, Dutchess county, N. Y. Fruit medium, whitish shaded and mottled with light and dark red; flesh white, a little stained next the skin, pleasant subacid, good; season November.

We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

REPKA.

REFERENCES. 1. Barry, 1883:334. 2. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:79. 3. Gibb, *Am. Pom. Soc. Rpt.*, 1887:57. 4. Bailey, *Am. Hort.*, 1892:248. 5. Beach, *N. Y. Sta. An. Rpt.*, 13:584. 1894.

SYNONYMS. No. 139 (2, 3). REPKA (1, 2, 3, 4, 5). *Rieпка* (3).

A medium sized yellow apple of good quality in season in August and early September; inferior to Primate both in appearance and quality. The tree comes into bearing moderately early and is an annual cropper yielding fair to good crops. Not recommended for planting in New York.

This is distinct from either Repka Aport, Repka Sweet or Repka Malenka.

Historical. Origin Russia. Received in 1884 from Ellwanger and Barry, Rochester, N. Y., for testing at this Station. It has been but little disseminated in New York.

TREE.

Tree rather small; not a vigorous grower. *Form* spreading or roundish. *Twigs* short, straight, stout with large terminal buds; internodes short. *Bark* brown or reddish-brown, heavily coated with gray scarf-skin. *Lenticels* scattering, medium to small, round, slightly raised. *Buds* large, prominent, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium or below. *Form* roundish oblate inclined to conic, regular; sides unequal. *Stem* rather short, moderately slender. *Cavity* acute, medium in depth to deep, rather wide, heavily russeted and with outspreading russet rays. *Calyx* rather small, closed. *Basin* moderately deep, wide, somewhat abrupt, usually furrowed or wrinkled.

Skin moderately thin, tough, clear pale yellow or whitish. *Dots* small, white, pale and submerged or russet.

Calyx tube long, narrow, funnel-form.

Core small; cells closed; core lines clasping. *Carpels* broadly roundish. *Seeds* medium size, wide, flat, obtuse.

Flesh white, rather firm, fine, juicy, crisp, rather mild subacid, good.


Season August and September.

RIBSTON.

REFERENCES. 1. Forsyth, 1803:52. 2. Diel, 11:93. 1813. 3. Cox, 1817:125. fig. 4. Forsyth, 1824:124. 5. Buel, *N. Y. Bd. Agr. Mem.*, 1828:476. 6. Fessenden, 1828:130. 7. *Pom. Mag.*, 3:No. 141. 1830. col. pl. 8. *London Hort. Soc. Cat.*, 1831:32. 9. Ronalds, 1831:54. 10. Kenrick, 1832:52. 11. Floy-Lindley, 1833:59. 12. Manning, 1838:54. 13. Manning, *Mag. Hort.*, 7:49. 1841. 14. Russell, *ib.*, 10:403. 1844. 15. Downing, 1845:131. 16. Downing, *Horticulturist*, 2:416. 1847. 17. *ib.*, 3:421. 1848. 18. Thomas, 1849:152. 19. Cole, 1849:126. 20. *Horticulturist*, 6:16, 292. 1851. 21. Emmons, *Nat. Hist. N. Y.*, 3:75. 1851. 22. Elliott, 1854:155. 23. *Am. Pom. Soc. Cat.*, 1854. 24. *Mag. Hort.*, 21:62. 1855. 25. Hooper, 1857:80. 26. Oberdieck, *Ill. Handb. Obst.*, 1:353. 1858. 27. *Mag. Hort.*, 26:116. 1860. 28. Warder, 1867:612. 29. Regel, 1868:463. 30. Berghuis, 1868:No. 3. col. pl. 31. Mas, *LeVerger*, 1868:99. 32. Leroy, 1873:750. figs. 33. Lauche, 1:No. 25. 1882. col. pl. 34. Barry, 1883:353. 35. Hogg, 1884:194. 36. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 37. Hick, *Can. Hort.*, 15:157. 1892. 38. Bailey, *Am. Hort.*, 1892:248. 39. Bredsted, 1893:137. 40. Gaucher, 1894:No. 19. col. pl. 41. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:355. 42. *Can. Hort.*, 22:510. 1899. 43. Eneroth-Smirnoff, 1901:224. 44. Waugh, *Rural N. Y.*, 61:285, 286. 1902. fig. 45. Budd-Hansen, 1903:165. fig. 46. Beach and Clark, *N. Y. Sta. Bul.*, 248:141. 1904.

SYNONYMS. *Beautiful Pippin* (25). *Englische Granat-Reinette* (40). *Essex Pippin* (40). *Formosa* (25). *Formosa Pippin* (7, 10, 15, 22, 32, 40). *Glory of York* (10, 15, 22, 25, 32, 35, 40, of some 7). *Granat-Reinette* (40). *Nonpareille* (40). *Pepin Ribston* (40). *Reinette Grenade Anglaise* (40). *Reinette de Traver* (32). RIBSTON (2, 8, 9, 11, 20, 23, 24, 26, 27, 29, 30, 31, 33, 36, 37, 39, 42, 43, 44, 45). RIBSTON PIPPIN (1, 3, 4, 5, 7, 10, 13, 14, 15, 16, 17, 18, 19, 21, 22, 25, 28, 32, 34, 35, 38, 41, 44). *Ribston Pippin* (45, 46). RIBSTON PEPPING (40). *Ribston's Pepping* (40). *Ribstone* (32). RIBSTONE PIPPIN (6, 12). RIDGE (46 by error). *Rockhill's Russet* (32). *Travers* (15, 22, 25, 32). *Travers Apple* (7, 10, 40). *Travers Peppin* (40). *Travers Pippin* (35). *Travers Reinette* (40).

Ribston evidently belongs in the same group as Hubbardston. It is much esteemed for its rich flavor and fine quality and it is desirable either for dessert or culinary uses. The fruit is pretty smooth and uniform but often it averages below medium size and is ordinary in appearance. Heat ripens it quickly and it is not considered a very good keeper. In cold storage, if properly handled before storing, it is possibly equal to Tompkins King or Hubbardston as a keeper (46). Its season in Southern New York extends from late September to November or December, and in the northern and more elevated regions from late fall to early or mid-winter and sometimes a portion of the fruit may be kept



RIBSTON

till spring in ordinary storage. The tree is pretty hardy, vigorous, healthy and long-lived. It comes into bearing rather young and usually bears some fruit every year. Occasionally the crops are heavy but more often they vary from moderate to rather light. Generally speaking, it is hardly satisfactory as a cropper and Hubbardston is much to be preferred for planting in commercial orchards in New York.

Historical. Ribston originated more than two hundred years ago in Yorkshire, England (1, 7). In that country it has long been considered the standard of excellence among dessert apples. It has long been known in cultivation in America but has not gained the standing here that it holds in England. It is not grown to any considerable extent in New York but succeeds better farther north, as in portions of Northern New England and of Canada, where it is of some commercial importance (16, 20, 24, 44).

TREE.

Tree medium in size or sometimes rather large, moderately vigorous to vigorous with rather stout, stocky branches. *Form* rather upright and spreading or roundish, not very regular. *Twigs* medium to rather long, rather slender to moderately stout; internodes medium to long. *Bark* bright dark reddish-brown and olive-green, somewhat mottled with grayish scarf-skin. *Lenticels* conspicuous, scattering, small to medium, elongated or roundish. *Buds* medium to large, broad, plump, obtuse, nearly free, very pubescent. *Foliage* rather dense; leaves broad.

FRUIT.

Fruit medium or above, pretty uniform in shape and size. *Form* roundish, rather broad and flattened at the base, narrowing somewhat toward the basin, occasionally a little inclined to roundish oblong, often broadly and obscurely ribbed. *Stem* pubescent, medium to short, occasionally moderately slender, more often rather thick, sometimes irregularly swollen or inserted under a lip. *Cavity* rather large, acute, moderately shallow to rather deep, wide to moderately narrow, sometimes furrowed or compressed, occasionally smooth and green but often faintly russeted and with some outspreading russet. *Calyx* variable, small to rather large, closed or partly open; lobes sometimes separated at the base, erect or converging, tips usually somewhat reflexed. *Basin* small to medium, shallow to moderately deep, moderately narrow, more or less abrupt or occasionally obtuse, often slightly furrowed and wrinkled.

Skin smooth or slightly roughened with russet, deep yellow or greenish-yellow more or less overspread with rather dull red which in highly colored specimens deepens to a distinct red with some obscure carmine stripes and splashes. *Dots* scattering, conspicuous toward the base, more numerous and smaller toward the basin, pale, sometimes whitish, often areolar with russet center. *Prevailing effect* sometimes rather attractive but more often the colors are rather dull.

Calyx tube rather wide, cone-shape or sometimes funnel-form. *Stamens* basal.

Core below medium to small, axile or with a narrow hollow cylinder at the axis; cells pretty regular, closed; core lines clasping to nearly meeting. *Carpels* roundish to nearly elliptical, emarginate, slightly tufted. *Seeds* variable, some abortive, usually but few are plump, light and dark brown, rather large, moderately narrow to wide, medium to long, obtuse or sometimes approaching acute, sometimes slightly tufted.

Flesh tinged with yellow, firm, very crisp, medium in texture, juicy, pleasantly aromatic, rich, sprightly subacid, very good.

Season late September to December or later.

RICHARD GRAFT.

REFERENCES. 1. *Mag. Hort.*, 18:492. 1852. 2. Downing, 1857:101. fig. 3. Warder, 1867:457. fig. 4. Thomas, 1875:204. 5. *Am. Pom. Soc. Cat.*, 1877:14. 6. Barry, 1883:340. 7. *Mich. Hort. Soc. Rpt.*, 1888. 8. Hendricks, *Rural N. Y.*, 47:759, 811. 1888. 9. Bailey, *Am. Hort.*, 1892:248. 10. Budd-Hansen, 1903:166.

SYNONYMS. *Derrick and Ann* (8). *Derrick's Graft* (2, 4). *Red Spitzenberg* (3). *Red Spitzenburgh* (2, 4, 6). *RICHARD* (1). *RICHARD GRAFT* (2, 3, 4, 5, 6, 7, 8, 9). *Strawberry* (2, 3). *Wine* (2, 3).

This is a very fine fall apple of superior dessert quality. It begins to ripen during late August or early September; the crop ripens in succession during a period of several weeks, and some portion of the fruit may be kept till late autumn. Several pickings are required in order to secure the fruit in prime condition. The tree is upright, of medium size, moderately vigorous, long-lived and a reliable cropper yielding good crops biennially. It is an excellent variety for home use and is being grown to a limited extent in commercial orchards with profit.

Historical. This variety was originated at Greenport, Columbia county, N. Y., by Richard Delamatter. It was introduced about 1860 by E. G. Studley, a nurseryman of Claverack, Columbia county, N. Y. Its cultivation is being extended somewhat in Columbia county, but as yet it is but little known outside of the Hudson valley.

TREE.

Tree of medium size, moderately vigorous. *Form* upright or roundish, open. *Twigs* moderately long, curved, moderately stout; internodes medium. *Bark* dark brown, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, medium size, round, not raised. *Buds* medium size, broad, acute to obtuse, free, pubescent.

FRUIT (2, 8).

Fruit medium size. *Form* roundish oblate. *Stem* of medium length, slender. *Cavity* large. *Calyx* small, closed. *Basin* medium size.

Skin yellow, nearly covered with stripes and splashes of deep red.

Flesh yellowish, very tender, juicy, aromatic, subacid, very good.

Season September.

ROLFE.

REFERENCES. 1. Downing, 1857:167. 2. Warder, 1867:725. 3. Thomas, 1875:505. 4. *Me. Pom. Soc. Rpt.*, 1876:149. 5. *Ib.*, 1885:135, 138. 6. *Ib.*, 1888:120. *fig.* 7. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:294. 8. Bailey, *Am. Hort.*, 1892:248. 9. Munson, *Me. Sta. Rpt.*, 1893:133. 10. *Ib.*, 1896:71, 81. 11. *Am. Pom. Soc. Cat.*, 1897:14. 12. Munson, *Me. Sta. Rpt.*, 1902:90, 92. 13. *Me. Pom. Soc. Rpt.*, 1902:49, 50. 14. Budd-Hansen, 1903:166.

SYNONYMS. MACOMBER (1, 2, 3, 7). *Macomber* (4, 9, 14). ROLFE (4, 5, 6, 8, 9, 10, 11, 12, 13, 14).

Fruit medium to large, of good quality and rather attractive for a yellowish apple. It is in season from late September to December or January. The tree is very hardy, vigorous and a reliable cropper. At the present time it is probably grown more extensively in Maine than in any other section of the country. It is there regarded highly wherever it is known and is gaining in popularity among fruit growers (1, 6, 10, 12). It is worthy of testing in those portions of the state where superior hardiness in a variety is a matter of prime importance.

Historical. Originated in the town of Guilford, Maine, about 1820. Said to be a seedling of the Blue Pearmain. It has, as yet, been disseminated but sparingly in this state and is but little known among New York fruit growers.

TREE.

Tree rather large, vigorous. *Form* roundish to spreading and drooping, rather dense; laterals slender, willowy. *Twigs* rather long, irregularly crooked, slender to moderately stout; internodes long to below medium. *Bark* brown to reddish-brown with an occasional tinge of olive-green, overlaid with scarf-skin, pubescent near tips. *Lenticels* moderately numerous, of a dull color but rather conspicuous, medium or above, roundish, not raised. *Buds* very deeply set in bark, medium in size, broad, flat, obtuse, appressed, pubescent.

FRUIT.

Fruit medium to sometimes large, pretty uniform in shape and size. *Form* roundish to roundish oblate, regular or somewhat angular, symmetrical. *Stem* short to medium and rather slender. *Cavity* acute to slightly acuminate, moderately deep, rather wide, sometimes slightly furrowed or compressed. *Calyx* small to above medium, closed; lobes short to moderately long, rather wide, acute. *Basin* below medium to rather large, pretty regular, shallow to moderately deep, narrow to rather wide, a little abrupt, slightly wrinkled.

Skin moderately thin, rather tough, glossy, clear pale yellow, sometimes faintly blushed or in well colored specimens distinctly shaded and striped with lively red. *Dots* numerous, inconspicuous, small. *Prevailing effect* yellow or yellow and red.

Calyx tube rather narrow, short, funnel-shape. *Stamens* median.

Core variable, below medium to large, abaxile; cells usually symmetrical, wide open; core lines clasping. *Carpels* often markedly concave, broadly ovate, emarginate, tufted. *Seeds* above medium, rather long and narrow, plump, acute or approaching acuminate, light brown.

Flesh whitish with slight tinge of yellow, moderately fine-grained, crisp, tender, juicy, briskly subacid, good.

Season late September to December or January.

ROMAN STEM.

REFERENCES. 1. Cox, 1817:132. fig. 2. Thacher, 1822:135. 3. *London Hort. Soc. Cat.*, 1831:33. 4. Downing, 1845:131. 5. Kirtland, *Horticulturist*, 2:545. 1848. 6. Thomas, 1849:185. 7. Phoenix, *Horticulturist*, 4:472. 1850. 8. Emmons, *Nat. Hist. N. Y.*, 3:86. 1851. 9. Elliott, 1854:155. 10. *Horticulturist*, 10:87. 1855. 11. Hooper, 1857:81. 12. *Mag. Hort.*, 24:110. 1858. 13. *Am. Pom. Soc. Cat.*, 1862. 14. Warder, 1867:579. fig. 15. Barry, 1883:354. 16. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 17. Bailey, *Am. Hort.*, 1892:248. 18. Hansen, *S. D. Sta. Bul.*, 76:94. 1902. 19. Budd-Hansen, 1903:166. 20. Beach and Clark, *N. Y. Sta. Bul.*, 248:142. 1904.

SYNONYMS. *French Pippin* of some (9). ROMAN STEM (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20).

Fruit about medium size, whitish-yellow, often somewhat blushed. The flesh is juicy, aromatic, subacid and very good in quality, particularly for dessert use. It is in season from midautumn to midwinter. "A good fruit but in a great measure superseded by other sorts" (15). The tree is moderately vigorous, spreading, irregular, very hardy and very productive. In the trying climate of the upper Mississippi valley it has proved hardier than most of the old varieties from the East and has succeeded well where the varieties of the grade of hardness of Baldwin and Rhode Island *Greening* have failed. It is not well suited for commercial purposes because it is yellow, lacks good size and is not a late keeper.

Historical. Cox published the following description of Roman Stem in 1817: "This apple was first propagated in the neighbourhood of Burlington, New-Jersey, where the original tree is now standing. It is an excellent early winter fruit, much admired for its tender, mild, juicy and agreeable properties; the size is small, the form round, the stalk of singular appearance, from a fleshy protuberance of the neighboring part, resembling an aquiline nose, whence the apple derives its name—the skin is rough, the color yellow, with black clouds and spots—the tree is of handsome and vigorous growth, with long shoots, and great fruitfulness; it is in every respect deserving of extensive cultivation."

Roman Stem has been pretty widely disseminated and considerably cultivated in various parts of the Southern, Central and Western states but it is now generally superseded by other kinds. It is but little grown in New York.

ROMNA.

REFERENCES. 1. Budd, *Am. Pom. Soc. Rpt.*, 1883:73. 2. *Ib.*, *Id. Agr. Coll. Bul.*, 1883:29. 3. *Ib.*, 1885:20, 23, 29. 4. Gibb, *Am. Pom. Soc. Rpt.*, 1887:50. 5. Lyon, *U. S. Pom. Bul.*, 2:45. 1888. 6. Budd, *Id. Agr. Coll. Bul.*, 1890:24. 7. *Ib.*, 1892:7. 8. *Ib.*, *Id. Sta. Bul.*, 18:520. 1892. 9. *Ib.*, 19:541. 1892. 10. Troop, *Ind. Sta. Bul.*, 53:124. 1894. 11. Budd, *Id. Sta. Bul.*, 31:333. 1895. 12. Beach, Paddock and Close, *N. Y. Sta. An. Rpt.*, 15:275. 1896. 13. Thomas, 1897:295. *fig.* 14. Waugh, *Id. Sta. Bul.*, 61:31. 1897. *fig.* 15. Budd, *Id. Sta. Bul.*, 41:69, 70, 80. 1899. 16. Troop, *Ind. Sta. Rpt.*, 1899:81. 17. Hansen, *S. D. Sta. Bul.*, 76:94. 1902. *fig.* 18. Budd-Hansen, 1903:168. 19. Beach and Clark, *N. Y. Sta. Bul.*, 248:142. 1904.

SYNONYMS. *No. 11 M* (1, 2, 6, 7, 9). *No. 599 Dept.* (1, 2, 3, 4, 6, 7, 8, 9, 17, 18). *Omensk* (4). *ROMENSKOE* (2, 4, 5, 8). *Romenskoe* (1, 3, 17, 18). *ROMNA* (1, 3, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19). *Romnskoe* (3, 4, 5).

A Russian variety received from Dr. T. H. Hoskins, Newport, Vermont, in 1888 for testing at this Station. It was described in 1896 (12) as being in season that year during the last of August and the first of September. This statement was erroneous because it was incomplete. While the fruit began to come in season during the last of August and the first of September some portion of it was kept in ordinary storage till midwinter. In a subsequent report (19) it was correctly stated that as fruited at this Station the commercial limit of this variety is early October and its season in ordinary storage extends from September to January. It is properly classed as a fall and early winter apple here. The tree is vigorous, hardy, comes into bearing rather young and yields good crops biennially. It does not appear to be worthy of the attention of fruit growers in New York except possibly where superior hardiness is a prime requisite. The fruit corresponds very closely with the illustrated description given by Troop (10, 16) and Waugh (14) but it varies considerably from the descriptions of Budd and Hansen (3, 6, 9, 11, 15, 17, 18) particularly in that it is usually oblate conic and is in season during the autumn and early winter instead of late winter and spring.

The following is one of Budd's descriptions of Romna (9, 11). "This succeeds best on dry soil where its roots run very deep. Fruit medium in size, conical, smooth, handsomely colored. Flesh white, firm, quite acid and best for cooking, but when matured it is much better for dessert use than Willow or Missouri Pippin or other coarse sorts found in our markets. Season, midwinter here, and late winter north of 43d parallel."

TREE.

Tree moderately vigorous; branches short, stout, curved, crooked and drooping. *Form* spreading, drooping, flat. *Twigs* short to medium, straight, moderately slender to stout, with large terminal buds; internodes medium to long. *Bark* brown, somewhat tinged with red, streaked with grayish scarf-skin; slightly pubescent near tips. *Lenticels* scattering,

small to medium, roundish or oval, slightly raised. *Buds* moderately small to very large and prominent, broad, very plump, acute, free or nearly so, scarcely pubescent.

FRUIT.

Fruit medium or sometimes rather large, not very uniform in shape or size. *Form* usually oblate conic, irregularly elliptical or broadly and obscurely angular, often unsymmetrical with sides unequal. *Stem* medium length to short and stout, pubescent. *Cavity* medium to rather large, acute or sometimes nearly obtuse, medium in depth to rather deep, rather wide to narrow, somewhat furrowed, occasionally lipped, russeted and often with conspicuous outspreading russet. *Calyx* above medium to large, open or partly closed; lobes often separated at the base, medium in length, rather broad, acute. *Basin* medium to large, often oblique, moderately narrow to wide, sometimes compressed, abrupt, furrowed and wrinkled.

Skin thick, tough, smooth or partly roughened with flecks of russet; color greenish becoming yellow more or less blushed and overspread with thin bloom. Well colored specimens are covered to a considerable extent with pinkish-red, blushed and striped with bright carmine. *Dots* small, numerous, pale yellow or grayish, sometimes rather conspicuous.

Calyx tube rather long, wide, funnel-shape. *Stamens* median.

Core medium to small, axile or nearly so; cells closed; core lines meeting or slightly clasping. *Carpels* roundish ovate, somewhat emarginate, slightly tufted. *Seeds* medium to small, rather short, narrow, plump, obtuse to acute, rather dark brown.

Flesh yellowish, firm, moderately coarse, juicy, briskly subacid, slightly astringent, fair to good.

Season September to January (19).

RONK.

REFERENCES. 1. *Rural N. Y.*, 48:279. 1889. fig. 2. Lyon, *Mich. Sta. Bul.*, 143:201. 1897. 3. Farrand, *Ib.*, 205:46. 1903. 4. Beach and Clark, *N. Y. Sta. Bul.*, 248:142. 1904.

SYNONYMS. None.

Fruit of the Vandevere type, medium or above, rather dull red, pleasant subacid, good; season October to late winter. Commercial limit in ordinary storage January (4). The tree comes into bearing rather young and is moderately productive. It has not been sufficiently tested to determine its value for this state.

Historical. Originated about 1860 with Mr. Ronk, Boone county, Indiana. It is supposed to be a seedling of Vandevere which it much resembles (1).

ROSE RED.

REFERENCES. 1. *Rural N. Y.*, 1871 (cited by 5). 2. Downing, 1872:30 app. fig. 3. Thomas, 1875:511. 4. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 5. Ragan, *U. S. P. B. I. Bul.*, 56:267. 1905.

SYNONYMS. *Autumn Rose* (4). ROSE RED (1, 2, 3, 4, 5).

A variety of unknown origin. It is supposed to have originated in Egypt, Monroe county, N. Y. According to Downing (2) the tree is thrifty, a reliable cropper and very productive; the fruit medium, roundish oblate, whitish, striped and splashed with light and dark red; flesh yellowish, very tender, lively subacid, very good; in season during late September, October and November.

We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

SAFSTAHOLMS.

REFERENCES. 1. Regel, 1868:473. 2. Gibb, *Montreal Hort. Soc. Rpt.*, 1886 87:81. 3. Eneroth-Smirnoff, 1901:46. 4. Hansen, *S. D. Sta. Bul.*, 76:96. 1902.

SYNONYMS. SAFSTAHOLM (2, 4). SÄFSTAHOLMSÄPPE (1). SÄFSTAHOLMS-ÄPPE (3).

This is an apple of fairly good red color, not particularly bright yet not unattractive. The flesh lacks piquancy and is not very juicy but because of its distinct aroma and rich subacid flavor it is classed among the good dessert apples. It is hardly acid enough for culinary use. The tree is a pretty good grower, comes into bearing young and so far as tested at this Station promises to be productive. It is doubtful whether it has sufficient value for the New York fruit grower to make it worthy of trial in this state.

Historical. Originated in Sweden about 1835. It was received for testing at this Station from the United States Pomologist in 1901.

TREE.

Tree moderately vigorous with rather short, slender branches. *Form* upright spreading or roundish, open. *Twigs* short, rather slender to moderately stout, straight or nearly so; internodes short to medium. *Bark* rather dull brown tinged with red, mottled with heavy scarf-skin, pubescent. *Lenticels* very scattering, small to medium, roundish, not raised. *Buds* often rather deeply set in the bark, medium size or below, plump, obtuse to somewhat acute, free, pubescent.

FRUIT.

Fruit medium to large, pretty uniform in shape and size. *Form* oblong to oblong conic, somewhat elliptical, often indistinctly ribbed; sides often unequal. *Stem* short to medium, moderately slender to rather thick. *Cavity* large, acute to acuminate, moderately shallow to deep, wide, sometimes lipped, often russeted. *Calyx* medium or below, usually partly open; lobes moderately narrow, acute. *Basin* small to medium, obtuse to rather abrupt, shallow to moderately deep, moderately narrow to rather wide, gently furrowed.

Skin thin, smooth or sometimes slightly rough toward the apex, pale yellow mottled and blushed with red, becoming rather dull red over a considerable portion of the fruit, mottled and splashed with carmine and sometimes marked with flecks and irregular lines of russet. *Dots* rather conspicuous, often large, pale gray or with russet center. *Prevailing color* red.

Calyx tube cone-shape. *Stamens* basal.

Core medium to large, abaxile; cells often unsymmetrical, usually somewhat open; core lines meeting. *Carpels* roundish to broadly ovate, mucronate, sometimes emarginate, tufted. *Seeds* numerous, above medium to rather large, wide, usually obtuse, plump.

Flesh yellowish, firm, a little coarse, moderately juicy, peculiarly aromatic, mildly subacid, rich, good for dessert, rather mild for culinary use.

Season late October or November to midwinter; often some portion of the fruit may be kept in ordinary storage to March.

SAILEE RUSSET.

REFERENCE. 1. Waugh, *Vt. Sta. An. Rpt.*, 14:307. 1901.

SYNONYMS. None.

A local variety grown in the vicinity of Lake Champlain. The following account of it is given by Waugh (1):

"Sailee was a Frenchman who came over from France about a hundred years ago and who had a farm on Cumberland Head, Clinton county, N. Y., just across from Grand Isle. He had a large orchard and grew many varieties of apples, some of which he had brought from France, others of which came from other sources, and some of which he originated himself. From his having given his own name to this variety it is supposed to have originated in his own grounds. It was early distributed to Grand Isle, and may be found in several of the older and more complete collections. It is a good variety, but not superior to Roxbury.

"Fruit oblate, slightly conic, size small to medium, cavity very deep and broad, stem medium long, slender, basin deep, corrugated, calyx small, closed, color dull green with occasional blush and considerable russet, dots russet, skin tough, flesh white, core small, flavor subacid, quality good, season early winter."

SALLY AUTUMN.

REFERENCES. 1. Downing, 1857:187. 2. Thomas, 1875:511.

SYNONYMS. None.

A local variety which originated at Plattsburg, N. Y. Fruit medium, roundish conic, greenish-yellow frequently with a deep red cheek. Stalk short; cavity medium; calyx small, closed; basin small, narrow; flesh very tender, rich, aromatic, subacid, good. September (1, 2).

We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

ST. LAWRENCE.

REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 1187. 2. Kenrick, 1832:28. 3. *Mag. Hort.*, 1:149. 1835. 4. Hovey, *Ib.*, 13:539. 1847. 5. *Mag. Hort.*, 14:531, 539. 1848. 6. Thomas, *Cultivator*, 5:246. 1848. 7. Cole, 1849:104. 8. Thomas, 1849:152. 9. Barry, 1851:286. 10. Waring, 1851:30. 11. Elliott, 1854:158. 12. Downing, 1857:193. 13. Hooper,

1857:90. 14. Gregg, 1857:47. 15. *Am. Pom. Soc. Cat.*, 1862. 16. Warder, 1867:731. 17. Downing, 1872:10 index, app. 18. Leroy, 1873:799. fig. 19. *Montreal Hort. Soc. Rpt.*, 1876:11. 20. *Ib.*, 15:19, 27. 1890. 21. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 22. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57. 23. Bailey, *Am. Hort.*, 1892:249. 24. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 6:9. 1899. figs. 25. Waugh, *Ft. Sta. An. Rpt.*, 14:307. 1901. 26. Hansen, *S. D. Sta. Bul.*, 76:96. 1902. 27. Farrand, *Mich. Sta. Bul.*, 205:46. 1903. 28. Budd-Hansen, 1903:171. fig. 29. *Can. Hort.*, 27:51. 1904. fig. 30. Beach and Clark, *N. Y. Sta. Bul.*, 248:143. 1904. 31. Scriver, *Can. Hort.*, 28:277. 1905.

SYNONYMS. *Corse's St. Lawrence* (8). *Montreal* (4, 11, 18). *SAINT-LAURENT* (18). *ST. LAWRENCE* (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31). *Saint-Lawrence* (18). *York and Lancaster* (17).

When well grown, St. Lawrence is a large, handsome apple. It is better for dessert than for culinary use but does not excel standard varieties of its season for either purpose. While it does very well in some portions of Western New York, generally speaking, it reaches a higher degree of perfection in favorable locations in the St. Lawrence valley and in the Lake Champlain region than in other portions of the state. The crop ripens somewhat unevenly and should have more than one picking in order to secure the fruit in prime condition and prevent great loss from dropping. It does not stand heat well before going into storage and goes down quickly. The fruit may not remain on the tree till it is well colored, and unless it is well colored it fades in the barrel so much as to render it almost valueless for market. It varies greatly in keeping qualities in different seasons but usually October is its commercial limit in ordinary storage. In cold storage it may be held until December (30). The tree is a moderately strong grower, hardy, generally pretty healthy, moderately long-lived and a reliable cropper yielding good to rather heavy crops biennially. It is not a very good grower in the nursery. Some growers hold that it is desirable to topwork it upon some more vigorous stock as Northern Spy. Although many fruit growers regard it as a fairly profitable commercial apple it cannot be recommended for general cultivation.

Waugb remarks, that in Grand Isle county, Vermont, "It is rather common but not highly prized. It precedes Fameuse in season and is of the same general character" (25). Woolverton (29) states that it is not planted in the commercial orchards of Ontario bordering Lakes Ontario, Erie or Huron, but it is valued in orchards along the St. Lawrence river and parts of the Province between the latitudes 45 and 46. In the Niagara district it is considerably affected by scab and by codling moth.

Historical. As early as 1835 St. Lawrence was recommended as one of the American varieties which was worthy of cultivation in England (3). Its origin does not appear to be definitely known but some credit it to this country (12, 17, 26, 28, 29). In 1848 Thomas (6) described it as a newly introduced variety cultivated in the vicinity of Rochester and originally from Lower Canada. In 1862 it was entered in the catalogue of the American Pomological Society (15). It is frequently listed by nurserymen (23) but is now seldom planted in New York state.

TREE.

Tree medium size, moderately vigorous. *Form* upright spreading. *Twigs* smooth, rather dark reddish-brown.

FRUIT.

Fruit large to medium. *Form* oblate inclined to conic varying to roundish oblate, faintly ribbed. *Stem* short to medium in length, moderately slender. *Cavity* large, acute, deep, regular, greenish-russeted. *Calyx* small, closed. *Hasin* rather small, narrow, moderately deep, abrupt, wrinkled.

Skin pale yellow washed and marbled with bright red striped and splashed with bright dark carmine and overspread with thin white bloom. *Dots* numerous, rather obscure, fine, russet.

Calyx tube narrow, cone-shape to funnel-form. *Stamens* median.

Core medium size, somewhat abaxile; cells partly open; core lines clasping. *Carpels* obovate to elliptical, emarginate.

Flesh white, sometimes slightly stained with red, tender, fine-grained, crisp, juicy, mild subacid, good to very good for dessert; rather mild for culinary uses.

Season September and October.

ST. PETER.

REFERENCES. 1. Hoskins, *Ia. Hort. Soc. Rpt.*, 1879:414. 2. Webster, *Am. Pom. Soc. Rpt.*, 1883:113. 3. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:440. 4. Hoskins, *Rural N. Y.*, 45:673. 1886. figs. 5. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886-87:75. 6. *N. Y. Sta. An. Rpt.*, 7:91. 1888. 7. Bailey, *An. Hort.*, 1892:249. 8. Beach, *N. Y. Sta. An. Rpt.*, 13:591. 1894. 9. Ragan, *U. S. B. P. I. Bul.*, 56:231, 273. 1905.

SYNONYMS. *No. 80* (5, 9). *No. 372* (9). PETROVSKOE (5, 9). PETROWSKOE (2, 3). *Petrowskoe* (9). ST. PETER (1, 4, 6, 7, 8). *St. Peter* (2, 3, 9).

A small August apple, greenish-yellow streaked and splashed with dull carmine, mild subacid, fair quality. The tree is a moderately vigorous grower, rather slow in coming into bearing and not a reliable cropper. Not valuable for planting in New York.

Historical. A Russian apple imported by the United States Department of Agriculture in 1870. It was received in 1888 from Dr. T. H. Hoskins, Newport, Vt., for testing at this Station (6).

SANDY GLASS.

REFERENCES. 1. Budd, *Ia. Agr. Coll. Bul.*, 1885:23, 30. 2. Schroeder, *Montreal Hort. Soc. Rpt.*, 1886:72. 3. Budd, *Ia. Agr. Coll. Bul.*, 1890:22. 4. Munson, *Me. Sta. Rpt.*, 1896:76. 5. Budd, *Ia. Sta. Bul.*, 19:538. 1896. 6. Stinson, *Ark. Sta. Bul.*, 43:105. 1896. 7. Thomas, 1897:295. fig. 8. Munson, *Me. Sta. An. Rpt.*, 18:84. 1902. 9. Hansen, *S. D. Sta. Bul.*, 76:96. 1902.

SYNONYMS. *No. 24 M* (1, 2, 3, 5, 8, 9). SANDY GLASS (1, 2, 3, 4, 5, 6, 7, 8, 9). *Steklianka pesotchnaya* (1).

A rather attractive apple of greenish-yellow color and often faintly blushed; it is of pretty good quality but inferior to Fall Pippin and other standard varieties of its season. The tree is below medium size, not a strong grower, comes into bearing rather young and is a reliable cropper yielding pretty good crops nearly annually. It is not valuable for planting in New York except possibly in localities where superior hardiness is specially desired.

Historical. Imported from Russia by the Iowa Agricultural College (1), from which institution it was received in 1890 for testing at this Station.

TREE.

Tree below medium size, moderately vigorous. *Form* rather flat, spreading and somewhat inclined to droop. *Twigs* moderately long, curved, stout; internodes medium. *Bark* dull brown, heavily coated with rough gray scarf-skin; slightly pubescent near tips. *Lenticels* very conspicuous, numerous, large to medium, oval, raised. *Buds* medium size, broad, plump, acute to obtuse, free, slightly pubescent.

FRUIT.

Fruit above medium to large, quite uniform in size and shape. *Form* roundish oblate or inclined to ovate, regular, sometimes obscurely ribbed. *Stem* short to medium length, rather thick. *Cavity* rather small, acuminate to acute, moderately deep, narrow to medium in width, usually symmetrical, more or less russeted and often with outspreading rays of thin russet. *Calyx* small, closed; lobes medium to short, acute. *Basin* shallow to moderately deep, medium in width to wide, lightly furrowed, wrinkled.

Skin pale greenish-yellow often becoming clear yellow as it ripens, faintly blushed and overspread with whitish bloom. *Dots* numerous, light, small, submerged, mingled with a few that are large and russet.

Calyx tube very long, moderately wide, conical to cylindrical. *Stamens* median to marginal.

Core medium to small, axile; cells symmetrical, closed or slightly open; core lines meeting or clasping. *Carpels* roundish or somewhat ovate, deeply emarginate. *Seeds* large, wide, plump, acute to obtuse, dull dark brown.

Flesh white or with greenish tinge, rather fine, tender, juicy, brisk subacid, fair to good.

Season September to early winter.

SAXTON.

REFERENCES. 1. *Prairie Farmer*, 1860 (cited by 9). 2. *Am. Pom. Soc. Cat.*, 1871:8. 3. Downing, 1872:11 app. 4. Thomas, 1875:511. 5. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-7:94. 6. Bailey, *Am. Hort.*, 1892:239. 7. *Id.*, 1892:249. 8. Burrill and McCluer, *Ill. Sta. Bul.*, 45:321. 1896. 9. Ragan, *U. S. B. P. I. Bul.*, 56:107. 1905. 10. *Id.*, 56:275. 1905.

SYNONYMS. FALL STRIPE (3, 7, 3, 9). *Fall Stripe* (4, 5, 10). SEXTON (1, 2, 4, 5, 6, 10). *Saxton* (3, 7, 9).

An old New England variety (3). Fruit yellow, shaded and splashed with light and dark red; flesh a little coarse, subacid, good to very good; season September. It was put upon the list of the American Pomological Society in 1871 (2) and dropped from that list in 1897. It is still listed by some nurserymen (6) but so far as we can learn it is practically unknown among New York fruit growers.

SCARLET PIPPIN.

REFERENCES. 1. Jones, *Ont. Fr. Gr. Assn. An. Rpt.*, 27:13. 1895. 2. Craig, *Can. Hort.*, 19:381. 1896. fig. 3. *Id.*, 19:117. 1896. fig. 4. *Id.*, *Amer. Gard.*, 20:27. 1899. figs. 5. Waugh, *Vt. Sta. Bul.*, 83:91. 1900. 6. Macoun, *Can. Dept. Agr. Bul.*, 37:41. 1901. 7. Abbott, *Can. Hort.*, 24:18, 123. 1901. 8. Budd-Hansen, 1903:172.

SYNONYMS. CRIMSON BEAUTY (3). *Crimson Pippin* (2). *Crimson Scarlet Pippin* (2). *Leeds Beauty* (4, 5, 6, 8). SCARLET PIPPIN (1, 2, 4, 5, 6, 7, 8).

An apple of the Fameuse group which quite closely resembles McIntosh (2), but is firmer in flesh and slightly more acid. "A very attractive looking apple said to sell better than Fameuse, which it does not equal in quality. Tree a strong, upright grower and said to be a heavy bearer" (6). Season about the same as Wealthy or earlier (1). It appears to be worthy of testing in New York especially in those regions of the state where Fameuse and McIntosh succeed best.

Historical. Originated about 1860 at Lynn, Leeds county, Ontario, near Brockville, where it has been locally grown for some years (4, 7). Mr. Harold Jones, Maitland, Ont., Experimenter for Ontario for apples in the

St. Lawrence river district, has had most to do with bringing this variety to notice as an autumn dessert fruit of value (6), but the report that the variety originated with him is incorrect (7).

TREE.

Tree vigorous. *Form* upright. *Twigs* long, straight, stout; internodes short. *Bark* dark brown or reddish-brown, lightly streaked with scarf-skin, pubescent near tips. *Lenticels* numerous, very conspicuous, medium size, oval, slightly raised. *Buds* medium size, flat, obtuse, appressed, pubescent.

FRUIT (4, 6).

Fruit medium size. *Form* roundish inclined to oblate, regular. *Stem* short, stout to slender. *Cavity* acute, shallow to deep, moderately wide to wide, sometimes lipped. *Calyx* closed or open. *Basin* narrow, shallow, slightly wrinkled or almost wanting.

Core small.

Flesh white, firm, crisp, tender, melting, juicy, mild subacid with a pleasant but not high flavor, very good.

Season fall and early winter.

SCHUYLER SWEET.

REFERENCES. 1. Thomas, *Am. Pom. Soc. Rpt.*, 1871:49. 2. *Rural N. Y.*, 1871:108. 3. Downing, 1872:31 app.

SYNONYMS. None.

This variety is unknown to us. We have received no report concerning it from any of our correspondents. Thomas gave the following description of it in 1871 (1): "A large, showy apple, ripening in October, originated on the lands of Rensselaer Schuyler, Seneca Falls, N. Y. Tree in vigor and form resembles the Baldwin, and is productive.

"Fruit large, roundish, inclining to roundish oblate; pale yellow with a few scattering brown dots; stalk slender, inserted in a large deep cavity; calyx closed; basin large, deep, slightly corrugated; flesh whitish, half fine, tender, moderately juicy, pleasant, sweet; quality good to very good; core small."

SCOLLOP GILLIFLOWER.

REFERENCES. 1. Kenrick, 1835:73. 2. Elliott, 1854:156. fig. 3. Watts, *Horticulturist*, 10:98. 1855. 4. Hooper, 1857:83. 5. Warder, 1867:543. 6. Downing, 1869:348. 7. Downing, *Am. Pom. Soc. Rpt.*, 1875:68. 8. Thomas, 1875:511. 9. Downing, 1876:69, 70, app.

SYNONYMS. Cornish Gilliflower (3). Five-Quartered Gilliflower (7, 9). Jellyflower (7, 9). Red Gilliflower (3, 5, of some 7 and 9). Ribbed Gilliflower (7, 9). SCALLOPED GILLIFLOWER (1). SCALLOPED GILLIFLOWER (5). SCOLLOP GILLIFLOWER (2, 4, 6, 7, 8, 9). SCOLLOPED GILLIFLOWER (3). *Scalloped Gilliflower* (6).

An old variety of unknown origin which was formerly grown to some extent in this state but is now practically obsolete. It has sometimes been

confounded with the Red Gilliflower of Elliott (2) and sometimes with Striped Gilliflower (9). It has been much esteemed in some portions of Ohio (2). Downing describes it as a moderate or poor grower with young shoots much darker colored than those of Striped Gilliflower, the tree more spreading and unproductive and the fruit more ribbed, much darker, rather dull red with broader stripes and splashes, with flesh more yellow, mildly subacid, aromatic, richer in quality and a month or more later in ripening than Striped Gilliflower (9).

Elliott (2) describes the fruit as "medium to large, roundish conical, flattened at base, tapering toward the eye, sometimes angular, always much ribbed or scalloped; light yellow, striped and splashed with shades of light and dark red; stem short, slender; cavity deep, russeted, irregular; calyx with long segments; basin abrupt, deep, ribbed; core large, hollow; seeds ovate, rounded; flesh yellowish, firm, crisp, tender, juicy, slight tinge of sweet. November to February."

SCOTT BEST.

REFERENCE. 1. Downing, 1869:349.

SYNONYMS. None.

We are unacquainted with this variety and have received no report concerning it from any of our correspondents. According to Downing it originated on the farm of Luther Scott, Hinsdale, Cattaraugus county, N. Y. The tree is moderately vigorous, spreading; the fruit medium to large, yellowish, shaded and mottled with light red, striped and splashed with crimson; flesh whitish, fine, tender, subacid, good to very good. Season November and December (1).

SENECA FAVORITE.

REFERENCES. 1. *Mag. Hort.*, 19:165. 1853. 2. Warder, 1867:731.


SYNONYMS. None.

A large, attractive, pale yellow apple. It resembles Swaar, but is earlier and larger, and its texture is more crisp.¹ It is excellent for dessert as well as for culinary uses. The crop begins to ripen in early autumn, and continues ripening in succession through a period of several weeks. The later fruit may be kept into early winter or midwinter. The tree is of medium size, vigorous, round-headed. It is a desirable variety for the home orchard.

Historical. N. S. Page states² that the original tree of Seneca Favorite grew upon his father's farm, five miles southwest of Geneva, and was an old tree forty-five years ago. It has been grown to a limited extent as an apple for the home orchard in various localities in Ontario county, particu-

¹ Wilson, C. S., *Hist. of the Apple in N. Y. State*, unpublished thesis Cornell Univ. 1905.

² Letter, 1906.



SHARP

larly in the town of Seneca. Downing gives Seneca Favorite as a synonym for Quince¹ but the true Seneca Favorite is surely not identical with Quince. It is now seldom propagated.

FRUIT.

Fruit very large to medium, usually large. *Form* variable, roundish conic to oblong conic, nearly regular but somewhat ribbed and elliptical; axis sometimes oblique; sides usually unequal. *Stem* long to medium, moderately slender. *Cavity* acute, deep, broad, quite strongly furrowed and compressed, usually somewhat russeted. *Calyx* small to above medium, closed or somewhat open; lobes broad, acute to acuminate. *Basin* very small, shallow or very shallow, narrow, obtuse to somewhat abrupt, slightly furrowed and wrinkled.

Skin rather thick, tough, smooth or slightly roughened by russet dots, at first green but becoming attractive pale yellow often with faint blush which sometimes deepens to a moderately bright rather pinkish-red, not striped. *Dots* numerous, irregular, large and small, varying from prominent russet to obscure and submerged, often reddish on blushed cheek.

Calyx tube funnel-form usually with long, narrow cylinder but sometimes short. *Stamens* median to basal.

Core rather large, somewhat abaxile; cells symmetrical, closed or somewhat open; core lines clasping. *Carpels* elliptical, emarginate, smooth. *Seeds* numerous, medium to above, dark brown, plump, obtuse to acute.

Flesh tinged with yellow, moderately coarse, crisp, tender, juicy, agreeably subacid, sprightly, very good.

Season fall and early winter to midwinter.

SHARP.

REFERENCES. 1. Beach, *N. Y. Sta. An. Rpt.*, 11:602. 1893. 2. *Ib.*, *Gard. and For.*, 8:428. 1895. 3. Burrill and McCluer, *Ill. Sta. Bul.*, 45:311. 1896. 4. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:56. 1903. 5. Beach and Clark, *N. Y. Sta. Bul.*, 248:144. 1904.

SYNONYMS. None.

This at its best is an excellent dessert fruit of very attractive appearance and very good quality. It is less suitable for most culinary uses because it is mildly subacid or nearly sweet, and it is not a good market variety because very often it is below medium size and not highly colored. It resembles Maiden Blush somewhat in shape and color. In this region it comes in season early in October or late in September. In ordinary storage it commonly reaches its commercial limit in November, but sometimes a portion of the fruit keeps till March. The tree is not a strong grower, but it comes into bearing young and yields full crops biennially.

¹ Downing, 1872:10 index, app.

It is distinct from both Sharp Greening and Sharp Russet. Buckman believes that it is the same as the Butler or Butler Sweet of Pennsylvania.¹

Historical. Received from Benjamin Buckman, Farmingdale, Illinois, in 1889 for testing at this Station. Mr. Buckman obtained his stock from the Illinois Experiment Station. That Station secured the variety from A. N. Lawver, who received it from Halliday and Son, Baltimore, Maryland.

TREE.

Tree below medium size, a slow grower with short, moderately stout branches. *Form* upright spreading or roundish, open. *Twigs* short to below medium, stout to rather slender, straight; internodes medium. *Bark* dull brown or brownish-red with some olive-green, streaked with thin scarf-skin; slightly pubescent. *Lenticels* quite numerous, medium to small, oblong, slightly raised. *Buds* medium size or below medium, prominent, plump, obtuse, free or nearly so, pubescent.

FRUIT.

Fruit often below medium, sometimes above medium, uniform in shape and size. *Form* roundish oblate to roundish conic, sometimes approaching oblong conic, regular or very faintly ribbed, symmetrical. *Stem* often very short and not exerted. *Cavity* usually rather large, acute to acuminate, moderately deep to deep, moderately wide to wide, sometimes very slightly furrowed and often russeted. *Calyx* small to medium, closed or partly open; lobes long. *Basin* moderately shallow to rather deep, moderately wide, abrupt, smooth or sometimes slightly ridged or wrinkled.

Skin attractive pale yellow partly covered with a bright bluish. *Dots* minute, pale or brown.

Calyx tube funnel-form. *Stamens* median to basal.

Core medium in size, somewhat abaxile; cells open or closed; core lines clasping. *Carpels* broadly roundish or somewhat elliptical, emarginate. *Seeds* medium or above, moderately long, rather flat, obtuse, dark.

Flesh whitish, moderately firm, fine-grained, tender, crisp, juicy, mild subacid, nearly sweet, very good.

Season late September into October.



SHERMAN.

REFERENCES. 1. *Rural N. Y.*, 1870 (cited by 3). 2. Downing, 1872:31 app. 3. Ragan, *U. S. B. P. I. Bul.*, 56:281. 1905.

SYNONYMS. SHERMAN (3). SHERMAN'S FAVORITE (2). *Sherman's Favorite* (3). SHERMAN'S SWEET (1). *Sherman's Sweet* (2, 3).

A yellow sweet apple of good medium size and mild, rather rich flavor; in season from November to January (2). Downing states that it originated on the farm of E. C. Sherman, Wyoming, N. Y. We have received no report concerning this variety and find no account of its having been grown outside of the place of its origin.

¹ Letter, 1895.



SHIAWASSEE

SHIAWASSEE.

REFERENCES. 1. Lyon, *Mag. Hort.*, 26:64. 1860. 2. Hovey, *Ib.*, 27:67. 1861. fig. 3. *Ib.*, 29:126. 1863. 4. Warder, 1867:459. fig. 5. Downing, 1869:351. 6. *Mich. Pom. Soc. Rpt.*, 1872:459. fig. 7. Thomas, 1875:204. 8. *Am. Pom. Soc. Cat.*, 1875:12. 9. Lyon, *Mich. Hort. Soc. Rpt.*, 1881:220. 10. Barry, 1883:354. 11. Cook, *Rural N. Y.*, 45:257. 1886. 12. *Can. Hort.*, 11:43. 1888. 13. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 14. *Ib.*, *Amer. Gard.*, 12:244. 1891. 15. *Can. Hort.*, 15:117. 1892. 16. Bailey, *Am. Hort.*, 1892:249. 17. Miller, *Rural N. Y.*, 53:278. 1894. 18. Hoskins, *Amer. Gard.*, 15:299. 1894. 19. Munson, *Me. Sta. Rpt.*, 1896:72. 20. Burrill and McCluer, *Ill. Sta. Bul.*, 45:341. 1896. 21. Van Deman, *Rural N. Y.*, 56:830. 1897. 22. Macoun, *Can. Dept. Agr. Rpt.*, 1899:77. 23. Waugh, *Vt. Sta. Bul.*, 83:92. 1900. 24. *Ib.*, *Vt. Sta. An. Rpt.*, 14:308. 1901. 25. Macoun, *Can. Dept. Agr. Bul.*, 37:41. 1901. 26. Munson, *Me. Sta. An. Rpt.*, 18:84, 90, 95. 1902. 27. Farrand, *Mich. Sta. Bul.*, 205:46. 1903. 28. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:56. 1903. 29. Budd-Hansen, 1903:174. 30. Beach and Clark, *N. Y. Sta. Bul.*, 248:144. 1904.

SYNONYMS. *Michigan Beauty* (5). *Shiawassee* (21). *Shiawassee Beauty* (3, 7, 8, 20). *SHIAWASSEE* (14, 26). *SHIAWASSEE BEAUTY* (1, 2, 4, 5, 6, 9, 10, 11, 12, 16, 17, 18, 19, 22, 25). *SHIAWASSEE Beauty* (27). *Shiawassee Beauty* (23, 24, 28, 29, 30). *SHIAWASSEE BEAUTY* (15).

Fruit of the Fameuse type, of good size, quite attractive appearance and pleasant dessert quality. It has a flavor and aroma somewhat similar to that of McIntosh. Probably McIntosh would be preferred to Shiawassee by most fruit growers. The fruit of Shiawassee is fair, uniform and hangs pretty well to the tree. It ripens in October, and some portion of the fruit may be held till January. The tree is of uniform size, very hardy, vigorous, upright spreading, healthy and long-lived. It does not come into bearing very young, and when mature in some cases it is not a reliable cropper, but it is generally reported as yielding good to heavy crops biennially, or sometimes annually.

Historical. The original tree was planted as an ungrafted seedling in the orchard of Beche Truesdell, in Vernon, Shiawassee county, Mich., and came into bearing about 1850 (1). The variety is often listed by nurserymen but is seldom planted in this state.

FRUIT.

Fruit medium to nearly large, uniform in shape but not in size. *Form* oblate conic, pretty regular but sometimes elliptical. *Stem* medium in length, slender to moderately thick. *Cavity* acute, varying from nearly acuminate to somewhat obtuse, deep, broad, rather symmetrical, sometimes compressed, often with outspreading russet rays. *Calyx* small to below medium, closed or

slightly open; lobes rather short, moderately narrow. *Basin* rather shallow to moderately deep, rather wide, obtuse to somewhat abrupt, somewhat furrowed and wrinkled, often compressed.

Skin rather pale yellow, usually entirely overspread with attractive red, irregularly splashed and striped with carmine. *Dots* small to medium, grayish.

Calyx tube medium size, moderately wide, conical to short funnel-shape. *Stamens* median to nearly basal.

Core below medium size, widely abaxile; cells symmetrical, usually open; core lines meeting or slightly clasping. *Carpels* cordate to broadly ovate. *Seeds* rather dark brown, medium size, rather narrow, plump, acute.

Flesh white, fine, crisp, tender, juicy, pleasant subacid, aromatic, rather sprightly, good to very good.

Season October to January.

SINE-QUA-NON.

REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 1220. 2. Downing, 1845:76. 3. Thomas, 1849:143. 4. Cole, 1849:103. 5. Waring, 1851:30. 6. Barry, 1851:281. 7. Elliott, 1854:157. 8. Hooper, 1857:84. 9. Gregg, 1857:39. 10. *Horticulturist*, 14:425. 1859. 11. *Am. Pom. Soc. Cat.*, 1862. 12. Warder, 1867:732. 13. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296.

SYNONYMS. None.

An August apple of good dessert quality, now seldom found in cultivation. It originated on Long Island and was brought to notice by Wm. Prince (2). It was entered on the catalogue of the American Pomological Society in 1862 (11) and dropped from that list in 1871. The tree is a rather slow, crooked grower, in some cases an indifferent bearer (7), in others productive (4, 13). Fruit medium size, roundish ovate, pale greenish-yellow; flesh greenish-white, tender, juicy, mild subacid, sprightly, good; season late August.

SLINGERLAND.


REFERENCES. 1. *N. Y. Sta. Agr. Soc. Rpt.*, 1849:594. 2. Emmons, *Nat. Hist. N. Y.*, 3:42. 1851. col. pl. No. 32. 3. Downing, 1857:189. 4. Warder, 1867:732. 5. Thomas, 1875:512. 6. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:206.

SYNONYMS. SLINGERLAND (6). SLINGERLAND'S FALL PIPPIN (2). SLINGERLAND PIPPIN (1, 3, 4, 5).

An excellent flavored apple of the Green Newtown type in season during late fall and early winter. It is not as good a keeper as Green Newtown. Raised from seed of the Newtown about 1830 by a Mr. Slingerland of New Scotland, Albany county, N. Y. (1, 2).

FIGURE (1, 2, 3)

Fruit medium to large. *Form* roundish, often oblique. *Stem* exerted but short. *Skin* yellow splashed with reddish-orange over the base. *Dots* minute. *Calyx* small to medium, partly closed. *Core* small. *Flesh* whitish tinged with yellow, firm, tender, juicy, brisk, rather rich subacid, good to very good. *Season* December to February or later.



SOPS OF WINE

SOMERSET (N. Y.).

REFERENCES. 1. Downing, 1869:356. 2. Downing, *Tilt. Jour. Hort.*, 7:303. 1870. 3. *Am. Pom. Soc. Cat.*, 1877:14. 4. Lyon, *Mich. Hort. Soc. Rpt.*, 1881:318. 5. Hoag, *Am. Pom. Soc. Rpt.*, 1885:28. 6. Lyon, *Mich. Sta. Bul.*, 129:40, 43. 1896. 7. *Mich. Sta. Bul.*, 152:222, 226. 1898.

SYNONYMS. None.

An early apple of high sprightly flavor. A fine family fruit. The tree is an unusually early bearer, upright spreading, vigorous and productive. Season late August to October (4, 6, 7).

There is a distinct variety of Maine origin which is also called Somerset.

Historical. Brought to notice by C. L. Hoag, Lockport, N. Y. Origin unknown but supposed to be Somerset, Niagara county, N. Y. (2). This variety appears to have been but sparingly disseminated.

FRUIT (2, 5).

Fruit below medium. *Form* like Black Gilliflower, roundish conical. *Skin* deep golden-yellow to whitish-yellow with sometimes a few nettings of russet and sparsely sprinkled with brown dots. *Flesh* quite white, tender, juicy, with a rich aromatic flavor; quality very good or best.

SOPS OF WINE.

REFERENCES. 1. Ray, 1688:No. 21. 2. Knoop, 8:45. 1758. 3. Kenrick, 1832:28. 4. *Ib.*, 1835:98. 5. Floy-Lindley, 1833:25. 6. Downing, 1845:77. 7. Cole, 1849:103. 8. Thomas, 1849:141. 9. Emmons, *Nat. Hist. N. Y.*, 3:111, 33. 1851. 10. Barry, 1851:282. 11. Hovey, *Mag. Hort.*, 18:545. 1852. fig. 12. Elliott, 1854:157. 13. Gregg, 1857:39. 14. Hooper, 1857:85. 15. *Am. Pom. Soc. Cat.*, 1862. 16. Warder, 1867:615. fig. 17. Downing, 1869:356. 18. *Horticulturist*, 27:309. 1872. fig. 19. Fütz, 1872:121, 174. 20. Downing, 1872:10 index, app. 21. *Montreal Hort. Soc. Rpt.*, 1879:22. 22. Downing, 1881:11 index, app. 23. *Ib.*, 1881:12 index, app. 24. Hogg, 1884:215. 25. Hoskins, *Rural N. Y.*, 47:662. 1888. figs. 26. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:296. 27. Bailey, *Am. Hort.*, 1892:249. 28. Hoskins, *Rural N. Y.*, 53:248. 1894. 29. Burrill and McCluer, *Ill. Sta. Bul.*, 45:325. 1896. 30. Waugh, *Vt. Sta. An. Rpt.*, 14:309. 1901. 31. Budd-Hansen, 1903:177. fig.

SYNONYMS. *Bell's Early?* (17). *Bell's Favorite* (20). *Bennington* (17, 18). *Dodge's Early Red* (17). *Early Washington* (22). *Hominy* (17, 19). *Horning* (29). *Pie Apple* (10). *Red Shropshire?* (17). *RODE WYN APPEL* (2). *Rode Wyn Appel* (5, 6, 9). *SAPSON* (3). *Sapson* (6, 9, 11, 12, 18). *Sapsontine* (3). *Shropshirevine* (23). *SOPS IN WINE* (1, 24). *Sops in Wine* (5, 6, 9, 11, 12, 18). *SOPS OF WINE* (4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 29, 30, 31). *Sops of Wine* (24). *Strawberry* (30). *Warden's Pie Apple* (17, 18). *Washington* (17, 18).

A dark crimson apple; flesh fine, stained with red; in season in August and September. The tree is a good grower, comes into bearing rather young and is a biennial or nearly annual cropper. Of little value except as a dessert apple for family use.

Historical. "A very ancient English culinary and cider apple" (24). It is frequently listed by nurserymen but is now seldom planted in New York being superseded by other better varieties.

TREE.

Tree medium to large, moderately vigorous to vigorous. *Form* upright or inclined to roundish, rather dense. *Twigs* short to rather long, curved, moderately stout; internodes short. *Bark* dark brown, lightly streaked with scarf-skin; heavily pubescent. *Lenticels* very scattering, small, oblong, not raised. *Buds* medium size, broad, obtuse, free, pubescent.

FRUIT.

Fruit medium to sometimes large, uniform in shape but not in size. *Form* roundish to roundish conic, slightly ribbed; sides unequal. *Stem* short to rather long, moderately slender. *Cavity* acute, moderately deep, medium or sometimes narrow, sometimes slightly furrowed, sometimes with thin, radiating russet rays. *Calyx* medium to rather small, closed or slightly open; lobes rather short. *Basin* shallow, narrow, furrowed, somewhat wrinkled.

Skin moderately thin, moderately tender, slightly roughened, greenish-yellow almost entirely overspread with purplish-red, mottled, irregularly splashed and sometimes indistinctly striped with dark carmine, overspread with thin white bloom. *Dots* small, few, light russet or yellow.

Calyx tube short, wide, cone-shape. *Stamens* marginal to median.

Core medium size, somewhat abaxile; cells usually symmetrical but not uniformly developed, open to nearly closed; core lines meeting. *Carpels* broad ovate, rather concave, mucronate, tufted. *Seeds* rather large or medium size, moderately wide, plump, obtuse.

Flesh yellowish often stained with pink, soft, fine, juicy, aromatic, mild, pleasant, subacid, good.

Season August to October.

SOUR BOUGH.

REFERENCE. 1. Downing, 1869:357.

SYNONYMS. None.

This is an old Westchester county variety which, according to Downing, is of medium size, roundish conic, yellow with whitish, brisk subacid flesh, good for cooking. Season, September. "Often knotty and unprofitable" (1).

The name Sour Bough has also been applied sometimes to the Champlain; see page 30; and also to the Tart Bough; see page 220.

SPECTATOR.

REFERENCE. 1. Downing, 1869:357.

SYNONYMS. None.



SOUR BOUGH

Originated with J. W. Bailey, Plattsburg, N. Y. According to Downing (1) this is an apple of medium size, ribbed, yellow, shaded with red in the sun; flesh white, subacid, hardly good; season September. The variety is unknown to us.

STARKEY.

REFERENCES. 1. *Am. Pom. Soc. Rpt.*, 1875:65. 2. Downing, 1876:69 app. 3. Thomas, 1885:524. 4. Munson, *Me. Sta. Rpt.*, 1893:133. 5. *Am. Pom. Soc. Cat.*, 1897:14. 6. Budd-Hansen, 1903:179.

SYNONYMS. None.

Fruit of good size, fair, well colored, largely striped and splashed with red, pleasant subacid, excellent for dessert or culinary use. Season, October to midwinter. In Maine, where it originated, it is said to be popular in market. It has not been sufficiently tested to determine its value for this state, but it appears to be worthy of testing. Tree a hardy and vigorous grower in the nursery, becoming rather large and spreading in the orchard; comes into bearing rather young and is a good biennial cropper.

Historical. Originated on the farm of Moses Starkey, North Vassalboro, Kennebec county, Me.

TREE.

Tree medium to rather large, moderately vigorous to vigorous with long, moderately stout, curved branches. *Form* spreading or roundish, open. *Twigs* medium to long, stout, geniculate, often irregularly crooked; internodes long to below medium. *Bark* brownish-red mingled sometimes with olive-green, irregularly mottled with scarf-skin; pubescent near tips. *Lenticels* scattering, inconspicuous, medium size or below, round, raised. *Buds* medium to large, prominent, plump, acute, free, slightly pubescent.

FRUIT.

Fruit medium to rather large. *Form* oblate or oblate conical, regular or faintly ribbed, symmetrical. *Stem* medium length. *Cavity* medium size, rather shallow, partly covered with greenish russet. *Calyx* large, closed or nearly so. *Basin* medium size, shallow, somewhat wrinkled.

Skin pale yellow, washed or deeply blushed with lively red, splashed and striped with carmine. *Dots* pale gray or russet. *Prevailing effect* red.

Calyx tube conical or somewhat funnel-form.

Core rather small.

Flesh whitish, firm, rather fine, juicy, crisp, pleasant, lively subacid, becoming mild subacid, very good.

Season October to January.

STARR.

REFERENCES. 1. Downing, *Tilt. Jour. Hort.*, 6:347. 1869. fig. 2. Downing, 1869:360. 3. Thomas, 1875:512. 4. Hexamer, *Am. Pom. Soc. Rpt.*, 1895:70. 5. *Rural N. Y.*, 54:587. 1895. 6. Parry, Wm., Parry, N. J., *Cat.*, 1896.

SYNONYMS. None.

Fruit large, very attractive for a green or yellowish apple, and very good in quality, especially for dessert use. Season, August and September. The tree is a pretty good grower, comes into bearing young and as tested at this Station gives promise of being an annual bearer. Starr appears to be worthy of testing where a fruit of this type is desired.

Historical. The original tree was found on the grounds of Judge J. M. White, Woodbury, N. J., which property afterward came into the possession of Mrs. Starr. The propagation of the variety was begun by Wm. Parry in 1865 under the name of Starr (6). So far as we can learn it has been but little planted in New York.

TREE.

Tree moderately vigorous with short, moderately stout, curved branches. *Form* upright spreading or roundish, rather dense. *Twigs* long, curved, stout with large terminal buds; internodes long. *Bark* brownish-red, tinged with olive-green, lightly streaked with scarf-skin; pubescent near tips. *Lenticels* quite numerous, medium size, round, not raised. *Buds* prominent, large, long, broad, plump, acute, free, pubescent.

FRUIT.

Fruit very large to large, pretty uniform in size and shape. *Form* distinctly oblate to roundish oblate, regular or faintly ribbed. *Stem* short to medium, moderately thick, sometimes swollen. *Cavity* acute, varying from a little obtuse to somewhat acuminate, shallow to medium, broad, smooth or gently furrowed. *Calyx* medium size, closed, lobes long to medium, rather narrow, acuminate. *Basin* medium in depth, narrow, abrupt, somewhat furrowed.

Skin rather thick, tough, nearly smooth, green becoming yellowish-green, sometimes with indications of a faint blush. *Dots* numerous, small and large, pale or russet.

Calyx tube long, very wide to moderately wide, conical to cylindrical and large, extending to the core. *Stamens* nearly marginal.

Core medium to rather large, abaxile to nearly axile; cells closed or slightly open; core lines clasping. *Carpels* obovate, sometimes tufted. *Seeds* dark brown, medium to large, rather wide, plump, acute to nearly acuminate.

Flesh tinged with yellow, moderately fine, very tender, crisp, very juicy, sprightly subacid, aromatic, very good.

Season August and September.

STILLMAN EARLY.

REFERENCES. 1. Downing, 1857:193. 2. Thomas, 1875:512. 3. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298.

SYNONYMS. STILLMAN (3). STILLMAN'S EARLY (1, 2).

This variety originated in Clinton, Oneida county, N. Y. Downing (1) states that the tree is a moderate grower and productive; the fruit small, yellow, sometimes slightly blushed; flesh pleasant subacid, good; season late July and early August. We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

STRAWBERRY.

This name has been applied to a great many different varieties of the apple. Those mentioned in this volume are listed below:

Autumn Strawberry, see Late Strawberry. *Chenango Strawberry*, see Chenango. *Early Strawberry*. *Fall Strawberry*, see Late Strawberry. *Late Strawberry*. *St. John's Strawberry*, see Early Strawberry. *Strawberry*, see Chenango, Early Strawberry, Late Strawberry and Richard Graft. *Washington Strawberry*.

STRIPED GILLIFLOWER.

REFERENCES. 1. Warder, 1867:606. fig. 2. Downing, 1876:90 app. 3. Bailey, *In. Hort.*, 1892:250. 4. Burrill and McCluer, *Ill. Sta. Bul.*, 45:342. 1896. 5. Thomas, 1897:354. 6. Bodd-Hansen, 1903:181.

SYNONYMS. *Red Gilliflower* (2). *Scalloped Gilliflower* (incorrectly) (2). *Striped Gilliflower* (2). *Striped Gilliflower* (1, 2, 3, 4, 5, 6).

This variety appears to be practically obsolete in New York. The tree is a vigorous grower and generally productive (2).

FRUIT (1, 2, 6).

Fruit large to very large. *Form* variable roundish conic to oblong conic, often furrowed, angular, but less ribbed than Scalloped Gilliflower. *Stem* short, curved. *Cavity* acute, deep, wide, furrowed, brown or russeted. *Calyx* large, closed or partly open; lobes erect. *Basin* shallow, abrupt.

Skin yellowish-white or greenish, partly covered with dull red, striped and splashed with carmine. *Dots* few, indistinct, gray or white.

Calyx tube cone-shape. *Stamens* median.

Core sessile, abaxile, large; cells wide open; core lines meeting or slightly clasping. *Carpels* tufted. *Seeds* few, small, roundish, plump, obtuse, black.

Flesh yellowish-white, breaking, juicy, briskly subacid, fair to good.

Season September.

STROAT.

REFERENCES. 1. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 2. *London Hort. Soc. Cat.*, 1831:No. 1256. 3. Kenrick, 1832:39. 4. Downing, 1845:97. 5. *Horticulturist*, 2:545. 1848. 6. Thomas, 1849:158. 7. Emmons, *Nat. Hist. N. Y.*, 3:38. 1851. 8. Elliott, 1854:158. 9. Hooper, 1857:87.

SYNONYMS. STRAAT (2, 3). *Straat* (4, 8, 9). STROAT (1, 4, 5, 6, 7, 8, 9).

Stroat was formerly much esteemed among the descendants of the Dutch settlers on the North river (4). The fruit is described as above medium, roundish inclined to conic, yellowish-green; flesh yellow, very tender, rich, brisk subacid, good to very good; season September to November or December (4, 6, 8).

We have not seen this fruit nor has it been mentioned by any of our correspondents.

STRODE BIRMINGHAM.

REFERENCES. 1. Downing, 1857:193. 2. Warder, 1867:733. 3. *Pa. Sta. Hort. Assn. Rpt.*, 1886:50. 4. Powell and Fulton, *U. S. P. B. I. Bul.*, 48:57. 1903. 5. Beach and Clark, *N. Y. Sta. Bul.*, 248:145. 1904.

SYNONYMS. *Dumpling* (3). STRODE (4, 5). *Strode's* (1). STRODE'S BIRMINGHAM (1, 2, 3). *Strode's Birmingham* (4, 5).

A medium-sized yellow apple of mild subacid flavor and good quality. Commercial limit September. For home use it is in season in September and October and a few specimens may be kept till January (5). It is reported as a desirable variety for market and general purposes in Pennsylvania (3). The tree comes into bearing rather young and is productive yielding moderate to good crops nearly annually. As compared with standard varieties of its season it does not appear to be worthy of the attention of New York fruit growers.

FRUIT.

Fruit of medium size; uniform in size and shape. *Form* roundish conic or inclined to oblong, regular or faintly ribbed; sides unequal. *Stem* long, slender. *Cavity* acute to almost acuminate, usually rather deep, medium in width, symmetrical, russeted. *Calyx* medium to small, closed or partly open; lobes medium in length and width, acute, reflexed. *Basin* shallow, narrow to medium in width, obtuse, furrowed and wrinkled.

Skin thin, tender, smooth, clear yellow or greenish, often with faint blush and marked with russet flecks. *Dots* scattering, very minute, submerged, inconspicuous, red or russet.

Calyx tube short, wide, urn-shape to broadly conical. *Stamens* nearly basal.

Core medium to small, axile; cells almost closed; core lines meeting. *Carpels* broadly ovate to oblong narrowing toward either end, deeply emarginate. *Seeds* dark dull brown, medium to large, wide, plump, broadly acute, tufted.



STUMP

Flesh yellowish, rather firm, fine, rather crisp, tender, juicy, brisk subacid becoming mild and pleasant when fully mature, good to very good.

Season September to early winter.

STUMP.

REFERENCES. 1. Downing, 1881:110 app. fig. 2. Barry, 1883:341. 3. Thomas, 1885:525. 4. *Pa. Sta. Hort. Assn. Rpt.*, 1885:25. 5. *Can. Hort.*, 11:8. 1888. 6. *Rural N. Y.*, 47:692. 1888. figs. 7. Bailey, *Am. Hort.*, 1892:250. 8. *N. Y. Sta. An. Rpt.*, 14:255. 1895. 9. Beach, *Gard. and For.*, 8:428. 1895. 10. *Kan. Sta. Bul.*, 106:55. 1902. 11. Beach and Clark, *N. Y. Sta. Bul.*, 248:145. 1904.

SYNONYMS. None.

This very beautiful apple resembles Chenango in form, size and general appearance. In Western New York it is by some considered a good variety for commercial planting, but others find difficulty in marketing it with profit, since it is esteemed chiefly for dessert purposes and ripens at a season when there is comparatively little demand for apples, and shows bruises so readily that it is not well adapted for barrelling. It may be marketed in local markets or shipped in small packages. The crop ripens unevenly, and on this account should have more than one picking in order to secure the fruit in prime condition. It begins to ripen in late August or early September. Its commercial limit is September or early October, but its season for home use extends to November (11). The trees are upright, compact and stocky, so that they may stand closer together than common varieties in the orchard, or be used for alternating with more spreading trees. They are hardy, long-lived, and reliable croppers, yielding good to heavy crops biennially. The fruit is borne on short spurs close to the limbs. It is recommended for planting in the garden and commercial orchard where a variety of this type is desired.

Historical. Originated as a chance seedling in an old stump on the grounds of John Prue, Chili, N. Y. (1). It appears that it has been planted more in Western New York than in any other section of the state.

TREE.

Tree a slow, stiff, upright grower in the nursery, apt to crack at the collar, not popular with nurserymen; in the orchard it becomes a moderately vigorous or vigorous grower; branches long, moderately stout, curved, filled

with fruit spurs. *Form* very upright but eventually somewhat roundish. *Twigs* moderately long, curved, moderately stout; internodes medium. *Bark* brown tinged with green, lightly streaked with scarf-skin; pubescent. *Lenticels* numerous, medium size, round, not raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium or below, uniform in size and shape. *Form* roundish conic to oblong conic, regular or faintly ribbed; sides sometimes unequal. *Stem* very short, variable in thickness. *Cavity* acute or approaching obtuse, usually rather shallow, medium in width to rather broad, unsymmetrical, frequently furrowed, sometimes lipped, partly russeted. *Calyx* medium to small, slightly open or closed; lobes short, rather narrow, acute. *Basin* shallow, narrow, abrupt.

Skin smooth, clear pale yellow largely washed and mottled with bright pinkish-red, becoming deep red in highly-colored specimens, rather indistinctly striped and splashed with bright carmine. *Dots* numerous, rather large, areolar with russet point or with whitish point. *Prevailing effect* red and yellow contrasting beautifully.

Calyx tube broadly conical with fleshy pistil point projecting into the base. *Stamens* basal.

Core below medium to rather large, usually abaxile; cells open; core lines meeting or slightly clasping. *Carpels* elongated ovate tapering toward base and apex. *Seeds* rather dark brown, medium to small, usually very wide, short, very plump, obtuse to acute.

Flesh whitish or tinged with yellow, rather fine, tender, juicy, rich, aromatic, sprightly, pleasant subacid, very good.

Season September and October.

STYMUS.

REFERENCES. 1. *Am. Pom. Soc. Rpt.*, 1867 (cited by 3). 2. Downing, 1869:365. 3. Ragan, *U. S. B. P. I. Bul.*, 56:298. 1905.

SYNONYMS. *Stymer's* (3). STYMUS (1, 2, 3).

This variety was described by Downing in 1869 as a new and really excellent apple which originated on the farm of Jacob Stymus, Dobbs Ferry, N. Y. (2). Fruit medium size, yellowish, shaded, splashed and striped with light and dark crimson; flesh fine, tender, pleasant subacid, very good; season October and November.

We are unacquainted with this variety and have received no report concerning it from any of our correspondents.

SUFFOLK BEAUTY.

REFERENCE. 1. Downing, 1869:365.

SYNONYMS. None.

Described in 1869 by Downing (1) as a new variety from Deer Park, Long Island. Fruit medium, yellowish-white; flesh subacid; season August and September.

This variety is unknown to us and we have received no report concerning it from any of our correspondents.

SUMMER BELLFLOWER.

REFERENCES. 1. *Horticulturist*, 3:168. 1848. fig. 2. Thomas, 1849:143. 3. Cole, 1849:104. 4. Waring, 1851:31. 5. Elliott, 1854:159. 6. Downing, 1857:196. 7. Warder, 1867:733. 8. *Am. Pom. Soc. Cat.*, 1875:14.

SYNONYMS. SUMMER BELLEFLEUR (1, 8). *Summer Belle-fleur* (5). SUMMER BELLFLOWER (2, 3, 4, 5, 6, 7). *Summer Bellflower* (1).

This fruit bears considerable resemblance in form and color to Yellow Bellflower. Flesh tender, subacid, good. It is in season from the middle of August to the middle of September. The tree is erect, of medium size, a good grower, hardy and a good cropper. It is not considered a satisfactory variety for commercial planting and so far as we can learn is gradually becoming obsolete in New York. Downing described it in 1848 as a new variety of promise and stated that it was raised by John R. Comstock of Washington, Dutchess county, N. Y., from seed of the Esopus *Spitzenburg* (1, 6). It was entered on the list of the American Pomological Society in 1875 as a variety of value for Nebraska (8). It was dropped from that list in 1897.

A distinct variety of Pennsylvania origin has also been known under the name Summer Bellflower (6, 7).

SUMMER PEARMAIN.

REFERENCES. 1. McMahon, *Amer. Gard. Cal.*, 1806:585. 2. Cox, 1817:104. fig. 3. Cobbett, 1821:par. 300. 4. Thacher, 1822:138. 5. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 6. Fessenden, 1828:129. 7. Kenrick, 1832:25. 8. *Mag. Hort.*, 1:398. 1835. 9. Manning, 1838:47. 10. *Ib.*, *Mag. Hort.*, 7:49. 1841. 11. Downing, 1845:70. 12. *Horticulturist*, 2:544. 1848. 13. Thomas, 1849:136. fig. 14. Cole, 1849:103. 15. Phoenix, *Horticulturist*, 4:472. 1850. 16. Emmons, *Nat. Hist. N. Y.*, 3:11. 1851. fig. 17. Barry, 1851:279. 18. *Am. Pom. Soc. Cat.*, 1852. 19. Elliott, 1854:64. fig. 20. Hooper, 1857:12, 106, 108. 21. Gregg, 1857:35. 22. Warder, 1867:582. fig. 23. Downing, 1869:78. fig. 24. Fitz. 1872:143, 160. 25. Hogg, 1884:7. 26. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 27. Bailey, *Am. Hort.*, 1892:234. 28. *Ib.*, 1892:250. 29. Budd-Hansen, 1903:182. fig.

SYNONYMS. AMERICAN PEARMAIN (19). *American Pearmain* (20). *American Summer* (26). AMERICAN SUMMER PEARMAIN (7, 8, 11, 12, 13, 14, 16, 17, 20, 21, 22, 23, 24, 25, 27). *American Summer Pearmain* (19, 29). EARLY SUMMER PEARMAIN (2, 5, 6). *Early Summer Pearmain* (7, 11, 14, 16, 19, 23, 25, of COX 13). SUMMER PEARMAIN (1, 3, 4, 9, 10, 15, 18, 26, 28, 29). *Summer Pearmain* (14). *Watkins Early* (19, 20).

An amateur fruit which when perfect is beautiful and of mild, rich, excellent flavor (26). The tree being of slender, slow growth in the nursery is not a favorite with nurserymen and although it makes a large productive tree in the orchard it is not profitable as a market variety (22). It is desirable for family use because the fruit is suitable both for culinary and dessert purposes and the crop ripens in succession through a period of nearly two months.

There is also another Summer Pearmain or English Summer Pearmain which ripens somewhat later.¹

Historical. Supposed to be of American origin. It is an old variety; first described in 1817 by Cox. *e.*

FRUIT (11, 14, 22, 23).

Fruit of medium size. *Form* variable, oblong or roundish inclined to conic, sometimes oblate. *Stem* medium to long. *Cavity* deep, acute, regular. *Calyx* large, open or closed. *Basin* medium size, abrupt, slightly wrinkled.

Skin smooth, greenish-yellow, more or less covered with dull purplish-red, marbled, splashed and striped with brighter red. *Dots* minute.

Core medium to small, roundish; cells closed. *Seeds* small, pointed.

Flesh yellowish, very fine, tender, almost melting, juicy, aromatic, crisp, mild subacid, best.

Season August and September.

SUMMER QUEEN.

REFERENCES. 1. M'Mahon, *Amer. Gard. Cal.*, 1806:584. 2. Cox, 1817:102. 3. Thacher, 1822:133. 4. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 5. Wilson, 1828:136. 6. Kenrick, 1832:28. 7. Manning, 1838:46. 8. Downing, 1845:77. 9. Hovey, *Mag. Hort.*, 14:489. 1848, fig. 10. Thomas, 1849:141. 11. Cole, 1849:103. 12. Barry, 1851:282. 13. Waring, 1851:28. 14. Emmons, *Nat. Hist. N. Y.*, 3:14. 1851. 15. Elliott, 1854:158. 16. Hooper, 1857:88, 107. 17. Gregg, 1857:39. 18. *Am. Pom. Soc. Cat.*, 1860:240. 19. Warder, 1867:545. fig. 20. Downing, 1869:370. fig. 21. Fitz, 1872:143, 161. 22. *Ill. Hort. Soc. Rpt.*, 1874:295. 23. Wickson, 1889:243. 24. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 25. Bailey, *An. Hort.*, 1892:250. 26. Budd-Hansen, 1903:183. fig.

SYNONYMS. *Lancaster Queen* (15, 20). *Polecat* (20). *QUEEN* (3, 13). *Queen* (6). *Sharpe's Early* (20). *SUMMER QUEEN* (2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26). *Summer Queen* (3, 13). *SWEET'S HARVEST* (1). *Sweet's Harvest* (3, 4, 6). *Sweet's Harvest* (2).

A striped red apple of good size and excellent quality for culinary use, in season during late summer. In regions farther west it is a very popular variety for home use and is also considered by some desirable for market. The tree is a moderate grower with rather spreading habit and productive, yielding good crops almost annually.

Historical. This is an old variety which is supposed to have originated in this country (4, 19, 26). It is commonly listed by nurserymen in most parts

¹ Ragan, *U. S. B. P. I. Bul.*, 56:300. 1905.

of the country (25). So far as we have been able to learn it is not often grown in New York and is now seldom planted in this state.

FRUIT (10, 12, 15, 19).

Fruit medium to large. *Form* roundish conical, somewhat angular. *Stem* medium to long, slender. *Cavity* narrow to rather wide, regular, pretty deep. *Calyx* medium to large, open or closed. *Basin* shallow or none, furrowed.

Skin yellow, striped, splashed and shaded with mixed red. *Dots* minute, yellow.

Core medium size; cells open. *Seeds* numerous, acute, brown.

Flesh whitish-yellow, sometimes with tinge of pink, firm, aromatic, juicy, subacid, good to very good for culinary use.

Season August and September.

SUMMER RAMBO.

REFERENCES. 1. Switzer, 1725 (cited by 7). 2. Duhamel, 1:28. 1768. 3. Forsyth, 1803:49. 4. Cox, 1817:104. fig. 5. Forsyth, 1824:123. 6. Kenrick, 1832:37. 7. Floy-Lindley, 1833:12. 8. Manning, 1838:47. 9. Downing, 1845:94. 10. Thomas, 1849:141. 11. Elliott, 1854:178. 12. *Am. Pom. Soc. Cat.*, 1862. 13. Warder, 1867:733. 14. *Am. Pom. Soc. Cat.*, 1871:10. 15. Leroy, 1873:598. fig. 16. *Mo. Hort. Soc. Rpt.*, 1883:76. 17. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 18. Bailey, *Am. Hort.*, 1892:250. 19. (?) *Revue Horticole*, 66:202. 1894. 20. Taylor, *Am. Pom. Soc. Rpt.*, 1895:199. 21. Burrill and McCluer, *Ill. Sta. Bul.*, 45:337. 1896.

SYNONYMS. *Cambour des Lorrains* (15). *Charmant Blanc* (15). *De Lorraine* (15). *De Rambourg* (15). *De Rambure* (15). *De Notre-Dame* (15). FRANK RAMBOUR (1, 7). *Frank Rambour* (9). *Grosh* (16). *Gros-Rambour d'Ete* (15). *Herbstbreitling* (15). *Lothringer Rambour d'Ete* (15). *Pomme de Notre-Dame* (6). RAMBOUR (3). *Rambour* (15, 17). *Rambour Blanc* (15). (RAMBOUR D'AMERIQUE, 19)? RAMBOUR D'ETE (4, 8, 15). *Rambour d'Ete* (6, 9, 10, 11). RAMBOUR FRANC (2, 5, 6, 9, 12). *Rambour Franc* (4, 7, 10, 11, 15, 21). *Rambour Aigre* (15). *Rambour Gros* (7, of the English 6). *Rambour Raye* (6, 15). *Rambu* (15). *Remboure d'Ete* (15). SUMMER RAMBO (13, 14, 16, 17, 18, 20, 21). *Summer Rambo* (11). SUMMER RAMBOUR (10, 11). *Summer Rambour* (4, 9).

Fruit of the type of Grosh; very attractive in size, form and color; large, oblate, yellowish-green considerably striped and splashed with mixed red, good to very good. Season early autumn. Begins to ripen about a month earlier than Grosh. The tree is a strong grower, comes into bearing young and bears quite regularly yielding moderate to good crops. Although it has long been known in cultivation in this country it has not gained much recognition among New York fruit growers. The fine color and size of this variety combined with its comparatively good quality recommend it for home use or local market. It is an old variety and has been tested in many parts of the United States but its cultivation has never become extensive. These facts would indicate that it has weaknesses not apparent to the casual observer. Possibly it is worthy of further trial in this state.

Historical. This variety is said to have originated in France (15). It has long been known in this country having been described by Coxé in 1817 and Kenrick in 1832 (4, 6). It was listed in the catalogue of the American Pomological Society under the name Rambour Franc from 1862 to 1871 (12, 14, 20). It is still listed by nurserymen (18). It is comparatively little known in New York state but is more often found in cultivation in Ohio and regions farther west.

TREE.

Tree vigorous. *Form* upright spreading to roundish, open. *Twigs* moderately long, curved, moderately stout; internodes medium. *Bark* brown, tinged with green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* quite numerous, medium size, round, not raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit large to very large, uniform in size and shape. *Form* oblate to roundish oblate, sometimes slightly ovate, sometimes faintly ribbed, pretty regular; sides often unequal. *Stem* short to medium, rather thick. *Cavity* nearly acuminate, deep to medium in depth, rather broad, usually symmetrical, sometimes lipped, sometimes slightly russeted close to the stem. *Calyx* rather large, closed or sometimes slightly open; lobes medium to short, rather narrow, acute to obtuse. *Basin* deep to sometimes medium, wide to medium in width, abrupt, smooth, symmetrical.

Skin thick, tough, smooth, attractive clear bright yellow or greenish, in well colored specimens largely washed and mottled with lively pinkish-red, conspicuously marked with many broken stripes and splashes of bright carmine. *Dots* numerous, usually small and submerged, but some are scattering, large, brown or russet. *Prevailing effect* striped.

Calyx tube medium in length, rather wide, broadly conical. *Stamens* median to marginal.

Core rather small, axile; cells nearly closed; core lines meeting or slightly clasping. *Carpels* roundish to roundish ovate approaching elliptical. *Seeds* frequently abortive, rather large, wide, plump, acute, moderately dark brown.

Flesh yellowish-green, firm, breaking, coarse, tender, very juicy, mildly subacid, somewhat aromatic, good.

Season September to November.

SUMMER REDSTREAK.

REFERENCE. 1. Downing, 1869:371.

SYNONYMS. None.

A September apple which originated in Columbia county, N. Y. According to Downing (1) the tree is moderately vigorous and productive. The fruit medium, yellowish, shaded, striped and splashed with rich red; flesh white, sometimes a little stained next the skin, brisk subacid, valued for cooking.

We have neither seen this variety nor received any report concerning it.

SUMMER ROSE.

REFERENCES. 1. M'Mahon, *Amer. Gard. Cal.*, 1806:584. 2. Coxe, 1817:103. fig. 3. Buel, *N. Y. Bd. Agr. Mem.*, 1826:476. 4. Wilson, 1828:136. 5. Fessenden, 1828:131. 6. Kenrick, 1832:29. 7. Manning, 1838:47. 8. Downing, 1845:77. 9. *Horticulturist*, 2:483. 1848. 10. *N. Y. Agr. Soc. Trans.*, 1848:277. fig. 11. Thomas, 1849:141. fig. 12. Cole, 1849:101. fig. 13. Waring, 1851:31. 14. Barry, 1851:282. 15. *Am. Pom. Soc. Cat.*, 1852. 16. Elliott, 1854:107. fig. 17. Gregg, 1857:39. 18. Hooper, 1857:87. 19. *Horticulturist*, 14:425. 1859. 20. Warder, 1867:616. 21. Fittz, 1872:143, 160. 22. *Ill. Hort. Soc. Rpt.*, 1875:112. 23. Downing, 1881:11 index, app. 24. Van Deman, *U. S. Pom. Rpt.*, 1887:630. col. pl. 25. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 26. Bailey, *Am. Hort.*, 1892:250. 27. Alwood, *Va. Sta. Bul.*, 130:122. 1901. 28. Waugh, *Vt. Sta. An. Rpt.*, 14:309. 1901. 29. Far-
rand, *Mich. Sta. Bul.*, 205:46. 1903. 30. Budd-Hansen, 1903:183. fig.

SYNONYMS. *French Reinette* (10). *Harvest Apple* (2, 3, 6). *Lippincott* (10, 13, 16). *Lodge's Early* (23). *SUMMER ROSE* (2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30). *Woolman's Harvest* (16). *Woolman's Early* (11, 14, 17). *WOOLMAN'S HARVEST* (1). *Woolman's Harvest* (8, 10, 16). *Woolman's Striped Harvest* (11).

A little dessert apple. Thomas rated it better in quality for the table than *Early Harvest* but less productive and too small for general value (11). The tree is a moderately vigorous or slow grower but is hardy, comes into bearing young and is productive. Suitable for culinary use in July, ripe in August.

Historical. This is an old New Jersey apple which Coxe described as of singular beauty and excellent for both eating and stewing; the size is moderate, the form flat, the skin smooth, of a beautiful yellow resembling wax, blended with red in streaks and blotches (2). It is still occasionally listed by nurserymen (26) but is now seldom or never planted in New York.

FRUIT (8, 11, 16, 20, 23, 30).

Fruit small to nearly medium. *Form* roundish, somewhat oblate. *Stem* rather short to medium, varying from stout to slender. *Cavity* shallow, acute, regular. *Calyx* small, closed or partly open. *Basin* regular, wide, abrupt, slightly furrowed.

Skin smooth, waxen, very pale yellow, striped and splashed distinctly with bright red and carmine on the exposed cheek. *Dots* minute.

Core medium to large; cells closed; core lines meeting. *Seeds* ovate, numerous, short, plump.

Flesh white, fine-grained, crisp, very tender, sprightly, juicy, subacid, agreeable but not rich, suitable for either culinary or dessert use.

SUMMER SPITZENBURG.

REFERENCE. 1. Downing, 1872:36 app. fig.

SYNONYMS. *French Spitzenburgh* (1). *SUMMER SPITZENBURGH* (1).

This is a September apple of attractive color. It is but little grown in New York. The tree is large, upright or roundish, a good grower, hardy,

long-lived and reliably productive yielding good crops biennially. It is not considered valuable for commercial planting because the fruit is apt to be undersized and drops badly. By some it is esteemed for home use.

Historical. Downing states that it originated with Woolsey Ostrander, Plattekill, Ulster county, N. Y. (1).

FRUIT (1).

Fruit medium, whitish almost covered with red and overspread with thin bloom. *Flesh* moderately juicy, a little aromatic, good to very good. *Season* August and September.

SUMMER SWEET.

REFERENCES. 1. *Mag. Hort.*, 14:388. 1848. *fig.* 2. Cole, 1849:97. 3. Hooper, 1857:87. 4. Downing, 1869:372.

SYNONYMS. SUMMER SWEET (2, 3, 4). SUMMER SWEETING (1).


A yellow sweet apple ripe in August and September. The tree is of medium size, moderately vigorous, spreading, productive. Fruit medium size, roundish oblate inclined to conic; flesh whitish, tender, rich, sweet (2, 4).

Historical. An old Connecticut apple (1, 2, 4) now but very seldom found in cultivation in New York.

SWEET BOUGH.

REFERENCES. 1. (?) *Amer. Gard. Cal.*, 1806:584. 2. Coxe, 1817:101. *fig.* 3. (?) Thacher, 1822:121. 4. Buel, *N. Y. Bd. Agr. Mem.*, 1826:477. 5. Wilson, 1828:136. 6. Fessenden, 1828:131. 7. Kenrick, 1832:26. 8. Floy-Lindley, 1833:84. 9. Manning, 1838:46. 10. Downing, 1845:74. 11. Hovey, *Mag. Hort.*, 14:486. 1848. *fig.* 12. Cole, 1849:99. 13. Thomas, 1849:135. 14. Phoenix, *Horticulturist*, 4:472. 1850. 15. Barry, 1851:279. 16. *Am. Pom. Soc. Cat.*, 1852. 17. Elliott, 1854:109. *fig.* 18. Hooper, 1857:20. 19. *Ib.*, 1857:107, 111. 20. Gregg, 1857:35. 21. Warder, 1867:494. *fig.* 22. Downing, 1869:250. 23. Fitz, 1872:143. 24. *Ib.*, 1872:161. 25. Leroy, 1873:154. *fig.* 26. Downing, 1881:11 index, app. 27. Hogg, 1884:129. 28. Wickson, 1889:243. 29. Bailey, *Am. Hort.*, 1892:235. 30. *Ib.*, 1892:243. 31. *Ib.*, 1892:250. 32. *Amer. Gard.*, 15:404. 1894. *fig.* 33. Taylor, *Am. Pom. Soc. Rpt.*, 1895:192. 34. Lyon, *Mich. Sta. Bul.*, 143:200, 202. 1897. 35. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 4:4. 1897. *figs.* 36. Beach, *W. N. Y. Hort. Soc. Rpt.*, 1901:76. 37. Budd-Hansen, 1903:55.

SYNONYMS. *August Sweet* (22). *August Sweeting* (12). *Autumn Bough* (11). *BOUGH* (4, 5, 11, 13, 24, 25, 29, 34). *Bough* (7, 10, 12, 17, 22, 27). *BOUGH APPLE* (2, 6, 8, 20). *BOUGH, Early Sweet* (18). *BOUGH, Sweet* (37). (*BOW APPLE* 1, 3)? *EARLY BOUGH* (7, 9). *Early Bough* (11, 25, 27). *Early French Reinette* (4). *Early Sweet Bough* (10, 11, 13, 17, 22). *Early Sweet-heart* (36). *LARGE BOUGH* (21). *LARGE EARLY BOUGH* (28). *Large Early Yellow Bough* (27). *LARGE SWEET BOUGH* (19, 30). *Large Sweet Bough* (15). *LARGE YELLOW BOUGH* (10, 16, 22, 23, 26, 27, 33). *Large Yellow*



SWEET BOUGH

Bough (11, 13, 15, 17, 20, 21, 25, 30, 31, 34, 35, 37). *Niack Pippin* (17). *Pound's July* (26). *SWEET BOUGH* (12, 14, 17, 31, 32, 35, 36). *Sweet Bough* (7, 13, 20, 22, 24, 25, 27, 33, 34). *Sweetbough* (28). *Sweet Harvest* (10, 11, 17, 22, 25, 27). *Yellow Bough* (12). *Washington* (17, 25, incorrectly 22).

This variety is a universal favorite throughout the state for the home orchard. Hovey (11) well says of it: "The Bough is one of our finest summer apples, having all the good qualities which should recommend a fruit for general cultivation. The tree is moderately vigorous, making a handsome head, and bears abundant crops of large, very fair fruit, which begins to ripen the last of July, and remains in eating till the first of September. As a table apple, it will not rank as high as the Early Harvest; but, as a kitchen fruit, in its honied sweetness and tender flesh, it has no equal of its season. It should be found in every good collection."

It is handled to a limited extent in local markets, but is too soft to stand shipping to distant markets. It cannot be ranked among the profitable commercial varieties. The tree comes into bearing rather young, and under favorable conditions is long-lived, specimens being found sixty to eighty years old which are still quite productive. In unfavorable locations the tree is sometimes injured by winter, and the branches are attacked somewhat by the apple canker.

Historical. This was described by Coxe in 1817 under the name Bough apple (2). It is evidently of American origin.

TREE.

Tree moderately vigorous. *Form* upright spreading to roundish, dense. *Twigs* short, straight, moderately stout with large terminal buds; internodes short. *Bark* clear brown mingled with olive-green, lightly mottled with scarf-skin; not pubescent. *Lenticels* scattering, small, round, not raised. *Buds* medium in size, plump, acute, free, not pubescent.

FRUIT.

Fruit above medium to large, uniform in size and shape. *Form* roundish conic or ovate to sometimes slightly oblong conic with broad and rather flat base, pretty regular; sides often unequal. *Stem* short to medium, moderately thick, usually not exerted. *Cavity* acuminate, deep, rather broad, sometimes furrowed or compressed, usually smooth. *Calyx* small to medium, closed or partly open; lobes often leafy, sometimes separated at base, long,

narrow, acute. *Basin* rather small, medium to rather shallow, narrow, a little abrupt, smooth or slightly wrinkled.

Skin rather thick, tough, smooth, pale greenish-yellow often changing to yellowish-white, sometimes faintly blushed. *Dots* numerous, small, often light colored and submerged, sometimes russet.

Calyx tube long, wide at top, conical to funnel-shape. *Stamens* median.

Core rather large to medium, abaxile to nearly axile; cells closed or open; core lines clasping. *Carpels* roundish to cordate, slightly emarginate, slightly tufted. *Seeds* light brown, medium to rather small, plump, acute.

Flesh white, moderately firm, fine, somewhat crisp, very tender, juicy, sweet, slightly aromatic, good to very good.

Season August and early September.

SWEET FALL PIPPIN.

REFERENCES. 1. Downing, 1857:192. 2. Warder, 1867:733. 3. Thomas, 1875:513.

SYNONYMS. None.

A large, greenish-yellow apple, good either for dessert or for culinary uses; in season from October to January. The tree is large, spreading, vigorous to very vigorous, hardy and a reliable cropper yielding good crops annually or nearly annually. It is not a good variety for commercial planting. Downing refers to it as being grown in Westchester county (1). It is also occasionally found in Western New York. We do not find it listed by nurserymen and it is gradually going out of cultivation.

SWEET RUSSET.

REFERENCES. 1. Warder, 1867:528. fig. 2. Downing, 1869:377. 3. Bailey, *Am. Hort.*, 1892:250.

SYNONYMS. *Summer Russet* (2). SWEET RUSSET (1, 2, 3).

A small roundish or oblate apple, yellow, mostly covered with light russet tinged with red in the sun, juicy, sweet, very good; season September and October. Tree large, spreading and drooping, a moderately vigorous grower, hardy, long-lived and usually a reliable cropper yielding good crops biennially. Origin unknown. It was at one time disseminated by Parsons and Company, Flushing, N. Y.

The name Sweet Russet has also been applied to the variety described as Pumpkin Russet on page 170.

SWITZER.

REFERENCES. 1. Hoskins, *Montreal Hort. Soc. Rpt.*, 1880:50. 2. *Ill. Hort. Soc. Rpt.*, 1881:55. 3. *Montreal Hort. Soc. Rpt.*, 1883:91. 4. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:436. 5. *Ia. Agr. Coll. Bul.*, 1885:12. No. 304. 6. *Montreal Hort. Soc. Rpt.*, 1886-87:16. 7. *Can. Hort.*, 12:344. 1889. 8. *Am. Pom. Soc. Cat.*, 1889:14. 9. Budd, *Ia. Agr. Coll. Bul.*, 1890:18. No. 304. 10.

Can. Hort., 13:216. 1890. 11. Van Deman, *U. S. Pom. Rpt.*, 1890:418. col. pl. 12. Bailey, *Am. Hort.*, 1892:251. 13. *Amer. Gard.*, 13:639. 1892. 14. *Can. Hort.*, 16:359. 1893. 15. Hoskins, *Rural N. Y.*, 53:279. 1894. 16. *Amer. Gard.*, 16:332, 412. 1895. fig. 17. Beach, *N. Y. Sta. An. Rpt.*, 14:255. 1895. 18. *Can. Hort.*, 20:183, 412. 1897. 19. *Amer. Gard.*, 19:652. 1898. 20. Macoun, *Can. Dept. Agr. Rpt.*, 1901:98. 21. Hansen, *S. D. Sta. Bul.*, 76:103. 1902. 22. Munson, *Me. Sta. An. Rpt.*, 18:85. 1902. 23. Budd-Hansen, 1903:186. fig. 24. Thomas, 1903:303.

SYNONYMS. *SUISLEPPER* (3, 4). *Suislepper* (5). *SWITZER* (1, 2, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24). *Switzer* (3, 4).

When well grown this is a very handsome fruit of medium size or below, nearly white with beautiful blush. It is very good in flavor and quality either for dessert or culinary uses. As fruited at this Station the tree comes into bearing rather early and is a fairly reliable cropper yielding pretty good crops biennially. As compared with standard varieties of its season it does not appear to merit the attention of New York fruit growers.

Historical. A Russian apple imported by the U. S. Department of Agriculture in 1870. It was received in 1888 for testing at this Station from T. H. Hoskins, Newport, Vt.

TREE.

Tree moderately vigorous with short, moderately stout, curved and crooked branches. *Form* spreading, rather flat, open. *Twigs* short, curved, stout with large terminal buds; internodes medium. *Bark* dark brown, streaked with heavy scarf-skin, slightly pubescent. *Lenticels* scattering, small, round, not raised. *Buds* large, prominent, broad, plump, obtuse, free, heavily pubescent.

FRUIT.

Fruit below medium to above medium. *Form* roundish or oblate, regular. *Stem* medium to rather long, rather slender. *Cavity* acuminate, moderately shallow, narrow, lightly russeted with thin, greenish-russet. *Calyx* small, closed; lobes medium in length, narrow, acute. *Basin* shallow or almost none, narrow to wide, furrowed, often wrinkled.

Skin clear white or becoming yellowish, washed with bright pink which often deepens to crimson. *Dots* whitish, obscure.

Calyx tube variable, elongated conical to cylindrical or funnel-form. *Stamens* median to somewhat basal.

Core large, axile; cells closed or partly open; core lines clasping. *Carpels* round, deeply emarginate. *Seeds* large, dark brown, medium in width, long.

Flesh white, firm, fine, juicy, mild subacid, good.

Season late August to October.

SYLVESTER.

- REFERENCES. 1. *Horticulturist*, 17:150. 1862. 2. Warder, 1867:617. fig. 3. Downing, 1869:379. 4. *Ill. Hort. Soc. Rpt.*, 1871:154. 5. Fitz, 1872:170. SYNONYMS. None.

A waxen-white fruit with crimson blush and brisk subacid flesh; very good for culinary uses. Season September and October. The tree is large, roundish, moderately vigorous, hardy, long-lived and a reliable cropper yielding good crops biennially. It is not desirable for commercial planting because the fruit shows bruises very readily.

Historical. Originated at Lyons, N. Y. (3). It is but little grown even in the locality of its origin.

TART BOUGH.

- REFERENCES. 1. Thomas, 1849:142. 2. Elliott, 1854:178. 3. Warder, 1867:734. 4. Downing, 1869:380.

SYNONYMS. *Sour Bough* of some (4). TART BOUGH (1, 2, 3, 4).

An old variety which according to Downing (4) was originated and disseminated by Judge Buel of Albany, N. Y. Elliott (2) states that it resembles Early Harvest, but it ripens ten days later, has more acid and the trees are more rapid in growth. He regarded it as unworthy of cultivation.


Downing recognizes another Tart Bough the fruit of which is small and whitish with white, tender flesh, juicy, sprightly, pleasant subacid, good. Season August (4).

We do not know either of these varieties.

TETOFSKY.

- REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 1291. 2. Manning, *Mag. Hort.*, 7:50. 1841. 3. Downing, 1845:78. 4. Thomas, 1849:141. 5. Elliott, 1854:111. fig. 6. Hooper, 1857:93. 7. *Horticulturist*, 14:425. 1859. 8. *Am. Pom. Soc. Cat.*, 1862. 9. *Horticulturist*, 21:294. 1866. fig. 10. Warder, 1867:657. 11. Todd, 1871:120. fig. 12. *Mich. Pom. Soc. Rpt.*, 1872:458. fig. 13. *Horticulturist*, 29:317. 1874. 14. *Montreal Hort. Soc. Rpt.*, 1876:18. 15. *Ill. Hort. Soc. Rpt.*, 1876:251. 16. *Montreal Hort. Soc. Rpt.*, 1879:22. 17. Barry, 1883:335. 18. *Montreal Hort. Soc. Rpt.*, 1886-87:27. 19. Hoskins, *Rural N. Y.*, 45:593. 1886. 20. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 21. Bailey, *Am. Hort.*, 1892:251. 22. *N. Y. Sta. An. Rpt.*, 11:588. 1892. 23. Munson, *Me. Sta. Rpt.*, 1896:76. 24. *Can. Hort.*, 19:358. 1896. 25. Waugh, *Vt. Sta. Bul.*, 61:32. 1897. 26. Troop, *Ind. Sta. An. Rpt.*, 12:81. 1899. 27. Waugh, *Vt. Sta. An. Rpt.*, 14:310. 1901. 28. Craig, *Cyc. of Hort.*, 1901:1404. 29. Hansen, *S. D. Sta. Bul.*, 76:104, 132. 1902. fig. 30. Munson, *Me. Sta. Rpt.*, 1902:85. 31. Dickens and Greene, *Kan. Sta. Bul.*, 106:56. 1902. 32. Budd-Hansen, 1903:188. fig.

SYNONYMS. TETOFSKY (2). TETOFSKI (10, 12, 27, 29). TETOFSKY (1, 3, 4, 5, 6, 7, 8, 9, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 28, 30, 31, 32). *Tetofsky* (29).



TETOFKY

A Russian apple below medium size, yellow, striped with crimson, sprightly acid, suitable for culinary use in July, ripe in August. The tree is very hardy, very upright, deep rooted, comes into bearing young and yields full crops biennially or sometimes annually. Desirable for planting in regions where superior hardiness is particularly desired. It is but little earlier than Yellow Transparent which is superior to it in size and dessert qualities. There is a limited demand for the fruit in some local and special markets.

Historical. Tetofsky is one of the pioneers of the Russian varieties in this country having been imported by the Massachusetts Horticultural Society from the London, England, Horticultural Society about 1835 (28). Manning (2) described it as early as 1841 as a new fruit of value. It has been widely disseminated particularly in regions where its superior hardiness renders it peculiarly valuable. It has never been commonly grown in New York state and is now seldom planted here.

TREE.

Tree medium size to dwarfish, moderately vigorous, deep rooted with moderately stout branches filled with small spurs. *Form* very erect, rather dense. *Twigs* short, straight, moderately stout; internodes short. *Bark* dull brown with tinge of red, lightly coated with gray scarf-skin, slightly pubescent. *Lenticels* scattering, small, oblong, not raised. *Buds* small, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium to rather small but usually below medium; uniform in size and fairly uniform in shape. *Form* oblate or roundish, a little inclined to conic, pretty regular. *Stem* medium in length, rather slender. *Cavity* obtuse, medium to rather deep, moderately broad, usually symmetrical, gently furrowed. *Calyx* rather large, closed; lobes long, moderately broad. *Basin* shallow, medium in width, furrowed and wrinkled.

Skin moderately thick, somewhat tough, smooth, waxy, greenish-yellow, more or less striped and splashed with rather attractive bright red, overspread with very faint bloom. Many specimens have little or no red. *Dots* inconspicuous, pale or greenish, submerged.

Calyx tube long, wide, funnel-shape with broad cylinder varying to nearly urn-shape. *Stamens* median to nearly marginal.

Core large, axile or somewhat abaxile; cells open; core lines clasping. *Carpels* broadly roundish, concave. *Seeds* medium size, short, moderately plump, somewhat obtuse.

Flesh white, firm, a little coarse, crisp, tender, juicy, sprightly, slightly aromatic, subacid, fair to good.

Season late July to early September.

THALER.

REFERENCES. 1. Webster, *Montreal Hort. Soc. Rpt.*, 1881:54. 2. Tuttle, *Ib.*, 8:135. 1881-82. 3. *Ib.*, 1883:94. 4. Gibb, *la. Hort. Soc. Rpt.*, 1883:438. 5. Budd, *Ia. Agr. Coll. Bul.*, 1885:14. 6. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:14. 7. Schroeder, *Ib.*, 1886-87:79. 8. Bailey, *An. Hort.*, 1892:236. 9. *Ib.*, 1892:251. 10. Green, *Minn. Sta. Bul.*, 32:240. 1893. 11. Jack, *Gard. and For.*, 7:99. 1894. 12. Hoskins, *Ib.*, 7:138. 1894. 13. *Rural N. Y.*, 54:539. 1895. 14. Troop, *Ind. Sta. Rpt.*, 1899:81. 15. Hansen, *S. D. Sta. Bul.*, 76:38. 1902.

SYNONYMS. CHARLOTTENTHALER (2, 6, 7, 8, 11, 15). *Charlottenthaler* (3, 4). CHARLOTTEN THALER (13). *Charlotten Thaler* (10). *Charlottenthaler Apple* (1). *Charlottenthaler Golba* (5). No. 147 (7). SCHARLOTTENTHALER GOLBA (1, 3, 4). THALER (5, 9, 10, 12, 14). *Yellow Transparent* (11).

A Russian variety of the Yellow Transparent type. It resembles the Yellow Transparent so closely that some have considered them identical but they are distinct (6, 12). Since Yellow Transparent is superior in health, vigor and productiveness Thaler is not recommended for planting.

THOMPSON.

REFERENCES. 1. Hansen, *S. D. Sta. Bul.*, 76:104. 1902. 2. Munson, *Me. Sta. Rpt.*, 1902:85.

SYNONYMS. THOMPSON (1). THOMPSON No. 24 (2). *Thompson's Seedling No. 24* (1).

Fruit of pretty good size, striped red and yellow, rather attractive. Flesh subacid, fair or possibly good in quality. Season late winter. The tree is a pretty good grower, comes into bearing rather young and bears nearly annually yielding moderately good crops. This is a new variety which has gained recognition in the upper Mississippi valley on account of its hardiness in that region. It is also reported as promising in the Northern apple districts of Maine (2). It is not worthy of planting in this state where standard varieties of its season succeed but may be worth testing in those localities where a tree of superior hardiness is desired.

Historical. Originated by J. S. B. Thompson, Grundy county, Ia., and first propagated under the name Thompson Seedling No. 24 (1). In 1892 it was received for testing at this Station from the Jewell Nursery Company, Lake City, Minn., by which the variety was introduced.

TREE.

Tree moderately vigorous with moderately long, slender, curved, drooping branches. *Form* spreading, open, inclined to droop. *Twigs* moderately long, curved, moderately stout; internodes medium. *Bark* brown tinged with red, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* numerous, medium size, oval, raised, conspicuous. *Buds* medium size, broad, obtuse to acute, free, slightly pubescent.

FRUIT.

Fruit below medium to above or rather large. *Form* roundish or roundish oval, pretty regular but somewhat unequal. *Stem* medium in length, slender. *Cavity* acuminate to acute, deep, medium in width to rather narrow, more or less russeted. *Calyx* medium size, open; lobes separated at base. *Basin* deep, narrow to rather wide, abrupt, somewhat furrowed.

Skin pale yellow more or less overspread and mottled with pinkish-red striped and splashed with carmine. *Dots* grayish or brownish, few, small, inconspicuous.

Calyx tube short, conical to funnel-form. *Stamens* basal to median.

Core medium size, abaxile; cells fairly symmetrical, open; core lines nearly meeting. *Carpels* nearly cordate to elliptical. *Seeds* numerous, medium size, flat, moderately acute.

Flesh white, moderately fine, crisp, rather tender, juicy, subacid to mild subacid, fair to good.

Season October to early winter or midwinter.

TINMOUTH.

REFERENCES. 1. Downing, 1857:197. 2. *Am. Pom. Soc. Cat.*, 1862. 3. Downing, *Horticulturist*, 19:145. 1864. figs. 4. Warder, 1867:734. 5. Downing, 1869:383. 6. Thomas, 1875:514. 7. Taylor, *Am. Pom. Soc. Rpt.*, 1895:193. 8. Burrill and McCluer, *Ill. Sta. Bul.*, 45:343. 1896. 9. Waugh, *Vt. Sta. An. Rpt.*, 14:310. 1901.

SYNONYMS. TEIGNMOUTH (3). Teignmouth (5, 6). TINMOUTH (1, 2, 4, 5, 6, 7, 8, 9). Tinnmouth Sweet (8). Vermont Pippin (3, 5).

Tinmouth is a fall and early winter apple of good size, good quality and rather attractive appearance. The tree is a good grower, with an upright habit. It is a regular cropper and very productive. It does not seem to be desirable for general commercial planting in New York.

Historical. Downing remarked that this variety originated at Tinmouth, Vt., and was much esteemed in the region of its origin (5). Waugh calls it a good apple and states that it is almost unknown in Grand Isle county but is much more common in other parts of Vermont (9). So far as we have discovered it is but little known in New York.

FRUIT.

Fruit medium to large. *Form* oblate to roundish, regular or obscurely ribbed; sides often distinctly unequal. *Stem* often bracted, short to medium, usually not exerted. *Cavity* large, acute or approaching acuminate, deep, broad, russeted and with outspreading russet rays. *Calyx* small, partly closed or open; lobes often small, sometimes long, acute, recurved. *Basin* rather large, deep, moderately wide, abrupt to moderately abrupt, often somewhat wrinkled.

Skin tender, pale yellow or greenish, often with a bright deep blush and overspread with thin bloom. *Dots* numerous, greenish or russet, giving the surface a somewhat rough appearance. *Prevailing effect* yellow.

Calyx tube rather small, cone-shape or approaching funnel-form. *Stamens* median.

Core rather small, axile or nearly so; cells closed; core lines clasping. *Carpels* roundish to obcordate, emarginate, mucronate. *Seeds* rather large, broad, somewhat obtuse.

Flesh whitish tinged with yellow, moderately fine, tender, rather juicy, mild subacid with a peculiar flavor, sprightly, good in quality.

Season October to early winter or midwinter.

TITOVKA.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 8:44. 1881-82. fig. 2. *Ib.*, 8:74. 1881-82. 3. *Ia. Hort. Soc. Rpt.*, 1882:80. 4. Budd, *Ia. Agr. Coll. Bul.*, 1883:31. 5. Gibb, *Ia. Hort. Soc. Rpt.*, 1883:431. 6. *Mo. Hort. Soc. Rpt.*, 1883:173. 7. Barry, 1883:335. 8. Budd, *Ia. Agr. Coll. Bul.*, 1885:8. 9. Thomas, 1885:526. 10. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:15. 11. Schroeder, *Ib.*, 1886-87:78. 12. Hoskins, *Can. Hort.*, 13:175. 1890. 13. Bailey, *An. Hort.*, 1892:251. 14. Taylor, *Me. Pom. Soc. Rpt.*, 1892:58. 15. Troop, *Ind. Sta. Bul.*, 53:124. 1894. 16. Beach, *N. Y. Sta. An. Rpt.*, 13:584. 1894. 17. Munson, *Me. Sta. Rpt.*, 1896:77. 18. *Mass. Hatch Sta. Bul.*, 44:4. 1897. 19. *Am. Pom. Soc. Cat.*, 1897:15. 20. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 6:11. 1899. figs. 21. Troop, *Ind. Sta. Rpt.*, 1899:81. 22. Eneroth-Smirnoff, 1901:470. 23. Hansen, *S. D. Sta. Bul.*, 76:105. 1902. fig. 24. Munson, *Me. Sta. An. Rpt.*, 18:85. 1902. 25. Budd-Hansen, 1903:188. fig. 26. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:58. 1903. 27. Far-
rand, *Mich. Sta. Bul.*, 205:46. 1903.

SYNONYMS. No. 134 (11). No. 230 Gov. (4). ТИТОВКА (9). ТИТОВКА (1, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27). ТИТОВКА (5). ТИТУС APPLE (2, 3, 4). *Titus Apple* (1, 5, 7). *Titus Riga* (25).

A Russian apple, large, roundish or somewhat oblong, greenish-yellow, handsomely shaded and striped with red and covered with light bloom. When fully mature the ground color is yellow and the red is bright and dark often nearly covering the fruit. Quality good to very good for culinary use. Season August and September. The tree is a vigorous grower, comes into bearing rather young and is moderately productive. It is one of the best Russian apples of its season which we have tested. The crop does not ripen uniformly and more than one picking is required in order to secure the fruit in prime condition for market. Titovka is perhaps worthy of testing for market where fruit of this type and season is desired.

This appears to be the variety which Hansen describes under the name Titovka Department, which name he adopts to distinguish it on the one hand from a variety called Titovka with Titus Riga as a synonym and on the other from a variety called Titovka Speer (23).

Historical. Imported from Russia by the U. S. Department of Agriculture in 1870. It was received in 1883 from Ellwanger and Barry, Rochester, N. Y., for testing at this Station. In 1897 it was entered in the catalogue of the American Pomological Society as a variety of value in the pomological district which includes the Dakotas, Montana and Wyoming (19). It is but little known in New York and so far as we can learn its cultivation is not increasing in this state.

FRUIT.

Fruit large. *Form* roundish or somewhat oblong, a little inclined to conic, nearly regular, pretty symmetrical. *Stem* short, usually not exerted. *Cavity* acuminate, very deep, sometimes slightly russeted. *Calyx* medium size, closed; segments small, convergent. *Basin* moderately shallow to deep, very abrupt, wrinkled.

Skin smooth, pale green becoming yellow, shaded and striped with bright red and overspread with light bloom.

Calyx tube elongated cone-shape or funnel-form. *Stamens* median or below.

Core rather large, somewhat abaxile; cells pretty symmetrical, open; core lines clasping. *Carpels* broadly roundish or somewhat obovate, emarginate. *Seeds*, few, rather small, plump.

Flesh whitish, coarse, crisp, juicy, subacid, good to very good for culinary uses.

Season August and September.

TOM PUTT.

REFERENCES. 1. *London Hort. Soc. Cat.*, 1831:No. 1299. 2. Hogg, 1884: 229.

SYNONYMS. *Coelbrook* (2). *Marrow-bone* (2). *TOM PUT* (1). *TOM PUTT* (2).

A rather attractive red apple of good size, moderately coarse, subacid, fair to good for culinary uses. The tree is a moderately vigorous grower, comes into bearing rather early and yields full crops annually or nearly annually. As compared with standard varieties of its season it does not appear to be worthy of the attention of New York fruit growers.

Historical. This is an old English variety (1, 2). In 1892 it was received for testing here from W. and T. Smith, Geneva, N. Y.

TUFTS.

REFERENCES. 1. *Mag. Hort.*, 14:519. 1848. 2. Cole, 1849:107. 3. Hovey, *Mag. Hort.*, 16:496. 1850. fig. 4. Hovey, 2:23. 1851. col. pl. 5. Downing, 1857:198. 6. Warder, 1867:734. 7. Thomas, 1885:527. 8. Bailey, *Am. Hort.*, 1892:251. 9. *Ib.*, 1892:251. 10. Beach, *N. Y. Sta. An. Rpt.*, 11:589, 596. 1892. 11. Burrill and McCluer, *Ill. Sta. Bul.*, 45:344. 1896. 12. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:58. 1903. 13. Beach and Clark, *N. Y. Sta. Bul.*, 248:147. 1904.

SYNONYMS. TUFTS (3, 4, 7, 8, 10, 11, 12, 13). TUFTS BALDWIN (2, 5, 6, 9). *Tufts Baldwin* (7, 10, 12, of some 3 and 5). TUFTS SEEDLING (1).

An apple of the Baldwin group, somewhat like Baldwin in form, color and general appearance, but more mild in flavor and less desirable for market or culinary uses. In ordinary storage it is in season from October to January with October as the commercial limit, but in cold storage it may be held through the winter (13). In some years nearly all of the fruit is discolored at the core. The crop does not ripen evenly, and there is apt to be considerable loss from dropping of the fruit. The tree is large, a pretty good grower, comes into bearing rather early and yields full crops biennially. Not recommended for planting in New York.

Historical. This variety originated in Cambridge, Mass., about 1830. It was first exhibited at the annual show of the Massachusetts Horticultural Society in 1848 (4). It is now rarely listed by nurseymen (8, 9) and is seldom or never planted in this state.

TREE.

Tree large, moderately vigorous. *Form* very spreading, flat, open. *Twigs* moderately long, curved, slender; internodes short. *Bark* brown, heavily coated with gray scarf-skin; pubescent. *Lenticels* scattering, medium size, oblong, not raised. *Buds* medium size, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit large or nearly so, quite uniform in size and shape. *Form* roundish to roundish oblate, sometimes with a broad protuberance above the cavity on one side, often flattened at apex, broadly ribbed. *Stem* long, rather slender. *Cavity* acute to acuminate, medium in depth to deep, rather broad, often furrowed, usually with outspreading, irregular, greenish-russet. *Calyx* medium to rather small, usually closed. *Basin* rather shallow to moderately deep, medium in width to wide, obtuse to somewhat abrupt, slightly wrinkled, somewhat ridged.

Skin moderately thin, tough, smooth, green or yellowish nearly covered with bright deep red like the Baldwin, not striped or with indistinct stripes of purplish carmine. *Dots* conspicuous, medium to rather small, gray or russet.

Calyx tube medium in length, conical or funnel-shape. *Stamens* median to marginal.

Core large, axile or nearly so; cells closed or partly open; core lines clasping. *Carpels* broadly roundish, often discolored, rather flat, emargin-

ate. *Seeds* often few, not well developed, dark brown, large, long, somewhat acute, slightly tufted.

Flesh tinged with yellow or greenish, firm, moderately coarse, crisp, rather tender, moderately juicy, rather mild subacid, fair to good.

Season October to December or January.

TWENTY OUNCE.

REFERENCES. 1. *Mag. Hort.*, 10:210. 1844. 2. Downing, 1845:140. 3. Hovey, *Mag. Hort.*, 13:70. 1847. fig. 4. Thomas, 1849:153. 5. Cole, 1849:120. 6. Elliott, 1854:126. 7. Hooper, 1857:94. 8. Downing, 1857:198. 9. *Am. Pom. Soc. Cat.*, 1862. 10. Warder, 1867:510. fig. 11. Downing, 1869:388. 12. Fitz, 1872:168. 13. Leroy, 1873:261. fig. 14. Barry, 1883:356. 15. Wickson, 1889:245. 16. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 17. Bailey, *Am. Hort.*, 1892:236. 18. *Id.*, 1892:251. 19. Budd-Hansen, 1903:191. 20. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:59. 1903. 21. Beach and Clark, *N. Y. Sta. Bul.*, 248:147. 1904.

SYNONYMS. *Aurora* (8, 11, 13). *Cabashaw*, incorr. (21). *CAYUGA RED STREAK* (6, 10, 17). *Cayuga Red Streak* (3, 5, 8, 11, 12, 13, 14, 15, 16, 19, 20, 21, 22). *Coleman* (8, 11, 13). *De l'in du Conn.* (13). *DIX-HUIT ONCES* (2, 13). *Eighteen Ounce* (4, 6, 13). *Eighteen Ounce Apple* (2, 8, 11). *Gov. Seward's* (3, 6). *Lima* (8, 11, 13). *Morgan's Favorite* (8, 11, 13). *TWENTY OUNCE* (1, 2, 3, 4, 5, 7, 8, 9, 11, 12, 14, 15, 16, 18, 19, 20, 21). *Twenty Ounce* (6, 10, 13). *Twenty Ounce Apple* (2, 3, 6, 8, 11, 12). *Twenty Ounce Pippin* (3, err. 4 and 6). *Wine* (21, of Conn. 11).

This is one of the most satisfactory of the fall varieties for commercial planting in New York. It is also highly esteemed for home use. The fruit is large, attractive, green becoming yellowish with broad stripes and splashes of red. It is in season from September to early winter. It keeps well for a fall variety and stands shipping well. Usually it should be handled direct to the consumer without going into cold storage. In common storage the fruit goes down rapidly during October and November. In cold storage it may be kept till midwinter (21). It hangs pretty well to the tree for so large an apple, is pretty uniform in size and generally reliable and satisfactory in color and quality. The fruit is in good demand in general market and sells at good prices. It is especially esteemed for culinary uses but it is inferior to other varieties for evaporating. The tree is a rather vigorous grower with main branches erect and laterals rather willowy and more or less drooping. It seems to succeed particularly well in favorable locations in the

apple belt south of Lake Ontario. It is especially subject to sunscald and canker on the trunk and larger limbs. For this reason it is desirable to topwork it upon some hardy and thrifty stock such as Tolman *Sweet* or Northern Spy. Careful attention should be given to treating the canker.¹ The top is inclined to grow rather dense and requires constant attention to keep it properly pruned to admit the light to the foliage in all parts of the tree so that the fruit may color properly. The tree when full grown is of medium size or below medium and may be planted closer in the orchard than Baldwin, Rhode Island *Greening* or Northern Spy. It comes into bearing rather young and is almost an annual bearer yielding moderate to good or even heavy crops. The skin of the fruit is apt to be roughened by the application of spray mixtures. In spraying Twenty Ounce after the fruit is set, it is therefore desirable to use an abundance of lime in the bordeaux mixture and make the application uniform and thorough but not excessive.

In different localities in New York Twenty Ounce is known under the various synonyms of Cayuga Redstreak, Wine Apple and Limbertwig. It is quite distinct from Twenty Ounce Pippin; see Volume I, page 349.

Historical. This variety was brought to the notice of pomologists about sixty years ago (1, 3), when fruit of it grown in Cayuga county, N. Y., was exhibited before the Massachusetts Horticultural Society. At that time its cultivation appeared to be mostly confined to Cayuga county and its origin was unknown. In 1857 Downing (8) reported that it originated in Connecticut but upon what authority we are unable to state. Twenty Ounce is well known in most of the apple-growing sections of the state and in certain districts its cultivation for commercial purposes appears to be increasing. It is pretty generally listed by nurserymen (17).

TREE.

Tree moderately vigorous with branches moderately long and moderately stout. *Form* upright becoming roundish, dense; laterals willowy, slender and more or less drooping. *Twigs* short, straight, slender; internodes medium. *Bark* reddish-brown tinged with green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* quite numerous, medium size, round, not raised. *Buds* medium size, broad, flat, obtuse, appressed, pubescent.

¹ N. Y. Sta. An. Rpt., 18:399-418. 1899. *Ib.*, 22:321-386. 1903.

PLANT. ANTH.



FRUIT.

Fruit very large. *Form* variable, usually roundish or roundish conic, sometimes broadly ribbed. *Stem* deep set, short to medium, moderately thick or rather slender. *Cavity* acuminate, very deep, sometimes lipped, sometimes russeted. *Calyx* below medium size to above, usually closed. *Basin* often oblique, often shallow but occasionally deep, moderately narrow to rather wide, rather abrupt, broadly and deeply furrowed.

Skin thick, tough, greenish becoming rather yellow, washed, mottled and splashed with bright red or deepening to dark or purplish-red with carmine stripes. *Dots* grayish or russet, small to large, often raised, sometimes whitish and submerged.

Calyx tube large, long, wide, conical or funnel-shape extending to the core. *Stamens* usually basal.

Core large, axile or somewhat abaxile; cells symmetrical, usually closed, sometimes wide open; core lines clasping the funnel cylinder. *Carpels* elongated ovate, slightly emarginate, somewhat tufted. *Seeds* medium size, round to narrow, obtuse to acute, variable.

Flesh whitish somewhat tinged with yellow, coarse, moderately tender, juicy, subacid, good for culinary use, second rate for dessert.

Season late September to early winter.

RED TYPE OF TWENTY OUNCE.

A red Twenty Ounce which originated at Hilton, Monroe county, N. Y., is being propagated under the name "Collamer." See page 36.

TYRE BEAUTY.

REFERENCES. 1. Downing, 1869:388. 2. *Horticulturist*, 24:52. 1869. fig. SYNONYMS. None.

This was brought to notice about thirty-five years ago as a new seedling apple of value in the locality of its origin, Tyre, Seneca county, N. Y. Fruit medium, pale yellow, splashed, marbled and shaded with crimson. Flesh tender, slightly acid, neither rich nor peculiarly sprightly. Season early September (2).

We are unacquainted with this variety and so far as we know it has not been grown outside of the locality of its origin.

UTTER.

REFERENCES. 1. *Horticulturist*, 10:528. 1855. 2. *Ib.*, 23:10. 1868. fig. 3. Downing, 1869:389. 4. *Am. Pom. Soc. Cat.*, 1873. 5. *Montreal Hort. Soc. Rpt.*, 5:35. 1879. 6. Budd, *1a. Hort. Soc. Rpt.*, 1879:472. 7. Thomas, 1885:527. 8. *Montreal Hort. Soc. Rpt.*, 1886:87:97. 9. Bailey, *Am. Hort.*, 1892:251. 10. Harris, *U. S. Pom. Rpt.*, 1892:271. 11. Burrill and McCluer, *Ill. Sta. Bul.*, 45:311, 344. 1896. 12. Macoun, *Can. Dept. Agr. Rpt.*, 1901:98. 13. Hansen, *S. D. Sta. Bul.*, 76:108. 1902. fig. 14. Budd-Hansen, 1903:193. fig.

SYNONYMS. *English Janneting* (6). *Fameuse* (6). *Seever's Red Streak* (6). *UTTER'S* (1). *UTTER* (2, 3, 4, 6, 7, 8, 9, 10, 11, 13, 14). *UTTER'S LARGE RED* (9). *UTTER'S RED* (5, 12). *Utter's Red* (13, 14).

Fruit of good size, yellow and red, rather attractive in appearance. The tree is a rather upright regular grower, forming a full rounded head, healthy and productive (11). It is very hardy and on this account has been grown to some extent in regions where standard varieties do not succeed (13, 14).

Historical. Originated in Wisconsin where it was known as early as 1855 (1). It has been much grown in that state and in other parts of the Middle West (13, 14) but it is little known in New York.

FRUIT.

Fruit above medium to large. *Form* usually roundish oblate varying to roundish, sometimes a little inclined to oblong, often somewhat broadly ribbed, pretty regular. *Stem* short to medium, moderately stout. *Cavity* acute to acuminate, deep, medium in width to wide, furrowed gently if at all, sometimes partly russeted. *Calyx* small, closed or partly open; lobes small, short, obtuse. *Basin* rather shallow to moderately deep, medium in width, abrupt, slightly furrowed and wrinkled.

Skin moderately thick, tough, clear, rather pale yellow usually washed with orange-red and narrowly streaked with bright carmine. Some fruits show little or no red but on highly colored specimens the prevailing color is red. *Dots* not conspicuous, numerous, often submerged or whitish or occasionally with russet point.

Calyx tube elongated funnel-form. *Stamens* median or below.

Core sessile, medium size, abaxile; cells symmetrical, open or closed; core lines clasping. *Carpels* broadly roundish or approaching elliptical, but slightly emarginate if at all, smooth or slightly tufted. *Seeds* moderately numerous, medium to rather large, somewhat narrow to rather wide, obtuse or approaching acute, moderately light reddish-brown.

Flesh whitish tinged with yellow, somewhat coarse, crisp, tender, juicy, mild subacid, pleasantly flavored, good.

Season October to December or later.

VANDEVERE.

REFERENCES. 1. *Am. Gard. Cal.*, 1806:585. 2. *Coxe*, 1817:141. *fig.* 3. *Thacher*, 1822:139. 4. *Buel, N. Y. Bd. Agr. Mem.*, 1826:477. 5. *Floyd-Lindley*, 1833:85. 6. *Downing*, 1845:141. *fig.* 7. *Thomas*, 1849:173. 8. *Cole*, 1849:122. 9. *Horticulturist*, 10:87. 1855. 10. *ib.*, 11:89. 1856. 11. *Warder*, 1867:735. 12. *Fitz*, 1872:121, 149, 153. 13. *Am. Pom. Soc. Cat.*, 1873. 14. *Lyon, Mich. Hort. Soc. Rpt.*, 1890:298. 15. *Bailey, An. Hort.*, 1892:251. 16. *Budd-Hansen*, 1903:193.

SYNONYMS. *Oxeye* (8, 12). *Staalcubs* (2, 4). *Stalcubs* (3, 6). *VANDERVERE* (2, 6, 9, 10, 11). *VANDEVEER* (4). *VANDEVERE* (1, 3, 5, 7, 8, 12, 13, 14, 15, 16).

The old variety which is commonly called Vandevere in New York is known to pomologists by the name Newtown Spitzenburg, under which name it is described in Vol. I, page 225. It originated in Newtown, L. I. The true Vandevere which we are here considering is quite distinct from this Newtown Spitzenburg.

When in perfection Vandevere is a beautiful and fine apple, medium in size, marked with light red in indistinct streaks over a yellow background; well colored specimens become deep red; dots numerous, green or light gray; flesh yellowish, crisp and tender with a rich, sprightly, mild subacid flavor, valued especially for culinary purposes; in season from October to January. The tree is of medium size, spreading, moderately vigorous, not very productive (2, 6, 7).

Historical. An old variety native of Wilmington, Del. (2, 6). It is sometimes called the Vandevere of Delaware or the Vandevere of Pennsylvania. It has never been much cultivated in New York and is now seldom or never planted here.

VANDEVERE PIPPIN.

REFERENCES. 1. Phoenix, *Horticulturist*, 4:471. 1849. 2. Elliott, 1854:113. fig. 3. Downing, 1857:199. 4. Hooper, 1857:94. 5. Warder, 1867:462. 6. Thomas, 1875:204. 7. Budd-Hansen, 1903:193. fig.

SYNONYMS. *Baltimore* of some, incorrectly (2). *Big Vandevere* (3). *Fall Vandevere* (2). *Gibbon's Smokehouse?* (2). *Gibbon's Smokehouse?* (2). *Imperial Vandevere* (2). *Indiana Vandevere* (3, 6). *Large Vandevere* (5). *Millcreek* (2). *Millcreek Vandevere* (2). *Pennsylvania Vandevere* (2). *Red Vandevere* (2). *Smokehouse?* (2). *Spiced Oxye* (2). *Staalclubs* (2). *Striped Ashmore?* (2). *Striped Vandevere* (2). *VANDEVERE* (2). *Vandevere* (5). *VANDEVERE PIPPIN* (1, 5). *Vandevere Pippin* (2). *VANDEVERE PIPPIN* (3, 6, 7). *Vandevere Pippin* (4). *VANDEVERE YELLOW* (4). *Watson's Vandevere* (2, 5). *Watson's Vandevere* (3, 6). *Windower* (1). *Yellow Vandevere* (2, 5).

A large, coarse apple, yellow, more or less covered with marbled red and scarlet stripes; flesh of rather sharp acid flavor, excellent for culinary use but not esteemed for dessert (5); in season from September or October to early winter. The tree is vigorous, large, spreading, a reliable cropper and productive. The twigs and leaves much resemble those of Vandevere (5). It appears that it is no longer listed by nurserymen.

Historical. Origin unknown (3, 5, 7). It has been grown to some extent in the West but has never been much cultivated in New York.

VICTORIA.

REFERENCES. 1. Downing, 1881:111 app. fig. 2. Bailey, *An. Hort.*, 1892: 251. 3. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:59. 1903. 4. Beach and Clark, *N. Y. Sta. Bul.*, 248:148. 1904.

SYNONYMS. *VICTORIA* (3, 4). *VICTORIA SWEET* (1). *Victoria Sweet* (4). *VICTORIA SWEETING* (2).

This variety belongs in the same group with *Mabie*. The fruit is of good medium size, dark red, with conspicuous, large dots somewhat like those seen on *Westfield Seek-No-Further* and *Blue Pearmain*. The flesh is moderately coarse, very tender, rather juicy, sweet, good to very good, for either dessert or culinary uses. In ordinary storage it is in season from October to January, with October as the commercial limit; in cold storage it may be held till January (4). The tree is a pretty good grower, comes into bearing rather young and yields full crops biennially. It is recommended for trial in Central and Eastern New York where a variety of this type is desired either for the home or for the local market.

Historical. Origin uncertain. It is supposed by some to have originated in Chenango county. Probably it is nowhere grown extensively but it is cultivated more in Chenango and adjacent counties than in any other region. It is occasionally listed by nurserymen (2).

TREE.

Tree moderately vigorous with rather short, rather stocky, crooked branches. *Form* spreading. *Twigs* moderately long, rather slender; internodes medium. *Bark* dull reddish-brown or olive-green, slightly mottled with scarf-skin; pubescent. *Lenticels* scattering, usually large and elongated. *Buds* medium size, rather narrow, plump, appressed, acute, pubescent. *Leaves* medium size, rather broad.

FRUIT.

Fruit above medium to medium, fairly uniform in size and shape. *Form* roundish inclined to conic, somewhat flattened at base, faintly and broadly ribbed. *Stem* usually short, moderately thick. *Cavity* moderately deep, medium in width to broad, symmetrical, often lipped, sometimes red and smooth, but often bright yellow russet or greenish-russet overspreads the cavity and radiates irregularly over the base in broken lines and splashes. *Calyx* small to medium, closed or partly open; lobes usually short and not separated at base, acute. *Basin* medium in depth to moderately deep, narrow to wide, somewhat abrupt, slightly wrinkled.

Skin tough, nearly smooth, yellow, blushed and faintly mottled with rather dull red and marked with numerous, narrow stripes of deeper red. Highly colored specimens are purplish-red with obscure stripes. *Dots* or flecks conspicuous, gray or russet, becoming smaller and more numerous toward the cavity.

Calyx tube short, medium size, conical or somewhat funnel-form. *Stamens* median to basal.

Core medium to somewhat distant, usually abaxile; cells closed or partly open; core lines clasping the funnel cylinder. *Carpels* roundish cordate to



VICTORIA SWEET

elliptical, decidedly concave, tufted. *Seeds* numerous, dark, medium size, flat, acute to obtuse.

Flesh tinged with yellow, firm, moderately coarse, very tender, juicy, sweet, good to very good.

Season October to January.

VICTUALS AND DRINK.

REFERENCES. 1. Downing, 1845:141. 2. Thomas, 1849:163. 3. Emmons, *Nat. Hist. N. Y.*, 3:88. 1851. 4. Elliott, 1854:179. 5. Hooper, 1857:94. 6. Warder, 1867:499. *fig.* 7. *Am. Pom. Soc. Cat.*, 1873. 8. Taylor, *Am. Pom. Soc. Rpt.*, 1895:200.

SYNONYMS. *Big Sweet* (1, 4, 5). *Fall Green Sweet* (6). *Green Sweet* of Indiana (6). *Pompey* (1, 4, 5, 6). VICTUALS AND DRINK (1, 2, 3, 4, 5, 6, 7, 8).

A large, somewhat rough, dull green or yellowish apple often veined with russet. The flesh is sweet, very tender, fine-grained, very good to best in quality; in season from October to January or later. The tree is medium to rather large, upright or roundish, stocky, vigorous, very productive, often carrying so heavy a load of fruit that many of the apples are small. Downing states that it originated in the neighborhood of Newark, N. J., about 1750 (1). In 1873 (7) it was entered in the catalogue of the American Pomological Society but was dropped from that list in 1897. Bailey does not mention it in his *Inventory of Apples Offered by American Nurserymen* in 1892.¹ It has been popular in some portions of the West but so far as we know has never been much cultivated in New York. It is undoubtedly an excellent variety for the home orchard.

VINEUSE ROUGE.

REFERENCES. 1. Leroy, 1873:230. *fig.* 2. *Ib.*, 1873:846. *fig.* 3. Hoskins, *la. Hort. Soc. Rpt.*, 1879:414. 4. Gibb, *Ib.*, 1883:442. 5. *N. Y. Sta. An. Rpt.*, 3:20. 1884. 6. Budd, *la. Agr. Coll. Bul.*, 1885:16. 7. Craig, *Montreal Hort. Soc. Rpt.*, 12:102. 1886-87. 8. Beach, *N. Y. Sta. An. Rpt.*, 11:587. 1892. 9. Bailey, *Am. Hort.*, 1892:240. 10. Beach, *N. Y. Sta. An. Rpt.*, 12:600. 1893. 11. *Ib.*, 13:581. 1894. 12. Dempsey, *Ont. Fr. Stas. An. Rpt.*, 1:24. 1894. *fig.* 13. Hansen, *S. D. Sta. Bul.*, 76:110. 1902.

SYNONYMS. *Aromatic Spike* No. 354 (13). COMPTE ORLOFF (1). COUNT ORLOFF (8). Count Orloff (13). *De Revel* (2). GRAND SULTAN (3, 7, 9, 10, 12). *Grand Sultan* (2). *Green Transparent* (13). GROSKOE SELENKA GRÜNER (5, 11). ORLOFF (6). Orloff (4). *Orlovskoe* (6). ORLOWSKOE (4). *Red Transparent* (13). *Revelstone Pippin* (2). TRANSPARENTE JAUNE (2). *Transparente Rouge* (1). *Transparente de Sainte-Léger* (2). *Transparente Verte* (1). VINEUSE ROUGE (13). *Vineuse Rouge* (1).

Hansen gives the following description of this variety (13): "Origin, Russia; as fruited in the Iowa Experiment Station orchard, this variety and Red Transparent, Count Orloff, Grand Sultan, Green Transparent and

¹ *An. Hort.*, 1892.

Aromatic Spike No. 354 are identical or very similar. Tree a strong grower, round topped, a heavy annual bearer. Fruit medium to large, round oblate conic, regular; surface greenish-yellow, rarely faintly splashed with red on sunny side, overlaid with white bloom; dots large, white, few; cavity narrow, abrupt, with irregular patch of russet, stem short, stout, often clubbed; basin small, shallow; calyx, small, closed. Core closed or nearly so, clasping; tube long, funnel-shaped; flesh white, firm, juicy, fragrant, subacid, good for table, very good for cooking. Season very early, about one week before Yellow Transparent, but perishable and should be picked early to prevent water-coring and rotting on the tree. Evidently the name is a misnomer as it means Red Wine Colored."

As grown at this Station Count Orloff, Grand Sultan and Groskoe Selenka Grüner are identical or very similar, and none of them is very desirable.

WASHINGTON STRAWBERRY.

REFERENCES. 1. *N. Y. Agr. Soc. Trans.*, 1849:117. 2. *Mag. Hort.*, 23:26. 1857. 3. Hovey, *ib.*, 24:79. 1858. fig. 4. *Am. Pom. Soc. Cat.*, 1869. 5. Downing, 1869:396. fig. 6. Thomas, 1875:205. 7. Downing, 1881:11, 12 index, app. 8. Barry, 1883:341. 9. Wickson, 1889:244. 10. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 11. Bailey, *Am. Hort.*, 1892:252. 12. Burnett, *La. Sta. Bul.*, 27:926. 1894. 13. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:60. 1903. 14. Budd-Hansen, 1903:197. fig. 15. Farrand, *Mich. Sta. Bul.*, 205:47. 1903. 16. Beach and Clark, *N. Y. Sta. Bul.*, 248:150. 1904. SYNONYMS. *Junata* (7). WASHINGTON (2, 3). *Washington of Maine* (7). *Washington County Seedling* (2). WASHINGTON STRAWBERRY (1, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16). WASHINGTON Strawberry (14). *Washington Strawberry* (3).

Fruit smooth, of good size and pretty good color, fairly uniform in shape but somewhat variable in size, desirable for either dessert or culinary uses. It is quite variable in season in different years and in different localities. As fruited at this Station it comes in season in September or October, and some portion of the fruit may be kept in fair condition into the winter or sometimes through the winter. In ordinary storage its commercial limit is October, and in cold storage November (13, 16). The fruit hangs well to the tree. The tree is vigorous, hardy, healthy, moderately long-lived, comes into bearing rather early and is a reliable cropper, yielding good crops biennially or almost annually. It is a good variety for home use, but evidently is not wanted in market. Its season is rather short, and it begins to mature at a time when the markets are filled with other fruits.

Historical. Washington Strawberry was first exhibited at the Fair of the State Agricultural Society in Syracuse in 1849 (1, 3). It originated on the farm of Job Whipple, Union Springs, Washington county, N. Y. (1). It was included in the catalogue of the American Pomological Society in 1869 (4). It is still listed by nurserymen and has been disseminated to some extent in various parts of the continent. Although it has been known for a half century it has failed to establish itself in the commercial orchards of this state and is but little known among New York fruit growers.

TREE.

Tree medium to large, vigorous to moderately vigorous. *Form* rather flat, spreading, open. *Twigs* below medium to short, straight or nearly so, rather slender to stout with large terminal buds; internodes medium or below. *Bark* clear brownish-red or with more or less olive-green, lightly streaked with scarf-skin; pubescent. *Lenticels* scattering, small to medium, round or somewhat elongated, slightly raised. *Buds* medium to large, broad, plump, obtuse, free or nearly so, pubescent.

FRUIT.

Fruit medium to large or very large. *Form* globular, usually inclined to conic, base rounding or sometimes flattened, nearly regular; sides often a little unequal. *Stem* short and rather thick or sometimes long. *Cavity* below medium to medium, acute to somewhat acuminate, rather shallow to deep, narrow to moderately broad, occasionally lipped, often somewhat furrowed, usually thinly russeted. *Calyx* below medium to rather large, usually somewhat open; lobes a little separated and broad at the base, narrow above, long, acute to acuminate. *Basin* small to medium, narrow to moderately wide, medium in depth, abrupt, somewhat furrowed, wrinkled.

Skin rather thin, tough, smooth, somewhat waxy, greenish or yellow, washed and mottled with red, conspicuously splashed and striped with bright carmine and overspread with thin bloom. *Dots* numerous, russet or whitish and rather conspicuous, often submerged. *Prevailing effect* striped red.

Calyx tube rather large, wide, cone-shape with core lines meeting, sometimes becoming funnel-form with clasping core lines. *Stamens* basal or nearly so.

Core below medium to above, axile or sometimes abaxile; cells not uniformly developed, usually symmetrical and more or less open, sometimes closed; core lines meeting if the calyx tube is cone-shape, clasping if it is funnel-form. *Carpels* variable, ovate to broadly obovate, sometimes a little emarginate, often tufted. *Seeds* dark, medium in size, rather long, somewhat acute; often many are abortive.

Flesh whitish tinged with yellow, firm, rather fine to a little coarse, crisp, tender, very juicy, pleasant subacid, sprightly, good to very good.

Season from September or October into early winter.

WATER.

REFERENCES. 1. Downing, *Horticulturist*, 19:172. 1864. figs. 2. Warder, 1867:735. 3. Downing, 1869:397. fig. 4. Thomas, 1875:315. 5. *Am. Pom. Soc. Cat.*, 1877:16. 6. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 7. Bailey, *Am. Hort.*, 1892:252.

SYNONYMS. None.

A mild flavored dessert apple of medium size, pale yellow or greenish with attractive blush of lively red; in season from October to December. The tree is a rather moderate grower, does not come into bearing young and is a biennial bearer yielding from fair to good crops. Not recommended for commercial planting.

Historical. Origin Durham township, Bucks county, Pa. (1, 3). It was entered in the catalogue of the American Pomological Society in 1877 (5) and dropped from that list in 1897. It is but little known in this state.

TREE.

Tree moderately vigorous with short, slender, curved branches. *Form* erect or roundish, rather dense. *Twigs* long, curved, moderately stout; internodes short. *Bark* dark brown, heavily coated with gray scarf skin; pubescent near tips. *Lenticels* scattering, small, round, not raised. *Buds* rather prominent, medium to large, plump, obtuse, free, pubescent.

FRUIT.

Fruit medium size. *Form* broadly ovate to roundish conic varying to oblong conic with flattened ends, nearly regular. *Stem* short to medium, slender. *Cavity* acuminate, rather narrow to moderately broad, moderately shallow to deep, often compressed, sometimes thinly russeted, the russet not extending beyond the cavity. *Calyx* small to medium, closed or open; lobes long, narrow, acute to acuminate. *Basin* varying from shallow to rather deep and abrupt, narrow to medium in width, furrowed and wrinkled.

Skin pale yellow or greenish with very attractive, lively pinkish-red blush, in well colored specimens deepening to dark or purplish-red, not striped, over-spread with thin bloom. *Dots* numerous, medium to small, grayish or whitish, often submerged.

Calyx tube short, wide, broadly conical. *Stamens* basal to median.

Core rather small, somewhat abaxile; cells closed or slightly open; core lines slightly clasping. *Carpels* small, slightly obovate to obcordate, emarginate. *Seeds* medium in size, few, dark brown, varying from blunt and good flat to acute and rather narrow.

Flesh nearly white, fine, crisp, tender, juicy, pleasant mild subacid, good.

Season October to December or later.

WEALTHY.

REFERENCES. 1. Downing, 1869:398. 2. Foster, *Horticulturist*, 25:362. 1870. 3. *Am. Pom. Soc. Cat.*, 1871:10. 4. Thomas, 1875:515. 5. *Montreal Hort. Soc. Rpt.*, 1876:19. 6. *Ib.*, 1879:33. fig. 7. *1a. Hort. Soc. Rpt.*, 1879:



WEALTHY

453. 8. Sheppard, *Montreal Hort. Soc. Rpt.*, 8:140. 1881-82. 9. Foster, *Id. Hort. Soc. Rpt.*, 1882:100. 10. Barry, 1883:356. 11. Gideon, *Am. Pom. Soc. Rpt.*, 1885:114. 12. *Montreal Hort. Soc. Rpt.*, 1886-87:17. 27. 13. *Rural N. Y.*, 46:494, 751. 1887. 14. *Am. Pom. Soc. Rpt.*, 1887:92. 15. *Can. Hort.*, 11:31. 1888. 16. *Rural N. Y.*, 48:177. 1889. 17. Gibb, *Montreal Hort. Soc. Rpt.*, 15:19. 1890. 18. Hamilton, *Id.*, 15:27. 1890. 19. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:298. 20. *Can. Hort.*, 14:119, 131, 331. 1891. 21. Taylor, *Me. Pom. Soc. Rpt.*, 1892:57. 22. *Rural N. Y.*, 51:705. 1892. 23. *Ont. Fr. Gr. Assn. An. Rpt.*, 26:65. 1894. 24. *Rural N. Y.*, 54:682. 1895. 25. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 3:17. 1896. figs. 26. Waugh, *Vt. Sta. Bul.*, 61:32. 1897. 27. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:357. 28. *Am. Gard.*, 20:404. 1899. 29. Waugh, *Vt. Sta. An. Rpt.*, 14:312. 1901. 30. Hays, *U. S. Dict. of Vegetable Physiology and Pathology Bul.*, 29:13. 1901. 31. *Can. Hort.*, 25:121. 1902. 32. Van Deman, *Rural N. Y.*, 61:249. 1902. 33. Munson, *Me. Sta. An. Rpt.*, 18:85, 90. 1902. 34. Hansen, *S. D. Sta. Bul.*, 76:112. 1902. fig. 35. Dickens and Greene, *Kan. Sta. Bul.*, 106:56. 1902. 36. *Can. Hort.*, 26:12. 1903. 37. Budd-Hansen, 1903:198. fig. 38. Farrand, *Mich. Sta. Bul.*, 205:47. 1903. 39. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:60. 1903. 40. Macoun, *Can. Hort.*, 27:71. 1904. fig. of tree. 41. Beach and Clark, *N. Y. Sta. Bul.*, 248:150. 1904. 42. Ragan, *U. S. B. P. I. Bul.*, 56:326. 1905.

SYNONYMS. None.

This variety is particularly valuable for cold climates because the tree is very hardy and the fruit sells well, being bright red and good in quality for either dessert or culinary uses. It is in season from October to early winter or mid-winter. In ordinary storage its commercial limit is October, but in cold storage it may be kept till January or later. It does not stand heat very well before going into storage, and goes down rather quickly (41). Young trees or trees which are making a thrifty growth produce fruit of good size, but mature, slow-growing trees are apt to yield a considerable percentage of undersized fruit, especially when they are overloaded, as is often the case. The crop ripens unevenly, and more than one picking should be made in order to secure the fruit in prime condition. If it is left upon the tree till fully colored there is apt to be considerable loss by dropping. The tree is a good thrifty grower when young, but with maturity it becomes a moderate or rather slow grower, forming a medium-sized or rather dwarfish tree. Wealthy is being planted for commercial purposes in many parts of the state, but in most localities the trees are as yet comparatively young; in some sections of the state it is being planted

more than any other apple of its season. Trees that have become old enough to develop the tendency to produce rather small fruit are in some places being grafted over to other sorts. In other places fruit growers, by adopting such treatment as thinning the fruit and keeping the soil fertile, continue to produce apples of good marketable size when the trees are mature.

Historical. Originated by Peter M. Gideon, Excelsior, Minn., from seed of the Cherry Crab, which he obtained about 1860 from Albert Emerson, Bangor, Me. (11, 28). Ragan (42) states that the fruit was first described in the Western Farmer in 1869. It has been extensively disseminated, particularly in those apple-growing districts where a tree of superior hardiness is especially desired. It is generally listed by nurserymen and its cultivation is gradually increasing.

TREE.

Tree somewhat dwarfish to medium size, moderately vigorous with short, moderately stout, curved branches. *Form* upright spreading or roundish, open and somewhat drooping. *Twigs* long, curved, slender; internodes long. *Bark* dark brown, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, medium to small, oblong, not raised, rather conspicuous. *Buds* medium size, broad, plump, obtuse, free, pubescent.

FRUIT.

Fruit above medium to large when well grown but often small on old trees; pretty uniform in shape and quality but more or less uneven in size. *Form* roundish conic, slightly flattened at base, regular, symmetrical. *Stem* usually short to medium, but rather long on small fruit and rather slender. *Cavity* decidedly acuminate, rather deep, moderately narrow to rather broad, russeted. *Calyx* medium size, closed or partly open; lobes broad, obtuse to acute. *Basin* medium in depth to rather shallow, rather narrow, abrupt, smooth, symmetrical.

Skin thin, tough, pale yellow or greenish, blushed and marked with narrow stripes and splashes of red, deepening in highly colored specimens to brilliant red, very attractive. *Dots* numerous, small, inconspicuous, pale or russet. *Prevailing effect* bright red.

Calyx tube conical approaching funnel-form. *Stamens* median.

Core medium to very small, axile or sometimes slightly abaxile; cells symmetrical, slightly open; core lines clasping. *Carpels* medium to rather small, roundish, narrowing toward base and apex, smooth, flat. *Seeds* moderately dark brown, above medium, rather acute.

Flesh whitish sometimes stained with red, moderately fine, crisp, tender, very juicy, agreeable subacid, sprightly, somewhat aromatic, good to very good.

Season October to January.

WESTERN BEAUTY.

REFERENCES. 1. *Gardeners' Monthly*, 1861:124 (cited by 2). 2. Kenrick, *Mag. Hort.*, 29:73. 1863. fig. 3. Warder, 1857:464. fig. 4. Downing, 1869:89. 5. *Ib.*, 1872:37 app. 6. *Am. Pom. Soc. Cat.*, 1877. 7. Barry, 1883:356. 8. Thomas, 1885:528. 9. Bailey, *Am. Hort.*, 1892:252. 10. Burrill and McCluer, *Ill. Sta. Bul.*, 45:345. 1896.

SYNONYMS. BEAUTY OF THE WEST (4). *Beauty of the West* (10). *Big Rambo* (3, 5). *Musgrove's Cooper* (3, 4, 5). OHIO BEAUTY (1). *Ohio Beauty* (3, 4, 5). WESTERN BEAUTY (2, 3, 5, 6, 7, 8, 9, 10).

The three varieties, Western Beauty, Grosh and Summer Rambo, resemble each other so closely in fruit that it is practically impossible to distinguish the one from the other from the examination of the fruit alone. The Summer Rambo, however, ripens about a month earlier than the other two varieties and it can consequently be readily distinguished in the orchard. Pomologists are in doubt as to whether the Grosh and Western Beauty are two distinct varieties or the same variety under two names. We have been unable to obtain sufficient evidence to determine this point.

For a technical description of the fruit, see Grosh, page 89.

Hyde King was received here for testing under the name Western Beauty and consequently is referred to under that name erroneously in some published accounts of its record at this Station. See Volume I, page 166.

Historical. Origin unknown. First introduced to notice by William F. English of Rhinehart, Auglaize county, Ohio (1, 2).

WHITE ASTRACHAN.

REFERENCES. 1. *Pom. Mag.*, 2:No. 96. 1829. col. pl. 2. *London Hort. Soc. Cat.*, 1831:No. 32. 3. Floy-Lindley, 1833:6. 4. *Mag. Hort.*, 1:391, 392. 1835. 5. Manning, *Ib.*, 7:52. 1841. 6. Downing, 1845:78. 7. Emmons, *Nat. Hist. N. Y.*, 3:17. 1851. 8. Elliott, 1854:179. 9. Warder, 1857:735. 10. Leroy, 1873:79. fig. 11. Thomas, 1875:516. 12. *Montreal Hort. Soc. Rpt.*, 1:9. 1876. 13. *Ib.*, 5:23. 1879. 14. Hogg, 1884:238. 15. Wickson, 1889:243. 16. *Am. Pom. Soc. Cat.*, 1889:14. 17. Hansen, *S. D. Sta. Bul.*, 76:113. 1903.

SYNONYMS. ASTRACHAN BLANCHE (10). *Astracan d'Ete* (10). *Astracanischer Sommer* (10). ASTRACHAN WHITE (2). *Blanche Glacee d'Ete* (10). *De Glace d'Ete* (10). *De Glace Hatice* (10). *De Moscovie d'Ete* (10). *Gelee d'Ete* (10). *Glace de Zelande* (1, 2, 3, 6, 7, 8, 10). *Glacee d'Ete* (10). *Naliki Jabloky* (10). *Pomme Astrachan* (3). *Pomme d'Astrachan* (2). *Pyrus Astracanica* (1, 2, 6, 7). *Taffitai* (10). *Transparent Apple* (14). *Transparente de Astracan* (10). *Transparente d'Ete* (10). *Transparente de Zurich* (10). *Transparente de Moscovie d'Ete* (10). *Transparent Moscovie* (7). *Transparent de Moscovie* (1, 2, 3, 6, 8). WHITE ASTRACHAN (1, 3, 4, 5, 7, 15). *White Astracan* (10). WHITE ASTRACHAN (6, 8, 9, 11, 12, 13, 14, 16, 17).

A Russian apple of little or no value for this region. Fruit medium size, roundish to roundish oblate, waxen yellow or whitish with faint streaks of red; flesh white, acid, good for culinary use; season August and September (6, 17).

Historical. It has been known in this country for many years (4, 5, 6, 7, 8, 9). It was not entered on the catalogue of the American Pomological Society till 1889 (16) and was dropped from that list in 1891. It is practically unknown in New York.

WHITE JUNEATING.

REFERENCES. 1. *Am. Gard. Cal.*, 1806:584. 2. Coxe, 1817:100. fig. 3. *London Hort. Soc. Cat.*, 1831:No. 612. 4. Floy-Lindley, 1833:3. 5. *Mag. Hort.*, 1:326. 1835. 6. Downing, 1845:78. fig. 7. Thomas, 1849:144. fig. 8. Cole, 1849:97. 9. Emmons, *Nat. Hist. N. Y.*, 3:17. 1851. 10. Elliott, 1854:162. 11. Hooper, 1857:20. 12. *Am. Pom. Soc. Cat.*, 1862. 13. *Horticulturist*, 21:292. 1866. fig. 14. Warder, 1867:417. 15. Fitz, 1872:143, 151, 161. 16. Hogg, 1884:117. 17. Wickson, 1889:242. 18. Bruner, *N. C. Sta. Bul.*, 182:23. 1903. 19. Budd-Hansen, 1903:200.

SYNONYMS. *Bracken* (8, 11, of Ohio 7). *Carolina* (10). *Caroline* (10). *Early Jenetting* (16). *Early May* (10, ?14). *Gennetting* (17). GINETTING (1). Ginetting (16). Jenetting (2). Jenetting (17). JOANETING (16). *Juneating* (1). JUNEATING (4). *Juneating* (6, 8, 10, 11, 14, 16). JUNETING (2). Juneting (16). *Owen's Golden Beauty* (3, 6, 10, 16). *Primiting* (16). WHITE JUNEATING (3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 17, 18, 19). *White Juneating* (16). *Yellow May* (18, 19). *Yellow June* (14).

This apple has little to recommend it except that it ripens very early in the season. It is small, roundish oblate, pale yellow, sometimes faintly blushed and has white, crisp, pleasant subacid flesh which becomes mealy if kept only a few days after it ripens. The tree is not large, only a moderate grower and not a great bearer (6, 8, 16).

Historical. Hogg (16) gives the following interesting account of the history of this apple and the probable derivation of the name Juneating:

"One of our oldest apples, and although generally known and popular, seems to have escaped the notice of Miller, who does not even mention it in any of the editions of his Dictionary. As I have doubts of this being the Geneting of Parkinson — his figure being evidently intended for the Margaret, which in some districts is called Joaneting — the first mention we have of this variety is by Rea, in 1665, who describes it as 'a small, yellow, red-sided apple, upon a wall, ripe in the end of June.'

"'Juneating,' as applied to this apple, is quite a misnomer. Abercrombie was the first who wrote it June-eating, as if in allusion to the period of its maturity, which is, however, not till the end of July. Dr. Johnson, in his Dictionary, writes it Ginetting, and says it is a corruption of Janetou (Fr.), signifying Jane or Janet, having been so called from a person of that name. Ray¹ says, 'Pomum Ginetinum, quod unde dictum sit met latet.' Indeed, there does not seem ever to have been a correct definition given of it.

"My definition of the name is this. In the Middle Ages, it was customary to make the festivals of the Church periods on which occurrences were to take place or from which events were to be named. Even in the present day we hear the country people talking of some crop to be sown, or some other

¹ *Hist. Plant.*, II, 1447.

to be planted, at Michaelmas, St. Martin's, or St. Andrew's tide. It was also the practice for parents to dedicate their children to some particular saint, as Jean Baptiste, on the recurrence of whose festival all who are so named keep it as a holiday. So it was also in regard to fruits, which were named after the day about which they came to maturity. Thus, we have the Margaret Apple, so called from being ripe about St. Margaret's Day, the 20th of July; the Magdalene, or Maudlin, from St. Magdalene's Day, the 22d of July. And in Curtius¹ we find the *Joannina*, so called, 'Quod circa divi Joannis Baptistæ nativitatem esui sint.' These are also noticed by J. Baptista Porta; he says, 'Est genus alterum quod quia circa festum Divi Joannis maturiscit, vulgus *Melo de San Giovanni* dicitur.' And according to Tragus,² 'Quæ apud nos prima maturantur, Sancti Johans Opfell, *Latine*, *Præcocia mala* dicuntur.'

"We see, therefore, that apples were called Joannina because they ripened about St. John's Day, and we have among the old French pears *Amiré Joannet*—the 'Wonderful Little John,' which Merlet informs us was so called because it ripened about St. John's Day. If, then, we add to Joannet the termination *ing*, so general among our names of apples, we have *Joanneting*. There can be no doubt that this is the correct derivation of the name of this apple."

WHITE SPANISH REINETTE.

REFERENCES. 1. *Pom. Mag.*, 3:No. 110. 1830. col. pl. 2. Floy-Lindley, 1833:61. 3. Downing, 1845:130. 4. Emmons, *Nat. Hist. N. Y.*, 3:88. 1851. 5. Elliott, 1854:162. 6. Downing, 1859:404. 7. Leroy, 1873:669. fig. 8. Thomas, 1875:231. 9. Hogg, 1884:190. 10. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:300.

SYNONYMS. *American Fall Pippin* (9). *Belle Joséphine* (9). *Blanche* (7). *Blanche d'Espagne* (7). *Canuesar* (1, 6, 9). *Camoisais du roi d'Espagne* (7). *Camoise Blanche* (7). *Camoisée Blanche* (7). *Canuezas* (7). *Camuzar* (7). *Cobbett's Fall* (1) *err.* *Cobbett's Fall Pippin* (2, 3, 6, 7) *err.* *Concombre Ancien* (1, 2, 5, 7, 9). *De Ratteau* (1, 2, 5, 7, 9). *D'Espagne* (1, 2, 3, 5, 6, 7). *Elgin Pippin?* (6). *Episcopale* (7). *Fall Pippin* (1, 2, 3, 7) *err.* (6) *err.* *Joséphine* (9). *Large Fall* (1). *Large Fall Pippin* (2, 3, 6, 7). *Philadelphia Pippin* (7). *Reinette A Gobelet* (7). *Reinette Blanche* (7). *REINETTE BLANCHE D'ESPAGNE* (3, 9). *Reinette Blanche d'Espagne* (1, 2, 5, 6, 7, 8). *REINETTE D'ESPAGNE* (7). *Reinette d'Espagne* (1, 6, 9). *Reinette Tendre* (7). *Saint-Germain* (7). *WHITE SPANISH REINETTE* (1, 2, 4, 5, 6, 8, 10). *White Spanish Reinette* (3, 7, 9). *York Pippin* (7).

This variety belongs in the group with Fall Pippin and Holland Pippin. It resembles Fall Pippin in the growth of the tree as well as in the color and character of the fruit, but is less regular in shape and keeps later (3, 6). Season here October to January or February; Hogg gives its season in

¹ *Hortorum*, p. 522.

² *Hist.*, p. 1043.

England as December to April (9). Lyon gives its season in Michigan as October to January (10).

Historical. This is an old European variety which has long been cultivated in Spain, France and England (1, 2, 3, 6, 7, 9). It was early imported into this country and is perhaps the parent of our Fall Pippin and Holland Pippin (6). According to Lyon (10) it is seldom seen under its own name. It appears to be but little cultivated in New York having been superseded by other varieties.

FRUIT.

Fruit very large. *Form* roundish oblate or inclined to oblong, angular, uneven at the crown where it is nearly as broad as at the base (3, 6, 9). *Stem* short. *Cavity* narrow, rather small, regular. *Calyx* large, open. *Basin* deep, broadly angular, irregular, oblique.

Skin smooth, waxy, yellowish-green, with orange tinge and brownish-red blush on the exposed cheek.

Calyx tube conical. *Stamens* marginal. *Cells* open, obovate.

Flesh yellowish-white, crisp, tender, juicy, subacid, very good for either dessert or culinary purposes.


Season October to January or February.

WILLIAMS.


REFERENCES. 1. Kenrick, 1832:29. 2. Manning, 1838:49. 3. *Ib.*, *Mag-Hort.*, 7:51. 1841. 4. Downing, 1845:79. 5. Hovey, *Mag. Hort.*, 14:118. 1848. *fig.* 6. Goodrich, 1849:51. 7. Cole, 1849:100. *fig.* 8. Thomas, 1849:142. *fig.* 9. Waring, 1851:32. *fig.* 10. Barry, 1851:282. 11. Emmons, *Nat. Hist. N. Y.*, 3:12. 1851. 12. Elliott, 1854:163. 13. *Am. Pom. Soc. Cat.*, 1854. 14. Hooper, 1857:100. 15. Warder, 1867:618, 736. 16. Fittz, 1872:161. 17. *Montreal Hort. Soc. Rpt.*, 1879:26. 18. *Id.* *Hort. Soc. Rpt.*, 1883:507. 19. Hogg, 1884:243. 20. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:300. 21. Bailey, *Am. Hort.*, 1892:252. 22. Hoskins, *Rural N. Y.*, 53:248. 1894. 23. Beach, *N. Y. Sta. An. Rpt.*, 14:255. 1895. 24. *Gard. and For.*, 8:248. 1895. 25. *Can. Hort.*, 20:183. 1897. 26. Bunyard, *Jour. Roy. Hort. Soc.*, 1898:357. 27. Waugh, *Vt. Sta. An. Rpt.*, 14:312. 1901. 28. Budd-Hansen, 1903:203. *fig.*

SYNONYMS. *Lady's Apple* (5). *Queen* (5). WILLIAMS (1, 5, 6, 7, 12, 20, 26, 27). *Williams* (7, 8). *Williams Early* (5, 12, 15, 19). *Williams Early Red* (7, 12). WILLIAMS FAVORITE (3, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 21, 22, 23, 24, 25). *Williams Favorite* (5, 7, 12, 27). *Williams Favorite Red* (5, 8, 12). WILLIAMS FAVORITE (28). WILLIAMS FAVORITE RED (2). WILLIAMS FAVOURITE (4, 19). *Williams Red* (8, 12, 19).

Williams is a very beautiful, bright red apple of mild agreeable flavor, good for dessert but not suitable for culinary uses. It is a favorite in Boston and other eastern markets, and is grown to a limited extent for commercial purposes in some portions of Eastern New York. It does not stand shipping very well, the skin being



WILLIAMS.



thin, tender and easily bruised, therefore best handled in small packages. It is in season during late August and early September. Under favorable conditions the fruit becomes rather large, but with very heavy crops it is apt to be rather small unless properly thinned. The crop ripens unevenly, and more than one picking is required to secure the fruit in prime condition. The tree being only moderately vigorous, it is an advantage to topwork it upon some thrifty hardy stock, such as Northern Spy, Rhode Island Greening, or Tolman Sweet. When topworked in this way the Williams becomes a rather vigorous grower, makes a tree of pretty good size, comes into bearing early and in favorable locations, under good treatment, is a reliable cropper, yielding good crops annually or nearly annually. It can be recommended for commercial planting where fruit of this type and season is desired.

Historical. Williams originated in Roxbury, Mass., more than 150 years ago. It was brought to the notice of the Massachusetts Horticultural Society in 1830 and then named Williams. It had previously been known in market under the name Queen and Lady's Apple (5). It was entered in the catalogue of the American Pomological Society in 1854 and is still retained on that list (13). It has become widely disseminated and is still often listed by nurserymen (21) but is nowhere being planted to any considerable extent.

TREE.

Tree rather small and a slow grower but when topworked on vigorous stock and properly tilled and fertilized it becomes rather large and vigorous. *Form* upright spreading or roundish, somewhat dense. *Twigs* short, curved, moderately stout, with large terminal buds; internodes short. *Bark* dark brown tinged with green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* quite numerous, small to medium size, oblong, raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium or under favorable circumstances rather large, pretty firm in size and shape. *Form* oblong conic to roundish conic, broadly ribbed; sides often unequal. *Stem* medium to long, moderately thick. *Cavity* obtuse, shallow, rather broad, furrowed, sometimes russeted. *Calyx* above medium size, usually closed; lobes long. *Basin* medium to rather shallow, rather narrow to moderately broad, a little abrupt, somewhat furrowed.

Skin moderately thick, rather tender, nearly smooth, pale yellow overlaid with bright deep red, indistinctly striped with dark red or crimson. *Dots* numerous, inconspicuous, small, grayish or russet.

Calyx tube long, narrow, funnel-shape or approaching cylindrical, sometimes extending to the core. *Stamens* marginal.

Core medium to rather large, axile; cells closed or slightly open; core lines clasping. *Carpels* ovate to roundish. *Seeds* above medium, rather narrow, long, moderately plump, acute or nearly acuminate, dark brown.

Flesh sometimes tinged with red, firm, a little coarse, moderately crisp, tender, rather juicy, becoming dry when overripe, pleasant mild subacid, aromatic, good.

Season late August and September.

WILLIS SWEET.

REFERENCES. 1. *Mag. Hort.*, 18:491. 1852. 2. ? Elliott, 1854:179. 3. Hicks, *Horticulturist*, 21:361. 1866. fig. 4. Warder, 1867:635. 5. Downing, 1869:408. fig. 6. *Am. Pom. Soc. Cat.*, 1869. 7. Thomas, 1875:516. 8. Bailey, *Am. Hort.*, 1892:252.

SYNONYMS. *Pear Lot* (5). *Pear-Tree Lot* (5). WILLIS SWEET (2, 4, 5, 6, 7, 8). WILLIS SWEETING (1, 3).

A rather large sweet apple in season in late summer and early autumn. According to Downing the tree is a good grower and productive; the fruit whitish with shade of light red washed with crimson; flesh crisp, juicy, tender, rich, sweet, very good; valuable for dessert, for culinary purposes and for market (5).

Historical. A chance seedling that originated at Oyster Bay, Long Island, about 1800, on the farm of Edmond Willis. It first had the local name of Pear-tree Lot or Pear Lot. Later it was named Willis Sweeting by Parsons & Co., of Flushing, N. Y. (3). In 1869 it was entered in the catalogue of the American Pomological Society (6), but was dropped from that list in 1899. It is still occasionally listed by nurserymen (8) but is now seldom planted. It is not generally known in New York.

WINE RUBETS.

REFERENCES. 1. Budd, *Ia. Agr. Coll. Bul.*, 1885:7. 2. Beach, *N. Y. Sta. An. Rpt.*, 12:600. 1893. 3. *Ib.*, 12:603. 1893. 4. Ragan, *U. S. B. P. I. Bul.*, 56:337. 1905.


SYNONYMS. CUT WINE (2). *Cut Wine* (3, 4). *No. 210* (1, 2, 3, 4). *Rubets Vinogradnui* (4). *Rubets vinogradnui* (1). *Rubetz vinogradnui* (4). *VINOGRAD* (1). *Vinograd* (3). WINE RUBETS (3, 4).

Fruit below medium size, nearly symmetrical, covered with delicate bloom. Skin green, lightly shaded with red and with a crimson cheek. Basin shallow, wrinkled. Stem medium length, slender, set in a deep cavity. Flesh mild subacid, fair to good in quality. Begins to ripen here about the 1st of August. Not recommended for planting in New York.

Historical. A Russian apple imported by the United States Department of Agriculture. It was received here in 1888 from Dr. T. H. Hoskins, Newport, Vt., under the name Cut Wine.



WOLF RIVER



WINTHROP GREENING.

REFERENCES. 1. Hovey, *Mag. Hort.*, 12:475. 1846. 2. Cole, 1849:104. 3. *Am. Pom. Soc. Cat.*, 1854. 4. *Mag. Hort.*, 21:63. 1855. 5. Downing, 1857:204. 6. Warder, 1867:737. 7. Thomas, 1875:209. 8. Downing, 1881:11 index, app. 9. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:300. 10. Burrill and McCluer, *Ill. Sta. Bul.*, 45:347. 1896.

SYNONYMS. *Howe Apple* (2, 5). *Kennebec Seedling* (8). LINCOLN (1). LINCOLN PIPPIN (4). *Lincoln Pippin* (2, 5). WINTHROP GREENING (2, 3, 5, 6, 7, 8, 9, 10).

Fruit yellow, tinged with red, of good size and good quality; season September to early winter. The flesh is tender, crisp, very juicy, sprightly, mild subacid (2). The tree is a shy bearer.

Historical. Originated in Winthrop, Me., about the year 1800 (1, 2). It was entered in the catalogue of the American Pomological Society in 1854 (3) and dropped from that list in 1897. It is but little known in New York.

WOLF RIVER.

REFERENCES. 1. *Wis. Hort. Soc. Rpt.*, 1875:107. 2. *Ia. Hort. Soc. Rpt.*, 1879:452. 3. Budd, *Ib.*, 1880:524. 4. Downing, 1881:113 app. 5. *Am. Pom. Soc. Cat.*, 1881:16. 6. Barry, 1883:357. 7. *Wis. Hort. Soc. Rpt.*, 1886:148. 8. Van Deman, *U. S. Pom. Rpt.*, 1886:271. fig. 9. *Am. Pom. Soc. Rpt.*, 1887:93. 10. *Can. Hort.*, 11:9. 1888. 11. Bailey, *Am. Hort.*, 1892:253. 12. *Can. Hort.*, 16:34. 1893. 13. *Ib.*, 20:34. 1897. 14. Thomas, 1897:286. fig. 15. Waugh, *Vt. Sta. Bul.*, 61:32. 1897. 16. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 6:32. 1899. 17. *Can. Hort.*, 22:121, 437. 1899. figs. 18. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 7:10. 1900. figs. 19. Waugh, *Vt. Sta. An. Rpt.*, 14:313. 1901. 20. Hansen, *S. D. Sta. Bul.*, 76:116. 1902. fig. 21. Dickens and Greene, *Kan. Sta. Bul.*, 106:56. 1902. 22. Budd-Hansen, 1903:207. fig. 23. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:61. 1903. 24. Beach and Clark, *N. Y. Sta. Bul.*, 248:151. 1904.

SYNONYMS. None.

This is a variety of the Aport group. It resembles Alexander very closely in size, form, and color. Hansen states (20) that it is "supposed to be a seedling of the Alexander, which it sometimes resembles, but is more round and less conical, and averages larger, as grown in the West. The Wolf River has largely superseded Alexander in the western states. Tree a strong spreading grower, not an early bearer, but productive in alternate years." As fruited at this Station it is in season from September to December, with October as the commercial limit in ordinary storage. In cold storage it may be held till January. It does not stand heat well,

and goes down quickly (24). The tree is very hardy and a good grower, and is a biennial or sometimes annual cropper, yielding moderate to good crops. The fruit, being large, shapely and highly colored, often sells well because of its attractive appearance; some fruit growers are finding it a profitable variety.

Historical. Originated by W. A. Springer, near Wolf River, Fremont county, Wis., hence its name. It was entered in the catalogue of the American Pomological Society in 1881 (5). It is frequently listed by nurserymen (11). Within recent years it has been planted to a limited extent in New York state and at the present time its cultivation is probably increasing somewhat.

TREE.

Tree large, moderately vigorous. *Form* much spreading, open and inclined to droop. *Twigs* short, straight, slender; internodes short. *Bark* brown, tinged with green, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, small, round, not raised. *Buds* small, plump, obtuse to acute, free, slightly pubescent.

FRUIT.

Fruit large, uniform in size and fairly uniform in shape. *Form* broad and flat at the base and somewhat inclined to conic or roundish, often somewhat irregular. *Stem* short to medium, rather thick, not exerted. *Cavity* acuminate, usually deep, rather wide and very heavily russeted. *Calyx* medium to large, open or closed. *Basin* medium to deep, moderately narrow, abrupt, usually smooth, somewhat broadly furrowed.

Skin rather thick, pale bright yellow or greenish, mottled and blushed with bright deep red and marked with conspicuous splashes and broad stripes of bright carmine. *Dots* numerous, medium to rather large, areolar, depressed, pale or russet.

Calyx tube conical. *Stamens* median to basal.

Core below medium to rather large, somewhat abaxile; cells closed or partly open; core lines clasping. *Carpels* broadly cordate, approaching elliptical, slightly emarginate, somewhat tufted. *Seeds* dark brown, of medium size, rather wide, short, moderately plump, obtuse.

Flesh slightly tinged with yellow, firm, moderately coarse, tender, juicy, subacid, a little aromatic, fair to good.

Season September to December.

WORKAROE.

REFERENCES. 1. N. Y. Sta. An. Rpt., 8:349. 1889. 2. Beach, *Ib.*, 11:588. 1892.

SYNONYMS. None.

A Russian apple of good size, pale yellow, blushed and striped with red and overspread with pinkish bloom. Flesh firm, crisp, tender, juicy, rather



WORKAROE (Reduced Size)

mild subacid with an agreeable but not high flavor and good quality. It is a good apple but hardly equal to other varieties of its season. The tree does not come into bearing very young but is a pretty good grower and eventually a good cropper yielding full crops biennially.

Received in 1884 from Ellwanger and Barry, Rochester, N. Y., for testing at this Station.

YELLOW CALVILLE.

REFERENCES. 1. Budd, *Ia. Agr. Coll. Bul.*, 1885:17. 2. Gibb, *Am. Pom. Soc. Rpt.*, 1887:48. 3. Beach and Paddock, *N. Y. Sta. An. Rpt.*, 13:584. 1894. 4. Beach, *W. N. Y. Hort. Soc. Rpt.*, 41:50. 1896. 5. Ragan, *U. S. B. P. I. Bul.*, 56:345, 353. 1905.

SYNONYMS. *Kakvil jeltui* (2, 5). *Kakville scholti* (1, 2, 5). *No. 442* (1, 2, 3, 5). *Voronesh No. 21* (3). YELLOW CALVILLE (1, 2, 3, 4, 5).

An August apple, medium to rather small, smooth, pale yellow, sometimes with faint blush, oblate to oblate conic. Cavity acute, wide, rather shallow; calyx closed; basin shallow, slightly wrinkled; flesh white, fine-grained, tender, moderately juicy, subacid, fair or sometimes good. The tree comes into bearing moderately young and is nearly an annual cropper. Not recommended for cultivation in this state being much inferior to standard sorts of its season.

Historical. This is a Russian apple, being No. 442 of the importation of the United States Department of Agriculture of 1870 (3, 5). Later it was imported by the Iowa Agricultural College under the designation Voronesh No. 21 (1). In 1888 it was received for testing at this Station from Dr. T. H. Hoskins, Newport, Vt. It is practically unknown in New York.

YELLOW TRANSPARENT.

REFERENCES. 1. Leroy, 1873:846. fig. 2. Budd, *Ia. Hort. Soc. Rpt.*, 1880:523. 3. Downing, 1881:114 app. fig. 4. *Am. Pom. Soc. Cat.*, 1881:16. 5. Webster, *Montreal Hort. Soc. Rpt.*, 7:54. 1881. 6. Gibb, *ib.*, 7:154. 1881. 7. *ib.*, *Ia. Hort. Soc. Rpt.*, 1883:437. 8. Barry, 1883:336. 9. Hoskins, *Rural N. Y.*, 43:651. 1884. fig. 10. Penhollow, *Montreal Hort. Soc. Rpt.*, 10:65. 1884. 11. Thomas, 1885:530. 12. *Rural N. Y.*, 44:185, 200. 1885. 13. Gibb, *Montreal Hort. Soc. Rpt.*, 1886-87:14. 14. *Am. Pom. Soc. Rpt.*, 1887:92. 15. *Rural N. Y.*, 46:107, 201, 382. 1887. fig. 16. *Can. Hort.*, 11:209, 223. 1888. 17. Fisk, *Montreal Hort. Soc. Rpt.*, 14:33. 1889. 18. *Can. Hort.*, 13:121, 132, 216, 272, 301. 1890. 19. Hamilton, *Montreal Hort. Soc. Rpt.*, 15:27. 1890. 20. *Can. Hort.*, 15:281. 1892. 21. Bailey, *Am. Hort.*, 1892:253. 22. Budd, *Ia. Sta. Bul.*, 19:535. 1892. 23. *Pa. Sta. Rpt.*, 1892:110. fig. 24. Hoskins, *Gard. and For.*, 7:138. 1894. 25. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 2:11. 1895. figs. 26. Van Deman, *Rural N. Y.*, 55:613. 1896. 27. Hoskins, *ib.*, 56:156. 1897. 28. Woolverton, *Ont. Fr. Stas. An. Rpt.*, 4:5. 1897. figs. 29. Waugh, *Vt. Sta. Bul.*, 61:32. 1897. 30. Alwood, *Va. Sta. Bul.*, 130:122. 1901. 31. Waugh, *Vt. Sta. An. Rpt.*, 14:313. 1901. 32. Hansen, *S. D. Sta. Bul.*, 76:119. 1902. fig. 33. Dickens and Greene, *Kan. Sta. Bul.*, 106:56. 1902. 34. Munson, *Me. Sta. An. Rpt.*, 18:85, 88. 1902. 35. *Rural N. Y.*, 61:626. 1902. fig. 36. Budd-Hansen,

1903:213. fig. 37. Farrand, *Mich. Sta. Bul.*, 205:47. 1903. 38. Bruner, *N. C. Sta. Bul.*, 182:24. 1903.

SYNONYMS. *De Revel* (1). *Grand-Sultan* (1). *Revelstone* (1). *SKWOSNOI SCHOTOI* (7). *Skwosnoi Schotoi* (5). *Transparente de Saint-Leger* (1). *TRANSPARENTE JAUNE* (1). *YELLOW TRANSPARENT* (2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38). *Yellow Transparent* (7).

This is one of the best of the extra early apples, being excellent for culinary use and acceptable for dessert. It is not equal in quality to Early Harvest, but it begins to ripen somewhat earlier and is a more reliable cropper, yielding good crops annually or nearly so. Generally speaking, it is grown in New York state for home use only, but in some places it is cultivated to a limited extent for market, particularly for local market. It is desirable for this purpose because it takes on a good clear yellow color before becoming overripe. On account of its delicate color and tender skin it shows bruises readily and must be handled with extra care. The crop ripens continuously through a period of three or four weeks, and two or more pickings are required in order to secure the fruit in prime condition. It begins to ripen in July, and continues in season in some cases till early September. On young or vigorous-growing trees the fruit may grow rather large, but on mature slow-growing trees, especially when they are overloaded, the fruit is apt to be below medium size unless thinned. The tree is a moderately vigorous grower, hardy, healthy and comes into bearing very young. In some portions of the West it suffers from twig blight (fire blight) but it appears to be quite free from this disease in New York.

Historical. Imported from Russia by the United States Department of Agriculture in 1870. Its merits were first brought to notice in this country by Dr. T. H. Hoskins, of Newport, Vt. (9). It has been disseminated throughout the apple-growing regions of the country from the Atlantic to the Pacific and is now commonly listed by nurserymen (21). In New York its cultivation for home use is gradually increasing, and occasionally it is grown to a limited extent for market.

TREE.

Tree of medium size, moderately vigorous, with short, stout, crooked branches filled with short spurs. *Form* upright at first but becoming spreading or roundish and rather dense. *Twigs* short, curved, stout with large



YELLOW TRANSPARENT

terminal buds; internodes medium. *Bark* conspicuously yellow or tawny, lightly streaked with scarf-skin; pubescent. *Lenticels* quite numerous, medium to small, oval or elongated, not raised. *Buds* medium size, broad, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit medium or above medium, sometimes large, pretty uniform in shape and size. *Form* roundish ovate to roundish conic or oblate conic, slightly ribbed; sides unequal. *Stem* medium to long, rather thick. *Cavity* acute or approaching obtuse, medium to deep, rather narrow, sometimes slightly lipped, sometimes russeted. *Calyx* medium size, closed; lobes medium in length, broad. *Basin* moderately shallow, rather narrow, somewhat abrupt, furrowed and slightly wrinkled.

Skin thin, tender, smooth, waxy, pale greenish-yellow changing to an attractive yellowish-white. *Dots* moderately numerous, greenish and light colored, often submerged.

Calyx tube conical. *Stamens* marginal.

Core medium to small; cells partly open to wide open; core lines clasping. *Carpels* broadly ovate. *Seeds* medium size, rather wide, rather flat, obtuse to slightly acute.

Flesh white, moderately firm, fine-grained, crisp, tender, juicy, sprightly subacid with a pleasant but not high flavor, good or sometimes very good.

Season late July and August.

YOPP.

REFERENCES. 1. Downing, 1857:205. 2. *Horticulturist*, 12:179. 1857. 3. Warder, 1867:737. 4. Fitz, 1872:175. 5. *Am. Pom. Soc. Cat.*, 1873. 6. Leroy, 1873:871. figs. 7. Thomas, 1875:517. 8. Bailey, *Am. Hort.*, 1892:253. 9. Clayton, *Ala. Sta. Bul.*, 47:7. 1893. 10. Budd-Hansen, 1903:213.

SYNONYMS. YOPP (10). YOPP'S FAVORITE (1, 2, 3, 4, 5, 6, 7, 8, 9).

A southern apple of little value in New York. It originated in Georgia. In 1873 it was entered in the catalogue of the American Pomological Society (5). It is practically unknown in this state.

As grown at this Station the fruit is medium to rather large, oblate conic to roundish conic, somewhat ribbed; stem short, slender; cavity acuminate, moderately wide, rather deep, usually russeted; calyx small, open; basin narrow, moderately deep to deep, abrupt, wrinkled. Skin light yellow usually with a dull red blush; dots numerous, small, russet. Calyx tube conical; stamens median. Core medium to small; cells partly open; core lines clasping. Carpels broadly roundish, emarginate, somewhat tufted. Flesh whitish, somewhat tinged with yellow, moderately fine, tender, juicy, breaking, subacid, fair to good; season October and November.

The tree is below medium size, a rather slow grower with spreading top. It comes into bearing moderately early and yields good crops biennially.

YORK.

A variety which is known to many in Central and Western New York under the name York Pippin is now called by

pomologists Golden Pippin, under which name it is described on page 78. It is an apple of the Fall Pippin group, large, quite yellow when fully ripe, often with a brownish blush on the exposed cheek.

Fall Pippin has also been known to some under the name York Pippin. It is described on page 61.

Both of these are distinct from the York which is a Massachusetts apple of medium size, pale yellow with shade of red, good to very good for culinary uses. Season October and November.¹

¹ Downing, 1869, 420.

SIBERIAN CRABAPPLES AND THEIR HYBRIDS AND OTHER CRABAPPLES.

BAILEY CRIMSON.

REFERENCES. 1. *Am. Pom. Soc. Rpt.*, 1871:50. 2. *Montreal Hort. Soc. Rpt.*, 1879:91. 3. *Ib.*, 1884:38. 4. Bailey, *Am. Hort.*, 1892:234.

SYNONYMS. BAILEY'S CRIMSON (2, 4, of New York 3). BAILEY'S CRIMSON CRAB (1).

In 1871 the Committee of the American Pomological Society on Native Fruits made the following report concerning this variety: "Originated with Wm. H. Bailey, Plattsburgh, N. Y. Tree vigorous, upright, very productive and very handsome.

"Fruit medium or large for its class; roundish, inclining to conic; skin yellow, shaded over the whole surface with deep rich crimson; flesh similar to other Siberian crabs."

We have not seen this variety nor have we obtained any report concerning it from our correspondents.

BRIER.

REFERENCES. 1. *Rural N. Y.*, 1870 (cited by 2). 2. *Am. Pom. Soc. Rpt.*, 1871:51. 3. Downing, 1872:39 app. 4. *Wis. Hort. Soc. Rpt.*, 1876:70. 5. Lyon, *Mich. Hort. Soc. Rpt.*, 1881:322. 6. Barry, 1883:358. 7. Gibb, *Montreal Hort. Soc. Rpt.*, 1884:34. 8. Keffer, *S. D. Sta. Bul.*, 23:141. 1891. 9. Bailey, *Am. Hort.*, 1892:235. 10. Thomas, 1897:629. 11. Budd-Hansen, 1903:216. fig.

SYNONYMS. BRIAR SWEET (1, 2, 9). BRIER (11). BRIER'S SWEET (4, 5, 6, 7, 8). *Brier Sweet* (11, erroneously 10). BRIER'S SWEET CRAB (3). VAN WYCK (10).

Tree vigorous, hardy, comes into bearing rather young and is productive.

Historical. Originated with B. B. Brier, of Baraboo, Wis., as a result of the fertilization of the Siberian crab with the Bailey apple (3).

FRUIT.

Fruit large. *Form* roundish to conic, ribbed. *Stem* long, slender. *Cavity* narrow, deep, russeted. *Calyx* small, closed or slightly open. *Basin* deep, narrow, abrupt, wrinkled.

Skin pale yellow washed with lively red, striped with carmine, dotted and flecked with yellow and covered with thin, whitish bloom.

Calyx tube conical to funnel-form. *Stamens* median.

Core medium in size, axile or nearly so; cells closed.

Flesh yellowish, rich, fine-grained, moderately juicy, pleasant, sweet, aromatic, not astringent, good in flavor and quality.

Season September and October.

CHERRY.

REFERENCES. 1. Warder, 1867:715. 2. Downing, 1869:422. 3. Barry, 1883:359. 4. *Montreal Hort. Soc. Rpt.*, 1884:39. 5. Thomas, 1897:298. fig. 6. Budd-Hansen, 1903:217.

SYNONYMS. CHERRY (2, 5, 6). CHERRY CRAB (1, 3, 4).

Cherry is an old variety of unknown origin. The tree is a pretty good grower, particularly on light soils, attains considerable size and is quite a regular bearer.

TREE.

Tree moderately vigorous with long, slender, curved branches. *Form* upright spreading to roundish, open. *Twigs* moderately long, straight, moderately stout; internodes long. *Bark* clear reddish-brown tinged with olive-green, mottled with scarf-skin especially at the tips; slightly pubescent. *Lenticels* very scattering, small, roundish, not raised. *Buds* prominent, medium to large, plump, acute, free, not pubescent.

FRUIT.

Fruit small. *Form* oblate or roundish, ribbed. *Stem* long to very long, slender, bracted. *Cavity* rather broad, shallow, obtuse to slightly acute, somewhat russeted. *Calyx* medium to large, usually closed or eventually deciduous. *Basin* wide, shallow, obtuse, wrinkled.

Skin pale yellow nearly covered with bright red, often striped with carmine and overspread with a thin bluish bloom. *Dots* distinct, numerous, large, whitish or russet.

Calyx tube funnel-form. *Stamens* marginal.

Core large, axile; cells closed. *Carpels* broadly roundish or elliptical, emarginate, mucronate.

Flesh yellowish, rather coarse, juicy, crisp, mild subacid, somewhat astringent.

Season last of August to October.

CORAL.

REFERENCES. 1. Warder, *Tilt. Jour. Hort.*, 5:208. 1869. 2. Downing, 1869:423. 3. Barry, 1883:359. 4. *N. Y. Sta. An. Rpt.*, 2:35. 1884.

SYNONYMS. None.

Fruit of pretty good size, brilliant color, sprightly subacid flavor, in season from October to February. The tree is a pretty good grower, rather spreading, comes into bearing early and is a reliable cropper yielding good crops annually.

Historical. In 1869 Warder described this as No. 4 of the Marengo Winter Siberian crabapples received from Charles Andrews, Marengo, Ill. (1). It originated in the vicinity of Marengo (2). It is but little cultivated in New York.

FRUIT.

Fruit medium size or above, about an inch and a half in diameter. *Form* roundish to somewhat oblong, regular. *Stem* medium to rather long, slender, bracted. *Cavity* somewhat acute, medium in width and depth, regu-

lar, usually russeted. *Calyx* small, closed; lobes reflexed. *Basin* very shallow, broad and obtuse, or none.

Skin smooth, yellow, blushed with scarlet. *Dots* numerous, medium to small, gray or russet.

Calyx tube long, narrow, funnel-form. *Stamens* median.

Core medium to rather small, axile with narrow cylinder in the axis; cells closed or nearly so; core lines clasp the funnel cylinder. *Carpels* roundish ovate. *Seeds* compactly fill the cells; small to above medium, obtuse to somewhat acute, plump, dark.

Flesh yellow, breaking, juicy, crisp, sprightly, mild subacid to nearly sweet.

Season October to February.

CURRENT.

REFERENCES. 1. Downing, 1857:229. 2. Barry, 1883:359. 3. Bailey, *An. Hort.*, 1892:237. 4. Gibb, *Montreal Hort. Soc. Rpt.*, 1884:39.

SYNONYMS. CURRENT (2). CURRENT CRAB (1, 3). CURRENT CRAB (4). *Pomme Grosseille* (1).

Fruit small, borne in clusters; said to be hardier than Transcendent. Of no commercial value. The tree is a good grower, comes into bearing young and is productive.

TREE.

Tree moderately vigorous with moderately long, slender, curved branches. *Form* upright spreading or roundish, open. *Twigs* long, curved, slender; internodes short. *Bark* dark brown, lightly mottled with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous, medium size, roundish, slightly raised. *Buds* medium size, plump, acute, free, slightly pubescent.

FRUIT.

Fruit small or below medium. *Form* somewhat oblate, regular, uniform. *Stem* medium to long, rather slender. *Cavity* obtuse, rather deep, broad, symmetrical, frequently russeted. *Calyx* sometimes deciduous, medium size, closed; lobes rather narrow, acute. *Basin* rather deep, wide, abrupt, obscurely furrowed.

Skin thin, tough, smooth, glossy, yellow, striped with brilliant red, over-spread with bluish bloom. *Dots* numerous, small, pale or whitish.

Calyx tube broadly cone-shaped, short. *Stamens* marginal.

Core medium to rather large, axile; cells closed; core lines clasping. *Carpels* roundish to elliptical, emarginate. *Seeds* light brown, medium to large, wide, somewhat obtuse.

Flesh yellowish, firm, moderately fine, tender, dry, subacid, medium to poor.

Season October and November.

DARTMOUTH.

REFERENCES. 1. Barry, 1883:359. 2. Beach, *N. Y. Sta. An. Rpt.*, 15:277. 1896. 3. Lyon, *Mich. Sta. Bul.*, 143:200. 1897. 4. Farrand, *Ib.*, 205:47. 1903. 5. Ragan, *U. S. B. P. I. Bul.*, 36:363. 1905.

SYNONYMS. None.

Fruit large, brilliantly colored, good in flavor and quality. The tree is not a vigorous grower, comes into bearing rather early and yields full crops in alternate years.

Historical. Origin New Hampshire (1).

TREE.

Tree a moderately vigorous or rather slow grower with moderately long, stout, crooked branches. *Form* upright spreading to roundish, open. *Twigs* short, curved, moderately stout; internodes short. *Bark* clear brown, mingled with olive-green, lightly mottled with scarf-skin; pubescent near tips. *Lenticels* scattering, small, round, slightly raised. *Buds* rather prominent, medium to large, long, narrow, acute, free, slightly pubescent.

FRUIT.

Fruit medium to large. *Form* oblate or roundish oblate, ribbed. *Stem* long and slender, often bracted. *Cavity* acute, broad, deep, russeted. *Calyx* small; lobes long, reflexed. *Basin* rather broad, shallow.

Skin pale yellow, almost entirely overlaid with bright red deepening to a dark red or purple on the exposed side, dotted with yellow and covered with a heavy bluish bloom.

Calyx tube elongated cone-shape approaching funnel-form. *Stamens* marginal.

Core large, abaxile; cells open; core lines clasping.

Flesh yellowish, tinged with red next the skin, fine-grained, juicy, mild subacid, good in quality and flavor.


Season August.

EXCELSIOR.

REFERENCES. 1. *Ill. Hort. Soc. Rpt.*, 1880. 2. Gideon, *Am. Pom. Soc. Rpt.*, 1885:26. 3. *Rural N. Y.*, 45:184. 1886. figs. 4. Bailey, *Am. Hort.*, 1892:238. 5. Beach and Paddock, *N. Y. Sta. An. Rpt.*, 13:580. 1894. 6. Munson, *Me. Sta. An. Rpt.*, 12:73. 1896. 7. Lyon, *U. S. Pom. Bul.*, 6:11. 1897. 8. *Am. Pom. Soc. Cat.*, 1897:11. 9. Munson, *Me. Sta. An. Rpt.*, 18:83. 1902. 10. Thomas, 1903:348. 11. Budd-Hansen, 1903:218. 12. Farrand, *Mich. Sta. Bul.*, 205:47. 1903. 13. Powell and Fulton, *U. S. B. P. I. Bul.*, 48:41. 1903.

SYNONYMS. None.

Fruit very large for a crabapple being nearly as large as a medium sized apple. It is very attractive in appearance and excellent in quality for either dessert or culinary uses. As grown at this Station it appears to be one of the most desirable varieties of its class during early September. The tree is a good strong grower, hardy, healthy, comes into bearing rather young and yields full crops in alternate years.



EXCELSIOR (CRAB)

Historical. Originated by Peter M. Gideon, Excelsior, Minn. "It came from seed of Wealthy which had a chance to be crossed with the Cherry Crab and also the Oldenburg" (2).

TREE.

Tree large with long, moderately stout branches. *Form* somewhat flat and spreading, rather dense. *Twigs* short, curved, stout; internodes medium. *Bark* dark brown tinged with red; slightly pubescent. *Lenticels* quite numerous, medium size, oval, slightly raised. *Buds* medium to large, plump, obtuse, free, slightly pubescent.

FRUIT.

Fruit very large. *Form* roundish ovate to roundish oblate, symmetrical. *Stem* rather long and slender, sometimes bracted. *Cavity* small, acute or approaching acuminate, narrow, rather shallow, often slightly russeted. *Calyx* rather large, closed; lobes reflexed. *Basin* shallow, moderately broad, obtuse, furrowed.

Skin smooth, yellow, shaded and splashed with red over much of its surface. *Dots* numerous, russet.

Calyx tube wide, cone-shape. *Stamens* median or above.

Core large, decidedly abaxile; cells unsymmetrical, wide open; core lines clasping. *Carpels* elongated ovate, sometimes tufted. *Seeds* above medium size, long, moderately narrow, acute, tufted.

Flesh whitish, firm, a little coarse, crisp, juicy, subacid, with some Siberian crab flavor yet agreeable for dessert use, good to very good in quality.

Season early September.

FLORENCE.

REFERENCES. 1. Stark, *Mo. Hort. Soc. Rpt.*, 1886:233. 2. *Am. Pom. Soc. Rpt.*, 1887:134. 3. Lyon, *Mich. Sta. Bul.*, 118:59, 60. 1895. 4. Thomas, 1897:298. 5. Lyon, *Mich. Sta. Bul.*, 152:219, 224. 1898. 6. Budd-Hansen, 1903:219.

SYNONYMS. None.

This variety seems to be very desirable for commercial planting because the trees commence bearing very young, are reliable croppers and very prolific and the fruit is of good size, very attractive in appearance and of good quality. Although not superior to Martha in quality, Florence is more beautiful and more prolific.

Historical. Originated by Peter M. Gideon, Excelsior, Minn.

TREE.

Tree moderately vigorous. *Form* at first upright spreading but eventually inclined to droop. *Twigs* long, curved, moderately stout; internodes long.

Bark bright reddish-brown, slightly tinged with olive-green, mottled with scarf-skin; slightly pubescent. *Buds* large, broad, plump, obtuse, free, not pubescent.

FRUIT.

Fruit medium in size, uniform in size and shape. *Form* oblate, faintly ribbed. *Stem* very long, slender. *Cavity* acute, deep, medium in width, symmetrical, sometimes slightly russeted. *Calyx* variable, usually small, closed. *Basin* very shallow, rather wide, obtuse, slightly furrowed.

Skin moderately thin, moderately tough, smooth, yellowish-white mostly overspread with brilliant pinkish-red, sometimes with whitish bands radiating from the cavity, overspread with faint bloom. *Dots* minute, whitish.

Calyx tube moderately long, moderately wide, varying from somewhat urn-shape to funnel-form. *Stamens* marginal.

Core large; cells closed; core lines clasping. *Carpels* broadly obovate, emarginate. *Seeds* medium to rather small, moderately wide, flat, obtuse.

Flesh tinged with yellow, coarse, crisp, rather tender, juicy, very brisk subacid, somewhat astringent, good.

Season late August and early September.

GIBB.

REFERENCES. 1. *Montreal Hort. Soc. Rpt.*, 1884:35. fig. 2. *Am. Pom. Soc. Rpt.*, 1885:29. 3. *Stark, Mo. Hort. Soc. Rpt.*, 1886:233. 4. *Ill. Hort. Soc. Rpt.*, 1889:22. 5. *Craig, Can. Hort.*, 15:225. 1892. fig. 6. *Am. Pom. Soc. Cat.*, 1897:11. 7. *Thomas*, 1897:298. fig. 8. *Farrand, Mich. Sta. Bul.*, 205:47. 1903. 9. *Powell and Fulton, U. S. B. P. I. Bul.*, 48:42. 1903. 10. *Budd-Hansen*, 1903:219. fig.

SYNONYMS. None.

Fruit large, yellow blushed with dull red. It is thinner skinned and much less astringent than Hyslop with remarkably yellow flesh. It is highly esteemed for canning; season last half of September. It is recommended for the home orchard and is worthy of trial for commercial planting where a crabapple of its season is desired. The tree is a slow, spreading grower, fairly hardy as far north as Montreal, very productive (5).


Historical. Originated with George P. Peffer, Pewaukee, Wis., being a seedling of an oblate Yellow Siberian crab crossed with Fall Greening (5, 10).

FRUIT (5, 7, 10).

Fruit large. *Form* roundish oblate. *Stem* short to medium length, thick. *Cavity* wide, deep, regular. *Calyx* medium size, open. *Basin* very wide, shallow, wrinkled.

Skin thin, yellow, blushed with dull red, attractive. *Dots* white, minute. *Flesh* remarkably yellow, firm, crisp, juicy, pleasantly acid, a little astringent, sprightly.

Season early.



HYSLOP (CRAB)

HYSLOP.

REFERENCES. 1. Warder, *Tilt. Jour. Hort.*, 1869:206. figs. 2. Downing, 1869:424. 3. Fitz, 1872:147. 4. Gibb, *Montreal Hort. Soc. Rpt.*, 1876:21. 5. Minn. *Hort. Soc. Rpt.*, 1876:110. 6. Barry, 1883:359. 7. Gibb, *Montreal Hort. Soc. Rpt.*, 1884:39. 8. Thomas, 1885:513. 9. Wickson, 1889:249. 10. Bailey, *Am. Hort.*, 1892:241. 11. Munson, *Me. Sta. Rpt.*, 1893:134. 12. *La. Sta. Bul.*, 27:926. 1894. 13. *Can. Hort.*, 26:489. 1903. figs. 14. Budd-Hansen, 1903:220. fig.

SYNONYMS. HYSLOP (1). HYSLOP (2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14). HYSLOP'S CRAB (8).

Fruit large, very brilliantly colored, dark red or purplish over-spread with thick blue bloom; borne in clusters. The tree is a good grower, very hardy, and a reliable cropper yielding good crops biennially or in some cases annually. It is desirable both for home use and for market.

Historical. Origin unknown. In 1869 Downing remarked: "This variety has been long and pretty extensively cultivated" (2). It is commonly listed by nurserymen throughout the country (10). It is one of the best known and most widely cultivated of the crabapples.

TREE.

Tree vigorous or moderately vigorous. *Form* upright spreading, rather open. *Twigs* long, curved, slender; internodes long. *Bark* olive-green, tinged with reddish-brown, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* numerous, very conspicuous, greenish-yellow, medium to large, oblong. *Buds* exceptionally large and prominent, very long, narrow, plump, acute, free, slightly pubescent.

FRUIT.

Fruit above medium to large, very uniform in size and shape. *Form* roundish ovate or obovate, sometimes a little inclined to oblong, regular or obscurely ribbed, symmetrical. *Stem* rather short to very long, slender. *Cavity* acuminate, rather small, shallow, narrow to medium in width, sometimes furrowed, often slightly russeted. *Calyx* medium to large, closed; lobes medium to long, narrow, acuminate, reflexed. *Basin* shallow, medium to wide, distinctly furrowed and wrinkled.

Skin clear pale yellow almost completely overspread with lively dark red shading to deep carmine or purplish carmine and covered with thick, blue bloom. *Dots* small, numerous, pale or gray.

Calyx tube short, narrow, cone-shape to urn-shape. *Stamens* median.

Core medium size, axile; cells symmetrical, closed; core lines meeting. *Carpels* elongated ovate, emarginate. *Seeds* small, narrow, short, plump, obtuse to acute, medium brown.

Flesh yellow, sometimes with tinge of red next the skin, very firm, mod-

erately fine, at first juicy but eventually becoming dry and mealy, subacid, astringent, good for culinary purposes.

Season late September and October.

LARGE RED SIBERIAN.

REFERENCES. 1. Downing, 1845:147. 2. Cole, 1849:137. 3. Fitz, 1872:147. 4. Barry, 1883:359. 5. Wickson, 1889:249. 6. Bailey, *An. Hort.*, 1892:243.

SYNONYMS. None.

This fruit is of medium size for a Siberian crab, being larger than Red Siberian, but smaller than either Transcendent or Hyslop. It is similar to Red Siberian in appearance and quality. The foliage is coarser than that of Red Siberian, and the tree is larger, being medium to rather large, a vigorous grower, erect or roundish, with long, slender twigs. It is very hardy, healthy, moderately long-lived, and a reliable cropper, yielding heavy crops biennially or sometimes annually. It has long been known in cultivation, and is still listed by nurserymen (6), but larger and handsomer varieties of more recent introduction are generally preferred in market.

FRUIT.

Fruit of medium size, uniform in size and shape. *Form* roundish to roundish ovate, regular. *Stem* medium to long, slender. *Cavity* acuminate, shallow, moderately broad, often furrowed, usually russeted. *Calyx* medium size, closed; lobes long, narrow, acuminate. *Basin* shallow or none, obtuse, wrinkled, having mammiiform protuberances.

Skin thin, tough, smooth, pale yellow, almost wholly overlaid with bright red and marked with obscure narrow stripes of dark red. *Dots* very small, light, inconspicuous.

Calyx tube short, wide, urn-shape. *Stamens* median to marginal.

Core medium size, axile; cells closed; core lines meeting. *Carpels* ovate to obovate, emarginate. *Seeds* glossy, dark brown, rather small, short, wide, obtuse.

Flesh yellowish, very firm, subacid, astringent, good for culinary uses.


Season September and October.

LARGE YELLOW SIBERIAN.

REFERENCES. 1. Warder, 1867:732. 2. Downing, 1869:425. 3. Barry, 1883:360. 4. Bailey, *An. Hort.*, 1892:243.

SYNONYMS. None.

Fruit large, similar in size to Large Red Siberian, clear pale yellow with a shade of red in the sun, roundish approaching oblong truncate; season September and October. Tree upright, somewhat irregular in form, of me-



LARGE RED SIBERIAN (CRAB)

dium size, vigorous or moderately vigorous, very hardy, healthy, comes into bearing young and is very productive. It is generally superseded in market by larger varieties.

MARENGO.

REFERENCES. 1. Elliott, *Horticulturist*, 23:136. 1868. fig. 2. Warder, *Tilt. Jour. Hort.*, 5:207. 1869. fig. 3. Downing, 1869:425. 4. *Horticulturist*, 26:371. 1871. 5. *Ill. Hort. Soc. Rpt.*, 1877:112. 6. *Ib.*, 1879:196. 7. Barry, 1883:360. 8. *Montreal Hort. Soc. Rpt.*, 1884:37. 9. Thomas, 1885:226, 517. 10. Bailey, *Am. Hort.*, 1892:244. 11. *Me. Sta. Rpt.*, 1893:134. 12. Budd-Hansen, 1903:222.

SYNONYMS. MARENGO (5, 6). MARENGO CRAB (3, 7, 9, 10, 11, 12). MARENGO NO. 1 (2, 8). MARENGO WINTER CRAB (1, 4). *Marengo Winter Crab* (2).

Originated at Marengo, Ill. (3, 6). This is a good variety for home use where a late-keeping crabapple is desired, and some have found it a profitable market variety but other varieties of its season which are more attractive in color are generally preferred for commercial purposes. The tree is of medium size, vigorous, spreading, very hardy, long-lived and a reliable cropper usually yielding heavy crops annually. The limbs are very tough and support heavy loads well. The fruit hangs well to the tree till very late in the season. It agrees well with the following description given by Warder (2): "Fruit globular, truncate, regular, or slightly flattened on the sides, one of the largest of its class; surface smooth, yellow, blushed or covered with crimson; dots minute; basin shallow, folded; eye small, closed; cavity medium, regular; stem long; core large, closed, meeting the eye; seeds few, small, plump, light brown; flesh yellow, firm, rather juicy; flavor subacid, aromatic. Use, kitchen and dessert; quality, good; season, winter, and till spring in the North."

MARTHA.

REFERENCES. 1. ? *Ia. Hort. Soc. Rpt.*, 1879:452. 2. *Ib.*, 1880:68. 3. *Rural N. Y.*, 45:284. 1886. figs. 4. *Am. Pom. Soc. Rpt.*, 1887:134. 5. Bailey, *Am. Hort.*, 1892:244. 6. Thomas, 1897:299. 7. *Am. Pom. Soc. Cut.*, 1897:11. 8. Budd-Hansen, 1903:222. 9. *Rural N. Y.*, 62:235. 1903. 10. Farrand, *Mich. Sta. Bul.*, 205:47. 1903.

SYNONYMS. None.

Fruit large, very handsome clear yellow more or less overspread with bright red; excellent in flavor and quality; one of the very best of its class for all culinary purposes. The tree is of medium size, moderately vigorous, roundish or spreading, very hardy, comes into bearing young, and is a reliable cropper, yielding good to heavy crops annually or nearly annually. The fruit hangs well to the tree, is uniform, reliable, and satisfactory in appearance and

quality. Season, September to late fall. It should be more generally grown in New York.

Historical. Originated with Peter M. Gideon, Excelsior, Minn. It has as yet been grown but little in this state.

TREE.

Tree medium in size, moderately vigorous. *Form* spreading, open and somewhat inclined to droop. *Twigs* long, curved, slender; internodes long. *Bark* reddish-brown tinged with green, lightly streaked with scarf-skin, not pubescent. *Lenticels* quite numerous, small, oblong, not raised. *Buds* very prominent, large, long, acute, free, not pubescent. *Leaves* rather long, somewhat twisted and drooping.

FRUIT.

Fruit usually rather large, uniform in size and shape. *Form* roundish or oblate, regular or very faintly ribbed, usually symmetrical; sides sometimes unequal. *Stem* long, slender. *Cavity* acute to somewhat obtuse, medium in depth to rather shallow, rather broad, sometimes furrowed, often thinly russeted. *Calyx* medium to small, closed or partly open, occasionally deciduous. *Basin* shallow, wide, obtuse, smooth.

Skin moderately thin, tough, smooth, clear pale yellow almost entirely covered with an attractive bright light red overspread with bluish bloom; sometimes faint narrow stripes extend from the cavity to calyx. *Dots* rather numerous, light-colored, small to medium size.

Calyx tube short, rather narrow, very small, conical or somewhat funnel-form. *Stamens* median to marginal.

Core medium size, axile; cells closed or nearly so. *Carpels* roundish or somewhat obovate, slightly tufted. *Seeds* medium size, rather narrow, acute to acuminate.

Flesh yellowish, firm, moderately coarse, crisp, juicy, rather brisk subacid, good to very good in flavor and quality.

Season September to November.

MINNESOTA.

REFERENCES. 1. Gideon, *Horticulturist*, 27:244. 1872. 2. Gibb, *Am. Pom. Soc. Rpt.*, 1883:125. 3. *Ib.*, *Montreal Hort. Soc. Rpt.*, 10:36. 1884. 4. Bailey, *Am. Hort.*, 1892:244. 5. Beach, *N. Y. Sta. An. Rpt.*, 15:277. 1896. 6. Budd-Hansen, 1903:223. 7. Thomas, 1903:349.

SYNONYMS. MINNESOTA CRAB (2, 3, 4, 5, 6, 7). MINNESOTA (1).

This variety originated in Minnesota. The fruit is very large for its class, roundish; skin pale yellow blushed or mottled on the sunny side and overspread with thin whitish bloom; flesh white, firm, crisp, juicy, fine-grained, subacid to mild subacid or nearly sweet, slightly astringent, good. Season September and October. As grown at this Station the tree is of spreading form, moderately vigorous and not very productive.



MARTHA (CRAB)

MONTREAL BEAUTY.

REFERENCES. 1. Downing, 1869:425. 2. Gibb, *Montreal Hort. Soc. Rpt.*, 1876:17. 3. *Ill. Hort. Soc. Rpt.*, 1877:112. 4. Barry, 1883:360. 5. *Montreal Hort. Soc. Rpt.*, 1884:39. 6. Wickson, 1889:249. 7. Bailey, *Am. Hort.*, 1892:244. 8. *Me. Sta. Rpt.*, 1893:134. 9. Thomas, 1897:299. 10. Budd-Hansen, 1903:223.

SYNONYMS. MONTREAL (10). *Montreal Beauty* (10). MONTREAL BEAUTY CRAB (1, 2, 3, 4, 5, 6, 7, 8, 9).

This very beautiful fruit originated in Quebec and was propagated as early as 1833. Gibb (2, 5) states the tree is less hardy than that of Transcendent, but is strong, vigorous, rather large, at first very close and upright, does not come into bearing early, but bears heavily. The fruit is large for its class, oblong conic to roundish oblong and truncate, yellowish-green, mostly covered with red; flesh yellowish-white, tender, rather juicy, unless overripe, subacid, very slightly astringent; season, late September and October. It is well known in the vicinity of Montreal and other portions of Quebec. In the United States it is quite frequently listed by nurserymen (7), yet it is not generally known.

OBLONG.

REFERENCES. 1. Desportes, *Horticulturist*, 6:507. 1851. fig. 2. Downing, 1869:425. 3. Barry, 1883:360.

SYNONYMS. *Baccata fructa oblonga* (1). OBLONG-FRUITED SIBERIAN CRAB (2). OBLONG SIBERIAN CRAB (1, 3).

Fruit medium size, oblong conic, yellowish-white, partly shaded with crimson, striped with carmine and covered with thin bloom; stem very long, slender, usually bracted. This variety has been sparingly disseminated in New York but is now seldom or never planted here. It is not superior to other varieties of its season.

ORANGE.

REFERENCES. 1. Downing, 1869:425. 2. *Montreal Hort. Soc. Rpt.*, 1884:36. fig. 3. Thomas, 1897:299. fig. 4. Budd-Hansen, 1903:224.

SYNONYMS. None.

A variety of American origin but little grown in New York. Some regard it as a desirable variety for both home use and market. The tree is roundish, spreading, dwarfish, a moderate or rather slow grower, hardy, moderately long-lived and a reliable cropper yielding good to heavy crops annually. It comes into bearing rather young. Season September to November.

FRUIT (1, 4).

Fruit medium size. *Form* roundish, slightly oblate. *Stem* very long, slender. *Cavity* open, deep, acute with trace of russet. *Calyx* closed. *Basin* very shallow or flat, wrinkled.

Skin orange-yellow often netted with russet. *Dots* minute, white, obscure.

Core open.

Flesh light salmon-yellow, a little dry, rather mild subacid with sweet after-taste, good.

Season September to November.

PAUL IMPERIAL.

REFERENCES. 1. Ellwanger and Barry, *Cat.*, 1888:14. 2. Thomas, 1897:299.
SYNONYMS. None.

A September variety in season about with Transcendent; somewhat irregular in shape; of very good general appearance but less attractive in size and color than Hyslop and inferior to Martha in quality. The tree is below medium size, comes into bearing rather young and is a reliable annual cropper.

Historical. Introduced by Paul and Son, Cheshunt, England. A cross between the Red Astrachan and Siberian Crab (1).

TREE.

Tree moderately vigorous with short, stout, crooked branches having numerous small spurs. *Form* spreading, flat, open. *Twigs* olive-green tinged with brown, lightly streaked with scarf-skin; slightly pubescent. *Lenticels* scattering, medium to large, oval, slightly raised. *Buds* large, prominent, plump, obtuse, free, pubescent.

FRUIT.

Fruit small to medium, uniform in size but not in shape. *Form* usually oblate, often irregularly elliptical, strongly ribbed; sides unequal. *Stem* long to medium, slender. *Cavity* obtuse to acute, moderately deep, broad, furrowed, not russeted. *Calyx* large, closed, prominent, persistent; lobes long, moderately broad. *Basin* very shallow, wide, obtuse, furrowed and wrinkled and sometimes mammillate.




Skin thin, rather tender, smooth, yellow, often entirely covered with dark bright red or with but little of the yellow ground color exposed, over-spread with blue bloom. *Dots* very small, numerous, indistinct, light.

Calyx tube small, short, moderately wide, urn-shape. *Stamens* nearly marginal.

Core rather large, axile or nearly so; cells closed or slightly open; core lines meeting. *Carpels* nearly roundish, narrowing toward apex, tufted. *Seeds* below medium size, moderately wide, acute, light brown.

Flesh yellowish sometimes stained with pink, firm, moderately coarse, crisp, rather tough, juicy, brisk subacid, less astringent than Red Siberian, good.

Season September and October.



MONTREAL BEAUTY (CRAB)

PICTA STRIATA.

REFERENCES. 1. Ellwanger and Barry, *Cat.*, 1888:14. 2. Beach, *N. Y. Sta. Am. Rpt.*, 12:601. 1893. 3. Ragan, *U. S. B. P. I. Bul.*, 56:370. 1905.
SYNONYMS. PICTA STRIATA (1, 2). PIETA (3). *Picta Striata* (3).

Fruit handsome, rather mild in flavor; season late fall and early winter. It is hardly large enough for a good commercial variety. The tree is a good grower, comes into bearing rather late and is an annual cropper yielding moderate to good crops.

Historical. Received from Ellwanger and Barry, Rochester, N. Y., in 1888 for testing at this Station.

TREE.

Tree rather large. *Form* upright spreading to roundish with rather drooping laterals. *Twigs* long, curved, slender; internodes short. *Bark* clear brown, tinged with green, lightly mottled with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous, rather conspicuous, medium size, oval, not raised. *Buds* medium size, plump, acute, free, not pubescent.

FRUIT.

Fruit medium or above, uniform in size but not in shape. *Form* oblate or roundish oblate, irregularly ribbed. *Stem* long, slender. *Cavity* obtuse, medium to rather deep, medium to broad, compressed, smooth or nearly so. *Calyx* usually small, closed; lobes separated at base, long, medium in width, acute to acuminate. *Basin* shallow to medium in depth, medium in width, obtuse, nearly smooth.

Skin thin, tender, smooth, rather glossy, pale greenish-yellow nearly covered with crimson, blushed and striped with carmine. *Dots* indistinct, small, gray.

Calyx tube moderately short, rather narrow, conical to urn-shape. *Stamens* marginal or nearly so.

Core medium to large, axile; cells slightly open or closed; core lines clasping. *Carpels* roundish to elliptical, concave, deeply emarginate, sometimes tufted. *Seeds* rather dark brown, medium size, wide, short, obtuse.

Flesh tinged with yellow, firm, a little coarse, tender, juicy, somewhat astringent, sprightly subacid, good.

Season October to early winter.

QUAKER.

REFERENCES. 1. *Rural N. Y.*, 1870 (cited by 7). 2. Bailey, *An. Hort.*, 1892:247. 3. Thomas, 1897:299. 4. Lyon, *Mich. Sta. Bul.*, 152:224. 1898. 5. Farrand, *ib.*, 205:48. 1903. 6. Budd-Hansen, 1903:224. 7. Ragan, *U. S. B. P. I. Bul.*, 56:371. 1905.

SYNONYMS. None.

This variety has been disseminated more in the western states than it has in New York. It is but little grown here. Farrand (5) describes it as

a late ripening variety of only fair quality, size medium to large; color yellow with a red cheek; tree handsome, vigorous but not very productive; season October.

QUEEN CHOICE.

REFERENCES. 1. Bailey, *An. Hort.*, 1892:247. 2. Ellwanger and Barry, *Cat.*, 1894:15. 3. Thomas, 1897:299. 4. Budd-Hansen, 1903:224.

SYNONYMS. QUEEN'S CHOICE (1, 2, 3). QUEEN CHOICE (4).

This crab has a fruit which is medium or above, roundish conical, of a beautiful crimson color, showy and attractive; flesh whitish and of pleasant flavor and quality; season October; tree vigorous, very prolific (2, 3, 4).

It is but little known in New York.

RED SIBERIAN.

REFERENCES. 1. Forsyth, 1803:57. 2. Cox, 1817:106. 3. Fessenden, 1828:129. 4. Kenrick, 1832:57. 5. *Ib.*, 1835:75. 6. Downing, 1845:147. 7. Thomas, 1849:158. 8. Cole, 1849:137. 9. Barry, 1851:287. 10. Elliott, 1854:157. 11. Gregg, 1857:62. *fig.* 12. Downing, 1869:426. 13. Fitz, 1872:147. 14. Thomas, 1875:208. 15. *Montreal Hort. Soc. Rpt.*, 10:39. 1884. 16. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:300. 17. Bailey, *An. Hort.*, 1892:248. 18. Lyon, *U. S. Pom. Bul.*, 6:11. 1897. 19. Budd-Hansen, 1903:224.

SYNONYMS. RED SIBERIAN (11, 15, 16, 17, 18, 19). RED SIBERIAN CRAB (5, 8, 9, 12). SIBERIAN CRAB (1, 2, 3, 4, 6, 7, 10, 13, 14).


Origin France (19). Fruit small, decidedly ornamental, borne in clusters. It is three-quarters of an inch to an inch in diameter, roundish oblate to somewhat oblong, irregularly elliptical; stem long and slender; cavity acute, medium in width and depth; calyx small to medium, often deciduous; basin but slightly depressed. Skin smooth, pale yellow striped and blushed with lively red and overspread with blue bloom; flesh subacid, astringent, good for culinary uses. Season September and October.

SEPTEMBER.

REFERENCES. 1. *Mich. Hort. Soc. Rpt.*, 1888:319. 2. Beach, *N. Y. Sta. An. Rpt.*, 12:602. 1893.

SYNONYMS. None.

A very handsome fruit of good quality for either dessert or culinary use. It ripens a few days later than Transcendent.



SMALL RED SIBERIAN

The tree is a good grower, comes into bearing young and yields full crops biennially.

This is distinct from the September apple described by Downing.¹

Historical. Originated with Peter M. Gideon, Excelsior, Minn., from seed of Cherry Crab. In 1888 Mr. Gideon sent stock of this variety to this Station for testing.

TREE.

Tree vigorous, with short, stout, crooked and twisted branches. *Form* rather flat, spreading, open. *Twigs* moderately long, curved, moderately stout with large terminal buds; internodes long. *Bark* clear brown, lightly mottled with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous, medium size, round, not raised. *Buds* prominent, large, long, plump, acute, free, not pubescent.

FRUIT.

Fruit medium to large for a crab, uniform in size but not in shape. *Form* roundish oblate to somewhat oblong, inclined to conic, frequently ribbed; sides usually unequal. *Stem* usually long and slender. *Cavity* somewhat obtuse, rather shallow to medium in depth, medium to broad, occasionally furrowed, sometimes russeted. *Calyx* medium to rather large, closed or partly open. *Basin* mammillate, very shallow to moderately deep, medium in width, slightly obtuse.

Skin thin, tender, smooth, pale yellow or greenish, striped with red, in well colored specimens becoming nearly covered with dark red, overspread with bluish bloom. *Dots* small, scattering, inconspicuous, gray or brownish.

Calyx tube rather short, medium in width, broadly funnel-form; pistil point persistent. *Stamens* median to nearly marginal.

Core medium size, axile or slightly abaxile; cells closed or wide open; core lines clasping. *Carpels* roundish, elongated. *Seeds* variable in shape, medium size, moderately wide, rather short, acute to broadly acute, rather light dull brown.

Flesh tinged with yellow, medium in grain, moderately tender, juicy, subacid, not astringent but with an agreeable crabapple flavor, good to very good in quality.

Season September.

SOULARD.

REFERENCES. 1. Soulard, *Gard. Monthly*, 10:199. 1868 (cited by 13). 2. *Ib.*, *Ill. Hort. Soc. Rpt.*, 1869:195, 260. 3. Downing, 1869:426. 4. *Am. Pom. Soc. Cat.*, 1871:8. 5. Soulard, *Ill. Hort. Soc. Rpt.*, 1873:291. 6. *Montreal Hort. Soc. Rpt.*, 5:91. 1879. 7. *Ib.*, 10:37. 1884. 8. Thomas, 1885:524. 9. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:300. 10. Bailey, *Am. Hort.*, 1892:249. 11. Budd, *Amer. Gard.*, 14:244. 1893. 12. Thomas, 1897:264. fig. 13. Bailey, *Evol. Native Fruits*, 1898:261. fig. 14. Craig, *la. Acad. Sci.*, 7:130. 1899. pl. 15. Budd-Hansen, 1903:225. fig.

SYNONYMS. None.

¹ Downing, 1869:350.

This is regarded by Bailey as a hybrid between the native prairie crab, *Pyrus ioensis*, and the common apple, *Pyrus malus* (13), in which opinion he is supported by Craig (14). It is distinct from the Soulard apple. According to Soulard's account (13) "it originated on a farm about twelve miles from St. Louis, Mo., where stood an American crab thicket not inclosed, near the farmhouse, about twenty-five years since. The thicket was cut down and the ground cultivated some two or three years; culture being discontinued, another crab thicket sprang up, and when bearing, one tree (the identical kind now called Soulard crab) was discovered. The fruit astonished me by its remarkably large size, being sent to me by a friend whose widowed mother, Mrs. Freeman Delauriere, occupied the farm. I immediately propagated it by grafting upon crab stock and upon our common seedlings. Upon both stocks producing the same fruit and thriving admirably, I disseminated it among my friends as a very desirable fruit, having nothing of the Siberian type. It is to me conclusive that this crab is the offspring of an accidental hybridization of the wild crab by our common apple. The tree, its foliage, habit, increased size of fruit and tree, and decreased acerbity, convince me it is a hybrid, and as far as I know, the first instance of such cross.

"I consider it the most desirable of all crabs that I have seen. Adding sweetness, it is delicious baked. It makes most excellent preserves, being large enough to be quartered, and unsurpassed by any crab for jams, jellies, etc., imparting its delicate taste and rich crab aroma. The largest have measured over seven inches around. In form, color and smell it is like the common crab, and it hangs on the tree until destroyed by frost. It will keep two years, with common care, in a cellar, and will stand repeated freezing and thawing in a dark place. It is agreeable to many palates in the spring.


"The tree is an immense grower in the nursery, coming early into fruit and making but little growth afterward, and is an immense and regular bearer. I have made some cider as clear as wine, with sugar or a quarter part of sweet apples. It will make delicious strong cider. Tree perfectly hardy, having stood the severest winters here and at St. Paul, Minn., for twenty-five years."

After giving the testimony of several observers with regard to the value of this hybrid for the Upper Mississippi valley Bailey concludes: "It is probable that too much was expected of the Soulard crab when it was first introduced, and that it afterward, suffered from the partial collapse. Such an array of apples has now been introduced into the cold Northwest—from the East, from Russia, offspring of the Siberian crab, and local seedlings of the common apple—that the Soulard crab and its kin have been obscured" (13).

The variety is still listed by some nurserymen (10). It is practically unknown in New York and has no value for this state.

TRANSCENDENT.

REFERENCES. 1. *Horticulturist*, 22:125. 1867. 2. Warder, *Tilt. Jour. Hort.*, 5:205. 1869. fig. 3. Downing, 1869:426. 4. Todd, 1871:83. fig. 5. Fitz,



TRANSCENDENT (CRAB)

1872:147. 6. Gibb, *Montreal Hort. Soc. Rpt.*, 1:21. 1876. 7. Barry, 1883:360. 8. Gibb, *Montreal Hort. Soc. Rpt.*, 10:39. 1884. 9. Wickson, 1889:249. 10. Bailey, *Am. Hort.*, 1892:251. 11. Thomas, 1897:300. 12. *U. S. Pom. Bul.*, 6:11. 1897. 13. *Can. Hort.*, 26:489. 1903. figs. 14. Budd-Hansen, 1903:226. fig.

SYNONYMS. TRANSCENDANT (5, 9, 10, 13). TRANSCENDENT (1, 2, 3, 4, 6, 7, 8, 11, 12, 14).

This beautiful fruit has for many years been one of the most popular of the crabapples cultivated in this state. The tree is a good grower, roundish, spreading, hardy and usually very productive yielding good to very heavy crops nearly annually. It is in season from late August to the middle of September or a little later.

Historical. The history of Transcendent seems to be unknown. Although our first reference to this variety is 1867, William Prince had it listed in his nursery catalogue as early as 1844.¹ It seems to have been in the hands of nurserymen for years before it came to the attention of pomological writers, hence the obscurity in regard to its origin.

TREE.

Tree large with stout, curved and drooping branches. *Form* very spreading, drooping, rather dense. *Twigs* moderately long, curved, slender, with large terminal buds; internodes long. *Bark* brown, tinged with green, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* quite numerous, medium to large, oval, raised, conspicuous. *Buds* medium to large, rather prominent, plump, obtuse or slightly acute, free, slightly pubescent.

FRUIT.

Fruit medium to rather large. *Form* roundish or roundish oblong, flattened at the ends, somewhat ribbed. *Stem* medium to long, rather stout, bracted. *Cavity* narrow, shallow, obtuse. *Calyx* large, closed; lobes long, leafy, reflexed. *Basin* shallow, wrinkled.

Skin thin, clear bright yellow with bright red cheek, overspread with bloom. Highly colored specimens are nearly covered with bright red.

Calyx tube conical. *Stamens* marginal.

Core medium size; cells closed.

Flesh yellow, crisp, juicy, moderately fine, somewhat astringent, subacid, very good for culinary uses.

Season late August to the middle of September.

VAN WYCK.

REFERENCES. 1. Downing, 1872:39 app. 2. Barry, 1883:360. 3. *Montreal Hort. Soc. Rpt.*, 10:37. 1884. 4. Bailey, *Am. Hort.*, 1892:251. 5. *Am. Pom. Soc. Cal.*, 1897:11. 6. Thomas, 1897:300. fig. 7. Budd-Hansen, 1903:227.

¹ Ragan, *U. S. P. B. I. Bul.*, 66:373. 1905.

SYNONYMS. *Brier Sweet* (6). VAN WYCK CRAB (2, 4, 5, 6, 7). VAN WYCK SIBERIAN (1). VAN WYCK SWEET (3, 4).

A sweet crabapple which according to Downing (1) originated as a chance seedling in Fishkill, Dutchess county, N. Y. Fruit large for a Siberian crabapple, whitish shaded with bright red, covered with bloom; flesh whitish, tender, moderately juicy, sweet, rich; it sometimes water-cores; core small, closed; season August and September (1, 2). This variety is occasionally listed by nurserymen (4). It is not generally known in New York and it appears that its cultivation in this state is not increasing.

WHITNEY.

REFERENCES. 1. Warder, *Till. Jour. Hort.*, 5:206. 1869. fig. 2. *Minn. Hort. Soc. Rpt.*, 1876:109. 3. *Am. Pom. Soc. Cat.*, 1881:16. 4. Barry, 1883:300. 5. Gibb, *Montreal Hort. Soc. Rpt.*, 1884:34. fig. 6. *ib.*, *Am. Pom. Soc. Rpt.*, 1885:29. 7. Thomas, 1885:529. 8. *Can. Hort.*, 11:42. 1888. 9. Wickson, 1889:249. 10. Lyon, *Mich. Hort. Soc. Rpt.*, 1890:300. 11. Bailey, *Am. Hort.*, 1892:252. 12. Beach, *N. Y. Sta. An. Rpt.*, 13:584. 1894. 13. Burrill and McCluer, *Ill. Sta. Bul.*, 45:346. 1896. 14. Farrand, *Mich. Sta. Bul.*, 205:48. 1903. 15. *Can. Hort.*, 26:489. 1903. figs. 16. Budd-Hansen, 1903:228.

SYNONYMS. WHITNEY (3, 10, 11, 14, 15, 16). WHITNEY CRAB (4, 7, 9). WHITNEY No. 20 (1, 2, 5, 6, 8, 11, 12, 13). *Whitney No. 20* (15, 16).

One of the most popular of the large crabapples particularly in the West and North. The fruit is attractive, yellow, striped with lively red, subacid, good for dessert and very good for culinary uses. It is in season in August and early September. The tree is a thrifty, upright grower, comes into bearing young and is very productive.

Historical. This variety originated with A. R. Whitney, Franklin Grove, Ill. (1). It was at first disseminated under the name Whitney No. 20, under which name it was described by Warder as early as 1869 (1). It has not been much planted in New York but in regions farther north and west its cultivation is gradually increasing.

TREE.

Tree below medium with moderately stout, moderately long, curved branches. *Form* upright becoming somewhat spreading after fruiting heavily. *Twigs* short, straight, stout with large terminal buds; internodes medium size. *Bark* bright brown tinged with green, lightly streaked with scarf-skin; slightly pubescent near tips. *Lenticels* numerous, small, oval, not raised. *Buds* medium to large, broad, obtuse, free, pubescent.

FRUIT.

Fruit large, pretty uniform in size and shape. *Form* roundish inclined to conic or to ovate. *Stem* slender. *Cavity* narrow to moderately wide,

rather deep, obtuse. *Calyx* medium to large, closed or partly open. *Basin* broad, shallow, wrinkled.

Skin light yellow largely shaded and striped with red.

Flesh yellowish, crisp, juicy, mild subacid or nearly sweet with slight crab-apple flavor, good to very good.

Season late August and early September.

YELLOW SIBERIAN.

REFERENCES. 1. Buel, *N. Y. Bd. Agr. Mem.*, 1826: 477. 2. Kenrick, 1832: 58. 3. Downing, 1845: 137. 4. Cole, 1849: 137. 5. Thomas, 1849: 158. 6. Waring, 1851: 30. 7. Barry, 1851: 287. 8. Gregg, 1857: 62. 9. Warden, 1867: 712. 10. Hadd Hansen, 1903: 228.

SYNONYMS. *Amber Crab* (3). *Golden Beauty*. *SIBERIAN CRAB* (6). *YELLOW SIBERIAN* (1, 2, 3, 4, 5, 7, 8, 9, 10).

This is sometimes called *Golden Beauty*. It is similar to *Red Siberian*, except in the color and size of the fruit, it being rather large and of a clear golden-yellow color. *Season* September. *Tree* medium size or below, a good grower, upright becoming roundish, and somewhat drooping, very hardy and healthy except that under certain circumstances it suffers from blight. It comes into bearing rather young and is a reliable cropper, yielding heavy to very heavy crops annually or nearly annually. It is grown principally for home use but sometimes a portion of the fruit is disposed of in local markets.

ERRATA—VOL. I.

- Page 24. Under Rhode Island Greening group, "Northwestern Greening" should be omitted.
25. Under Alexander group, "Bismark" should be "Bismarck."
71. Under Different Types of Ben Davis, in the last paragraph omit Rutledge.
76. Under Black Annette, second line, "1886" should be "1866."
95. First line of text, under Carlough, (2) should be (3).
141. Under Golden Medal, fourth line, reference "U. S. B. P. I. bulletin 56:124" should be "U. S. B. P. I. bulletin 56:125."
168. Colored plate facing. Legend "Jacob Sweet" should be "Jacobs Sweet."
196. McAfee, first line of second paragraph, insert (7). between Indian and Wyandotte; Wyandotte (7). should precede Zecke on p. 197.
270. Colored plate facing. "Genet," in italic should be omitted.
333. Under Sweet Russet all of the second line, including the words "Synonym" and "Summer Russet (1)," should be omitted.
392. Twenty-sixth line, second column, "Bismark" should be "Bismarck."
403. "Pomme Grise," first column, fifteenth line, should be roman instead of italic.
409. Eighth line from bottom of left column, ' Winter Sweet Paradise' should be roman capitals and small capitals instead of italic.

ERRATA—VOL. II.

- Page 222. Under Thompson, paragraph 3, line 2, "Season late winter" should be "Season early winter."
245. Eighth line from bottom, last word, "sometimes" should be "somewhat."
281. Second paragraph, insert "to 981 inclusive," so that the sentence will read: "The following numbers to 981 inclusive are those given the varieties included in the Department of Agriculture importations of 1870."
282. "Number 21 Veronesh" should be "No. 21 Voronesh."

INDEX TO VARIETIES.

(Accepted names appear wholly or in part in roman type; synonyms in italic.)

PAGE.		PAGE.
178	<i>Abe Lincoln</i>	7
1	ADIRONDACK	7
1	ADMIRABLE	32
2	ALBION	7
3	Albertin	14
3	ALEXANDER	14
3	<i>Alexander the First</i>	14
3	Alexandre	233
110	American	239
62	American Fall	239
241	American Fall Pippin	239
77	American Gloria Mundi	178
83	American Golden	178
83	American Golden Pippin	178
77	American Mammoth	178
143	American Mother	239
211	American Pearmain	178
178	American Red	155
55	American Red Juneating	3
211	American Summer	9
211	American Summer Pearmain	112
5	AMSTERDAM	53
5	Amsterdam Sweet	10
178	Anglesra Pippin	216
5	ANISIM	216
5	Anisim of Peterson	75
6	Antenovka	227
6	Antonowka	10
6	ANTONOVKA	11
6	Antony	12
6	APOINT	12
7	Apoint (synonym of Aport Ori- ent)	10
3	Aporta	
3	Aporta Nalicia	
	APOINT ORIENT	
	Aport Oriental	
	Arabka	
	ARCTIC	
	Arkad Krasieni	
	Arkad Krasieni	
	Arkad Krasieni	
	Aromatic Spike No. 354	
	Astracan Blanch	
	Astracan d'Ete	
	Astracanischer Sommer	
	Astracan Rouge	
	Astrachan	
	Astrachan Red	
	Astrachan Rouge	
	Astrachan White	
	Astrakhan Rouge	
	Astravaskoe	
	Aubertin	
	AUGUST	
	August (synonym of July)	
	August Apple	
	AUGUSTINE	
	August Sweet	
	August Sweeting	
	Aunt Ginnie	
	Aurora	
	Autumnal Bough	
	Autumnal Swaar (synonym of Autumu Swaar)	
	Autumnal Swaar (synonym of Autumn Sweet Swaar)	
	Autumn Sweet	
	Autumnal Sweet Swaar	
	AUTUMN BOUGH	

PAGE.	PAGE.
<i>Autumn Bough</i> 10	<i>Beel Solotofskaja</i> 82
<i>Autumn Bough</i> (synonym of Sweet Bough) 216	<i>Beitigheimer</i> 17
<i>Autumn Pearmain</i> 124	<i>BELBORODOSKOE</i> 15
<i>Autumn Pippin</i> 62	<i>Belle d'Angers</i> 21
<i>Autumn Rose</i> 190	<i>Belle de Boskoop</i> 25
<i>Autumn Seek-No-Further</i> 88	<i>Belle de Boskoop</i> 25
<i>Autumn Strawberry</i> 120	<i>Belle d'Orleans</i> 3
<i>AUTUMN STREAKED</i> 10	<i>Belle Dubois</i> 77
<i>AUTUMN SWAAR</i> II, 12	<i>Belle Josephine</i> (synonym of Gloria Mundi) 77
<i>Autumn Swaar</i> (synonym of Autumn Sweet Swaar) 12	<i>Belle Josephine</i> (synonym of White Spanish Reinette) . . . 241
<i>Autumn Sweet</i> 12	<i>Belle of Boskoop</i> 25
<i>Autumn Sweet Bough</i> 10	<i>Bellerodorskoe</i> 15
<i>AUTUMN SWEET SWAAR</i> 12	<i>Belle Rose</i> 168
<i>Avery Sweet</i> 175	<i>Bellflower of the West</i> 148
<i>Babouskino</i> 83	<i>Bell's Early</i> 203
<i>Babuscheno</i> 83	<i>Bell's Favorite</i> 203
<i>Babuschkino</i> 83	<i>BEN FORD</i> 45
<i>Babushkino</i> 83	<i>Benniger</i> 15
<i>BAILEY SPICE</i> 13	<i>BENNINGER</i> 15
<i>Bailey's Spice</i> 13	<i>Bennington</i> 203
<i>BAKER SWEET</i> 13	<i>BENONT</i> 16
<i>Baker's Sweet</i> 13	<i>Bergamot</i> 6
<i>Baltimore</i> (synonym of Gloria Mundi) 77	<i>Berry Apple</i> 39
<i>Baltimore</i> (synonym of Vande- vere Pippin) 231	<i>Bielborodorska</i> 15
<i>Baltimore Pippin</i> 77	<i>BIETIGHEIMER</i> 17
<i>BANKS</i> 14	<i>Big Rambo</i> (synonym of West- ern Beauty) 230
<i>Banks Gravenstein</i> 14	<i>Big Rambo</i> (synonym of Grosh) . 89
<i>Banks Red Gravenstein</i> 14	<i>Big Sweet</i> 233
<i>Band Apple</i> 49	<i>Big Vandevere</i> 231
<i>Barowski</i> 151	<i>BIRTH</i> 18
<i>Barowski</i> 151	<i>BISMARCK</i> 19
<i>Beard Burden</i> 49	<i>Bismark</i> 19
<i>Beaute de Kent</i> 15	<i>BLACK ANNETTE</i> 20
<i>BEAUTIFUL ARCAD</i> 14	<i>Black Apple</i> 46
<i>Beautiful Arcade</i> 14	<i>Black Detroit</i> 46
<i>Beautiful Pippin</i> 184	<i>Blanche</i> 241
<i>BEAUTY OF KENT</i> 15	<i>Blanche d'Espagne</i> 241
<i>Beauty of Queen</i> 3	<i>Blanche Glacee d'Ete</i> 239
<i>Beauty of the West</i> (synonym of Western Beauty) 239	<i>BLFNHEIM</i> 20
<i>Beauty of the West</i> (synonym of Grosh) 80	<i>Blenheim Orange</i> 21
<i>Beauty Red</i> 40	<i>Blenheim Pippin</i> 21
	<i>Blenheimsrenett</i> 21
	<i>BLOOM</i> 69
	<i>Blooming Orange</i> 21
	<i>Blumen Calvill</i> 85

PAGE.	PAGE.		
BLUSHED CALVILLE	22	<i>Camuscar</i>	241
<i>Blushed June</i>	181	<i>Camuzas</i>	241
<i>Blush June</i>	181	<i>Camuzar</i>	241
<i>Bogdanoff</i> (synonym of Grand- mother)	83	CANADA BALDWIN	69
<i>Bogdanoff Steklanka</i>	83	<i>Canada Peach</i>	159
<i>Bohannon</i>	118	<i>Canada Reinette</i> (synonym of Cheeseboro)	33
BONEM	23	<i>Carolina</i>	240
BOROVINKA	24	<i>Carolina June</i>	181
<i>Borovinka Angluskaja</i>	24	<i>Carolina Red</i>	181
<i>Borovitsky</i> (synonym of Boro- vinka)	24	<i>Carolina Red June</i>	181
<i>Borovitsky</i> (synonym of Olden- burg)	151	<i>Caroline</i>	240
<i>Borowicki</i>	151	CATHEAD	29
<i>Borowitski</i>	151	<i>Cathead</i> (synonym of Cheese- boro)	33
<i>Borowitsky</i>	151	<i>Cathead</i> (synonym of Fall Pip- pin)	62
<i>Borsdorfer</i>	5	<i>Cathead Greening</i>	29
BOSKOOP	25	<i>Catshred</i>	29
<i>Bough</i>	216	<i>Catshead Greening</i>	29
<i>Bough Apple</i>	216	<i>Cattell Apple</i>	148
<i>Bough, Early Sweet</i>	216	<i>Cayuga Red Streak</i>	227
<i>Bough Sweet</i>	216	CELESTIA	29
<i>Bow Apple</i>	216	CHAMPI VIX	30
<i>Bracken</i>	240	CHANDLER	31
<i>Bracy's Seek-No-Further</i>	88	<i>Chandler's Red</i>	31
BRESKOVKA	26	CHARLANOFF	32
BRILLIANT	69	<i>Charlamoski</i>	32
Buckley	33	<i>Charlamoska</i>	32
Buckram	97	<i>Charlamoski</i>	151
Buchanan	149	<i>Charlamoski d'Automne</i>	151
<i>Bullripe</i>	49	<i>Charlamoskircher Nalleoid</i>	151
BUNKER HILL	27	<i>Charlamoska</i>	32
BUTTER	27	<i>Charlamoski</i>	151
<i>Butter Pippin</i>	78	<i>Charlottenthaler</i>	222
<i>Cabane du Chien</i>	98	<i>Charlottenthaler Apple</i>	222
<i>Cabashaw</i>	227	<i>Charlottenthaler Golba</i>	222
CABASHEA	28	<i>Charmant Blanc</i>	213
<i>Cabashie</i>	28	<i>Cheesborough</i>	33
<i>Calkin's Pippin</i>	30	<i>Cheeseboro</i>	33
<i>Calville de Gravenstein</i>	85	<i>Cheeseboro's Russet</i>	33
<i>Calville Grafensteiner</i>	85	<i>Cheesborough</i>	33
<i>Calville Krasmui</i>	22	<i>Cheesborough Russet</i>	33
<i>Cambour des Lorrains</i>	213	CHENANGO	33
<i>Camoisas du roi d'Espagne</i>	241	<i>Chenango Strawberry</i>	33
<i>Camoisee Blanche</i>	241	<i>Chimney Apple</i>	65
<i>Camoisee Blanche</i>	241	<i>Christ Birth</i>	18
		<i>Christ Birth Apple</i>	18

	PAGE.		PAGE.
<i>Christmas</i>	18	<i>COX ORANGE</i>	42
<i>Cinnamon</i>	6	<i>Cox's Orange</i>	42
<i>CLAPPER FLAT</i>	34	<i>Cox's Orange Pippin</i>	42
<i>CLARKE</i>	35	<i>Cox's Pomona</i>	165
<i>Clarke Beauty</i>	35	<i>CRANBERRY PIPPIN</i>	43
<i>CLYDE</i>	36	<i>Crandall Seedling</i>	107
<i>Clyde Beauty</i>	36	<i>CREAM</i>	44
<i>Coalbrook</i>	225	<i>Crimson Beauty</i>	196
<i>Cobbett's Fall</i> (synonym of Fall Pippin)	62	<i>Crimson Pippin</i> (synonym of Scarlet Pippin)	196
<i>Cobbett's Fall</i> (synonym of White Spanish Reinette)	241	<i>Crimson Pippin</i> (synonym of Detroit Red)	46
<i>Cobbett's Fall Pippin</i> (synonym of Fall Pippin)	62	<i>Crimson Scarlet Pippin</i>	196
<i>Cobbett's Fall Pippin</i> (synonym of White Spanish Reinette)	241	<i>CROW EGG</i>	44
<i>COBLING</i>	117	<i>Crow's Egg</i>	44
<i>Codlin, Keswick</i>	117	<i>Cumming's Rambo</i>	89
<i>Coe's Spice</i>	49	<i>Cut Wine</i>	244
<i>Coleman</i>	227	<i>Czarshui Schip</i>	45
<i>Cole's Quince</i>	174	<i>CZAR THORN</i>	45
<i>COLLAMER</i>	36	<i>DEADERICK</i>	45
<i>Collamer Twenty Ounce</i>	36	<i>Dean's Codlin</i>	165
<i>COLTON</i>	37	<i>De Glace D'Ete</i>	239
<i>Colton Early</i>	37	<i>De Glace Hative</i>	239
<i>COLVERT</i>	38	<i>De La Madeleine Rouge</i>	55
<i>Compte Orloff</i>	233	<i>De Lorraine</i>	213
<i>Comte Woronsoff</i>	3	<i>Demary</i>	73
<i>Concombre Ancien</i> (synonym of Fall Pippin)	62	<i>De Moscovie d'Ete</i>	239
<i>Concombre Ancien</i> (synonym of White Spanish Reinette)	241	<i>De Neige</i>	65
<i>Conic June</i>	118	<i>De Notre Dame</i>	213
<i>CONSTANTINE</i>	39	<i>De Rambourg</i>	213
<i>COOPER</i>	40	<i>De Rambure</i>	213
<i>Copp's Mammoth</i>	77	<i>De Ratcau</i>	62
<i>Corail</i>	3	<i>De Ratteau</i>	241
<i>CORNELL</i>	40	<i>De Revel</i> (synonym of Yellow Transparent)	248
<i>Cornell Fancy</i>	41	<i>De Revel</i> (synonym of Vineuse Rouge)	233
<i>Cornell's Favorite</i>	41	<i>Derrick and Ann</i>	186
<i>CORNER</i>	41	<i>Derrick's Graft</i>	186
<i>Cornish Gilliflower</i>	197	<i>De Seigneur d'Automne</i>	29
<i>Corse's St. Lawrence</i>	193	<i>D'Espagne</i> (synonym of White Spanish Reinette)	241
<i>Cos Orange</i>	42	<i>D'Espagne</i> (synonym of Fall Pippin)	62
<i>Costard</i>	29	<i>Deterding's Early</i>	178
<i>Costard Ray</i>	29	<i>Detroit</i>	46
<i>Count Orloff</i>	233	<i>Detroit Black</i>	46
<i>Costard</i>	29		

PAGE.	PAGE.
DETROIT RED 46	Early May 249
D'Ere 141	EARLY PENNOCK 53
De l'in du Conn..... 227	Early Red 141
Diels Sommerkönig 85	Early Red Juneating..... 141
Dix-huit Onces 227	Early Red Margaret..... 141
Dodge's Early Red..... 203	Early Red Pippin..... 93
Douse 95	Early Redstreak..... 93
Dows 95	Early Red Streak..... 93
Dowse 95	EARLY RIPE..... 54
Doedje's Fame 21	EARLY STRAWBERRY..... 54
Duchess 151	Early Striped Juneating..... 141
Duchesse d'Oldenbourg..... 151	Early Summer Pearmain..... 211
Duchess No. 3..... 158	Early Sweet..... 98
Duchess of Oldenburg..... 48, 151	Early Sweet Bough..... 216
Duchess of Oldenburgh..... 151	Early Sweetheart 216
DUDLEY 48	Early Tart Harvest..... 168
Dudley Winter 48	Early Washington 203
Dudley's Winter 48	Egg Jap 44
Du Marechal 65	Egg Top 56
Dumpling 208	Eighteen Ounce 227
Dutchess 151	Eighteen Ounce Apple..... 227
Dutchess of Oldenberg..... 151	ELGIN PIPPIN 56
Dutchess of Oldenburgh..... 151	Elgin Pippin (synonym of White Spanish Reinette) 241
Dutch Mignonne (synonym of Blenheim) 21	Empereur Alexandre I..... 3
Duverson's June 141	Empereur Alexandre de Russie. 3
DYER 49	Empereur de Russie..... 3
Early Baldwin 108	Emperor Alexander 3
Early Bough 216	Englese Orange Appel..... 154
Early Colton 37	Englische Granat-Reinette..... 184
Early Congress 85	Englischer Pepping..... 57
Early French Reinette (synonym of Early Harvest)..... 51	English Borovinka 24
Early French Reinette (synonym of Sweet Bough)..... 216	ENGLISH COBLING..... 117
Early Golden Sweet..... 81	English Jannetling 230
EARLY HARVEST..... 50	English King 3
Early Jack..... 106	English Pearmain 125
Early Jannetling..... 240	EX-LINH PIPPIN 57
EARLY JOE..... 52	English Pippin (synonym of Longfield) 123
Early July Pippin..... 51	English Rambo 89
Early June (synonym of Early Harvest)..... 51	English Sweet (synonym of Ramsdell Sweet) 58, 175
Early June (synonym of Egg Top) 56	English Sweeting 175
Early June of South..... 141	Episcopal 62
Early Margaret 141	Episcopale 241
	Ernst's Apple 140
	Ernst's Pippin 149
	Essex Pippin 184

	PAGE.		PAGE.
<i>Eve</i>	56	<i>Forever Pippin</i>	33
<i>Eve Apple</i> (synonym of <i>Egg Top</i>)	56	<i>Formosa</i>	184
<i>Eve Apple</i> (synonym of <i>Margaret</i>)	141	<i>Formosa Pippin</i>	184
<i>Everbearing</i>	108	<i>Fourth of July</i>	112
<i>Fall Bough</i>	10	<i>Fraise</i>	55
<i>Fall Geneting</i>	59	<i>FRANCHOT</i>	71
<i>Fall Gennetting</i>	59	<i>Frank</i>	33
<i>FALL GREENING</i>	58	<i>Frank Rambour</i>	213
<i>Fall Green Sweet</i>	233	<i>French Pippin</i> (synonym of <i>Holland Pippin</i>)	101
<i>FALL HARVEY</i>	58	<i>French Pippin</i> (synonym of <i>Roman Stem</i>)	188
<i>Fall Jenetting</i>	59	<i>French Rambo</i>	89
<i>FALL JENNETING</i>	59	<i>French Reinette</i>	215
<i>Fall Jenetting</i>	59	<i>French Spitzenburgh</i> (synonym of <i>Summer Spitzenburg</i>)	215
<i>Fall Orange</i>	12, 60	<i>FULLERTON SWEET</i>	71
<i>FALL PIPPIN</i>	61	<i>GARDEN ROYAL</i>	72
<i>Fall Pippin</i> (synonym of <i>Fall Harvey</i>)	58	<i>Gardiner's Apple</i>	143
<i>Fall Pippin</i> (synonym of <i>Holland Pippin</i>)	101	<i>GARDNER SWEET PEARMAN</i>	73
<i>Fall Pippin</i> (synonym of <i>White Spanish Reinette</i>)	241	<i>Gelee d'Ete</i>	239
<i>Fall Queen</i>	91	<i>General Chandler</i>	31
<i>Fall Strawberry</i>	120	<i>GENESEE FLOWER</i>	73
<i>Fall Stripe</i>	106	<i>Geneva Pearmain</i>	30
<i>Fall Swaar</i>	11	<i>Gennetting</i>	240
<i>Fall Swaar of the West</i>	11	<i>German Calville</i>	6
<i>Fall Vanderveer</i>	231	<i>German Calville 324</i>	6
<i>FALL WINE</i>	63	<i>Georgia June</i>	181
<i>FAMEUSE</i>	65	<i>Gibbon's Smathhouse</i>	231
<i>Fameuse</i> (synonym of <i>Utter</i>)	230	<i>Gibbon's Smokehouse</i>	231
<i>Fameuse Baldwin</i>	98	<i>GIDEON</i>	74
<i>FAMEUSE GREEN</i>	69	<i>Gideon White</i>	74
<i>FAMEUSE GROUP</i>	68	<i>Ginetting</i>	240
<i>FAMEUSE NOIRE</i>	69	<i>GINNIE</i>	75
<i>FAMEUSE SUCRE</i>	69	<i>Glacé de Zelande</i>	239
<i>FANNY</i>	69	<i>Glacé d'Ete</i>	239
<i>Fin d'Automne</i>	3	<i>GLADSTONE</i>	75
<i>FISHKILL</i>	70	<i>Glazenwood</i>	77
<i>Fishkill Beauty</i>	70	<i>Glazenwood Gloria Mundi</i>	77
<i>Five-Quartered Gilliflower</i>	197	<i>GLORIA MUNDI</i>	76
<i>Flat</i>	34	<i>Glory of York</i>	184
<i>Flint Russet</i>	170	<i>Gloucester Pippin</i>	21
<i>Flower</i> (of <i>Genesee</i>)	73	<i>Golden Apple</i>	83
<i>Flushing Seek-No-Further</i>	88	<i>GOLDEN PIPPIN (I)</i>	78
<i>FORD</i>	71	<i>GOLDEN PIPPIN (II)</i>	79
		<i>Golden Pippin</i> (synonym of <i>Fall Pippin</i>)	62

PAGE.	PAGE.
Golden Pippin (synonym of Golding) 83	Groskoe Selenka Grüner..... 233
GOLDEN PIPPINS 78	Gros Pomier 91
GOLDEN REINETTE..... 79	Gros Pommier 91
Golden Spice 49	Gros-Rambour d'Ete..... 213
GOLDEN SWEET..... 81	Grosser Mogul..... 87
Golden Sweet (synonym of Northern Sweet) 147	Grosse-Schafnasé 29
Golden Sweeting 81	GRUNDY 90
GOLDEN WHITE..... 82	HAAS 91
GOLDING 82	HAGLOE 92
Goldreinette von Bleuheim..... 21	Hagloe Crab 92
Good Peasant (synonym of Longfield) 123	Hampers American 178
Good Peasant (synonym of Anisim) 5	Harmony 53
Gott. Seward's 127	Harvest (synonym of Early Harvest) 51
Grafen Apfel 85	Harvest (synonym of Primate). 168
Grafensteiner 85	Harvest Apple 215
Granat-Reinette 184	HARVEST REDSTREAK..... 93
Grand Alexander 3	Harvest Red Streak..... 93
Grand Alexandre 3	Harvey 58
Grand Duc Constantin..... 39	HASKELL 93
Grand Duke Constantine..... 39	Haskell Sweet 94
GRANDMOTHER 83	Hass 91
Grand Mother 83	Haverstraw Pippin 30
Grand Sultan (synonym of Vin- euse Rouge) 233	Hawkins Pippin 73
Grand Sultan (synonym of Yel- low Transparent)..... 248	HAWLEY 94
Grantham 108	Hawley (synonym of Hawthorn- den) 96
GRAVENSTEIN 84	Hawthorndean 96
Gravensteiner 85	HAWTHORNDEN 96
Gravenstein Rouge 180	Heickes Summer Queen..... 53
Gravenstine 85	Herbstbreitling 213
Grave Slige..... 85	Herbst Streifling..... 10
Grave Slige..... 85	Herbst Streifling..... 10
Greasy Pippin..... 128	Herr's June..... 141
GREAT MOGUL..... 87	HIBERNAL 96
GREEN SEEK-NO-FURTHER..... 88	HICKS 97
Green Sweet (synonym of Vic- tuals and Drink)..... 233	Highland Pippin..... 168
Green Transparent..... 233	HIGHTOP SWEET..... 97
Gros-Alexandre 3	High Top Sweet..... 98
GROSH 89	High Top Sweet (synonym of Amsterdam) 5
Grosch (synonym of Summer Rambo) 213	High Top Sweeting 98
Grosch's Mammoth 89	HILAIRE 98
	HILTON 99
	Himbeerapfel Lievlander..... 129
	Himbeerapfel Livlander 129
	HOADLEY 99

PAGE.	PAGE.
Hog Island Sweet..... 100	JEFFERSON COUNTY..... 109
Hogpen..... 60	Jeffries..... 108
Holden..... 60	Jellyflower..... 107
Holden Pippin..... 60	Jenneting..... 240
HOLLAND PIPPIN..... 101	Jennetting..... 109
Holland Pippin (synonym of Fall Pippin)..... 62	Jennetting (synonym of White Juneating)..... 240
Hollow Crown Pearmain..... 124	Jennings..... 166
Hominy..... 203	JERSEY SWEET..... 110
Hook..... 103	Jersey Sweeting..... 110
Horning..... 203	Joancting..... 240
Horse..... 91	Joc Precocce..... 52
Hoss..... 91	Jolly Gentleman..... 3
House..... 64	Jonathan of the North..... 5
HOWARD BEST..... 103	Jones Early Harvest..... 181
Howard Russet..... 33	Jones Pippin..... 60
Howard's Best..... 103	Josephine (synonym of Gloria Mundi)..... 77
Howard's Best Russian..... 103	Josephine (synonym of White Spanish Reinette)..... 241
Howe Apple..... 245	JUDSON..... 111
Hower..... 64	JULY..... 112
Hoypen..... 60	July Apple..... 168
Hudson Red Streak..... 125	July Early Pippin..... 51
HUNTER PIPPIN..... 104	July, Fourth of..... 112
Hurlburt..... 105	July Pippin..... 51
HURLBUT..... 104	June (synonym of Margaret) .. 141
Hurlbut (synonym of Ramsdell Sweet)..... 175	June (synonym of Red June) .. 181
Hurlbut Stripe..... 105	Juneating..... 240
Hurlbut Sweet..... 175	Juneating..... 109
Hurlbutt..... 105	Juneating (synonym of White Juneating)..... 240
Imperial Vandervere..... 231	Juneting..... 240
Imperial White..... 157	Junjata..... 234
Imperatrice Eugenie..... 21	
Imperatrice Josephine..... 77	KAIGHN..... 113
Indian Queen..... 53	Kaighn's Spitzemburg..... 113
Indiana Vandervere..... 231	Kaighn's Spitzenberg..... 113
ISHAM..... 106	Kaighn's Spitzenberg..... 113
Isham Sweet..... 106	Kaighn's Spitzenberg (synonym of Kaighn)..... 113
Isle of Wight Orange..... 154	Kaighn's Spitzenberg (synonym of Long Red Pearmain) 125
Isle of Wight Pippin..... 154	Kaighn's Spitzsburgh..... 113
	Kaighn's Spitzsburgh..... 113
JACK..... 106	Kaiser Alexander..... 3
Jack Apple..... 106	Kajabowka..... 114
Jackson..... 33	KALKIDON..... 114
JARVIS..... 107	
Jeffries..... 108	
JEFFERIS..... 108	

PAGE.	PAGE.
Kalkidonskæ 114	Landsburg 119
Kalkidonskæ 114	Langerfeldskæ 123
Kalkidonskæ 114	Lansberger Reinette 119
Kalkil jeltui 247	Large Bough 216
Kalkville scholli 247	Large Early 51
Karaboff 114	Large Early Bough 216
KARABOVKA 114	Large Early Harvest 51
Karabowka 115	Large Early Yellow Bough 216
Kempster's Pippin 21	Large Fall 241
Kennebec Seedling 245	Large Fall Pippin 241
Kent Beauty 15	Large Golden Pippin (synonym of Champlain) 30
Kent Fillbasket 115	Large Golden Pippin (synonym of Golden Pippin I) 78
Kentish Filbasket 115	Large Rambo 89
KENTISH FILLBASKET 115	Large Red and Green Sweeting 177
Kentish Pippin 15	Large Red Sweeting 177
Kentucky Bellflower 125	Large Summer Rambo 89
Kentucky Gilliflower 125	Large Sweet Bough 216
KESWICK 116	Large Vanderveere 231
Keswick Codlin 117	Large White Juneating 51
Keswick Codling 117	Large Yellow Bough 216
Khalkidonskæ 114	Late Bough 10
Kinderhook Pippin 77	Late Chandler 31
Kingsbury Russet (synonym of Cheeseboro) 33	Late Golden Sweet 13
Kingsbury Russet (synonym of Pumpkin Russet) 170	LATE STRAWBERRY 120
KIRKBRIDGE 118	LATHAM 121
Kirkbridge W'hile 118	LA VICTOIRE 69
Knight's Red June 181	LEAD 121
Korallen Apfel 3	Lead Apple 121
	Leeds Beauty 196
La Belle Fameuse 65	Lierland Raspberry 129
Lady Blush 139	Lima 227
Lady de Grey's 115	Lincoln 245
LADY FINGER 118	LINCOLN PIPPIN 122
Lady Finger (synonym of Kaighn) 113	Lincoln Pippin (synonym of Winthrop Greening) 245
Lady Finger (synonym of Long Red Pearmain) 125	Lincolnshire Pippin 96
Lady's Apple 242	LINDENWALD 122
Lady Suffield 126	Lippincott 215
Lady Sutherland 126	Liveland Raspberry 129
Lady Washington 40	Livesley's Imperial 126
La Fameuse 65	Livland Raspberry 129
Lammas 142	Lodge's Early 215
Lancaster Queen 212	LONGFIELD 122
LANDSBERG 119	Longfield's Apple 123
Landsberger Reinette 119	Long Island 60
	Long Island Graft 60

PAGE.	PAGE.
<i>Long Island Pearmain</i> 124	<i>Magdalene</i> 142
<i>Long John</i> (synonym of Kaighn) . . . 113	<i>Magnum Bonum</i> 23
<i>Long John</i> (synonym of Long Red Pearmain) 125	<i>MAGOG</i> 138
<i>Long Pearmain</i> (synonym of Kaighn) 113	<i>Magog Red Streak</i> 138
<i>Long Pearmain</i> (synonym of Long Red Pearmain) 125	<i>MAIDEN BLUSH</i> 139
<i>LONG RED PEARMMAIN</i> 125	<i>MAIDEN FAVORITE</i> 141
<i>LONG STEM</i> 125	<i>Maiden's Apple</i> 141
<i>Long Stem Sweet</i> 13	<i>Maiden's Blush</i> (synonym of Hawthornden) 96
<i>LONGWORTH</i> 125	<i>Malinowska</i> 177
<i>Longworth Red Winter</i> 125	<i>Mammoth</i> (synonym of Gloria Mundi) 77
<i>Lord Kingston</i> 96	<i>Mammoth</i> (synonym of Golden Pippin) 78
<i>Lord Nelson</i> (synonym of Blenheim) 21	<i>Mammoth Pippin</i> 77
<i>LORD SUFFIELD</i> 126	<i>Mammoth Rambo</i> 89
<i>Lothringer Rambour</i> 89	<i>Maralandica</i> 51
<i>Lothringer Rambour d'Ele</i> 213	<i>MARGARET</i> 141
<i>LOU</i> 126	<i>Margaret Early</i> 142
<i>Louis XI'III</i> 55	<i>Margaret, Early Red</i> 142
<i>LOUISE</i> 69, 126	<i>Margaretha Apfel</i> 142
<i>Louise, Princess</i> 127	<i>Marget Apple</i> 142
<i>LOWELL</i> 128	<i>Marguerite</i> 142
<i>Lowell Pippin</i> 128	<i>Marigold</i> 154
<i>LOWLAND RASPBERRY</i> 129	<i>Marigold, Creed's</i> 154
<i>LUBSK QUEEN</i> 129	<i>Marigold Pippin</i> 154
<i>Lubsk Requette</i> 130	<i>Marrow-bone</i> 225
<i>Lucius Apfel</i> 21	<i>Martin</i> 135
<i>Lyman's Large Yellow</i> 171	<i>Marygold</i> 154
<i>Lyman's Pumpkin Sweet</i> 171	<i>Matthews Stripe</i> 131
<i>LYSCOM</i> 130	<i>Maudlin</i> 142
<i>MARIE</i> 131	<i>Meachem Sweet</i> 146
<i>McAdoo's June</i> 112	<i>Melon</i> (synonym of Gloria Mundi) 77
<i>MCCARTY</i> 132, 173	<i>Michigan Beauty</i> 201
<i>McClellan</i> 135	<i>Millcreek</i> 231
<i>MACDONOUGH</i> 132	<i>Millcreek Vandervere</i> 231
<i>McINTOSH</i> 69, 132	<i>MILLER</i> 142
<i>McIntosh Red</i> 133	<i>Miller Seedling</i> 142
<i>Mackie's Clyde Beauty</i> 36	<i>MILLIGEN</i> 142
<i>McLELLAN</i> 134	<i>Mississippi</i> 77
<i>McLelan</i> 135	<i>Mr. Gladstone</i> 75
<i>McMahan</i> 136	<i>Montgomery Sweet</i> 10
<i>McMahan White</i> 136	<i>Monstreuse Pippin</i> 77
<i>McMAHON</i> 136	<i>Monstrous Pippin</i> 77
<i>McMahon White</i> 136	<i>Monstrous Rambo</i> 89
<i>Macomber</i> 187	<i>Montreal</i> 193
	<i>Montreal Peach</i> 159

	PAGE.
<i>Moose</i>	145
<i>Morgan's Favorite</i>	227
<i>MOSHER</i>	143
<i>Mosher Sweet</i>	143
<i>MOTHER</i>	143
<i>Mother Apple</i>	143
<i>Mother of America</i>	143
<i>Mountaineer</i>	145
<i>Mountain Flora</i>	77
<i>MOUNTAIN SWEET</i>	145
<i>MOUSE</i>	145
<i>Mudhole</i>	125
<i>MUNSON</i>	146
<i>Munson Sweet</i>	146
<i>Munson Sweeting</i>	146
<i>Musgrove</i>	89
<i>Musgrove's Cooper</i> (synonym of Grosch)	89
<i>Musgrove's Cooper</i> (synonym of Western Beauty)	239
<i>Mushroom</i>	24
<i>Musk Spice</i>	64
<i>Mygatt's Bergamot</i>	49
<i>Myer's Nonpareil</i>	148
<i>Nalini Jabloky</i>	230
<i>Naylor Rambo</i>	80
<i>Neige</i>	65
<i>Neige-Framboise de Gielen</i>	65
<i>New Brunswick</i>	151
<i>N. J. Red Strick</i>	53
<i>Newtown Greening</i>	83
<i>N. Y. Bellflower</i>	60
<i>N. Y. Gloria Mundi</i>	77
<i>N. Y. Greening</i>	83
<i>Niack Pippin</i> (synonym of Sweet Bough)	217
<i>Nonpareil</i>	148
<i>Nonpareille</i>	184
<i>North American Best</i>	168
<i>Northampton</i>	21
<i>Northern Golden Sweet</i>	147
<i>Northern Golden Sweeting</i>	147
<i>NORTHERN SWEET</i>	147
<i>Northern Sweeting</i> (synonym of Munson)	146
<i>Northern Sweeting</i> (synonym of Northern Sweet)	147

<i>North Star</i>	48
<i>Northwick Pippin</i>	21

The following numbers are those given by Schroeder in his shipment to the Iowa Experiment Station in 1870:

<i>No. 3 M</i>	121
<i>No. 4 M</i>	155
<i>No. 6 M</i>	83
<i>No. 9 M</i>	24
<i>No. 11 M</i>	189
<i>No. 14 M</i>	5
<i>No. 18 M</i>	5
<i>No. 21 M</i>	115
<i>No. 22 M</i>	22
<i>No. 23 M</i>	7
<i>No. 24 M</i>	195
<i>No. 26 M</i>	6
<i>No. 34 M</i>	87
<i>No. 56 M</i>	123
<i>No. 57 M</i>	123
<i>No. 80 M</i>	195
<i>No. 94 M</i>	114
<i>No. 105 M</i>	32
<i>No. 134 M</i>	224
<i>No. 139 M</i>	183
<i>No. 140 M</i>	45
<i>No. 147 M</i>	222
<i>No. 152 M</i>	27
<i>No. 161 M</i>	18

The following numbers are those given to varieties included in the Department of Agriculture importation of 1870:

<i>No. 161</i>	123
<i>No. 205</i>	115
<i>No. 206</i>	45
<i>No. 210</i>	244
<i>No. 224</i>	6
<i>No. 230</i>	224
<i>No. 236</i>	6
<i>No. 245</i>	24
<i>No. 252</i>	7
<i>No. 262</i>	32
<i>No. 288</i>	177
<i>No. 333</i>	182

	PAGE.		PAGE.
No. 340	129	Orange (synonym of Fall Orange)	60
No. 343	182	Orange (synonym of Lowell) ..	128
No. 351	170	Orange (synonym of Orange Pippin)	154
No. 372	195	Orange Blenheim	21
No. 442	247	Orange de Cox	42
No. 444	130	ORANGE (OF NEW JERSEY)	153
No. 453	14	ORANGE (OF PENNSYLVANIA) ..	153
No. 457	39	ORANGE PIPPIN	154
No. 460	83	Orange Pippin (synonym of Blenheim)	21
No. 472	155	Orange Russet	154
No. 477	18	ORANGE SWEET	154
No. 530	114	Orange Sweet (synonym of Golden Sweet)	81
No. 537	57	Orange Sweet (synonym of Munson)	146
No. 599	189	Orange Sweeting (synonym of Golden Sweet)	81
No. 964	10	Orange Sweeting (synonym of Orange Sweet)	154
No. 978	82	Orloff	233
No. 979	82	Orlovskoe	233
No. 981	82	Orlovskoe	233
No. 12 Orel	7	Orsimui	97
No. 21 Veronesh	247	Osgood's Favorite	131
No. 51 Vor.	79	Osimoe	97
No. 81 Vor.	83	Oskaloosa	106
Nyack	30	OSTRAKOFF	155
Nyack Pippin	30	Ostrakoff Glass	155
Oats	51	Ostrekhoff	155
OGDENSBURG ..	147	Ostrekhoff's Glass	155
Ohio Beauty (synonym of Western Beauty)	230	Ostrekovskaya Steklianka	155
Ohio Beauty (synonym of Grosh) ..	89	Ostrekovskaya Steklianka	155
OHIO NONPAREIL	148	Ostrokoff	155
Ohio Nonpareil (synonym of Gravenstein)	85	Ostrokoff's Glass	155
Ohio Nonpareil	148	Ox Apple	77
OHIO PIPPIN	149	Oxeye	230
Ohio Wine (synonym of Fall Wine)	64	Oxheart	33
Ohio Wine (synonym of Kaighn)	113	Owen's Golden Beauty	240
OKABENA	150	Ozark Pippin	45
Okobena	150	PALOUSE	156
OLDENBURG	150	Paper	30
Oldenburg, Duchess of	151	Paper-skin	30
Oldenburg	151	Paradies Apfel	85
Old Hawthorndean	96	Park	125
Omensk	189		
Oporto	7		
ORANGE	153		

PAGE,	PAGE,
PARRY WHITE..... 157	Plum Cider 164
PATTEN 158	Pointed Pipka 32
Patten's Duchess No. 3..... 158	Polecat 212
Patten Greening 158	Pomme Astrachan..... 239
PEACH (Montreal)..... 159	Pomme d'Astrachan 239
Peach Apple of Montreal..... 159	Pomme de Blenheim..... 21
Peach of Montreal..... 159	Pomme de Fameuse..... 65
PEACH POND..... 160	Pomme de Hawthornden..... 96
Peach Pond Sweet..... 160	Pomme de Neige..... 65
Peach Pound Sweet..... 160	Pomme de Notre-Dame..... 213
Pear Apple 174	Pomme d'Ete (of Canada)..... 51
Pear Lot 244	Pomme d'Orange..... 154
Pearmain 125	Pomme Fameuse 65
PEARSALL 161	Pomme Graefenstein..... 85
Pearsall's Sweet 161	Pomme Peche 150
Pear Tree Lot..... 244	Pomme Royal..... 49
PEASE 161	Pomme Royale..... 49
Pease, Walter 161	Pomme Roze..... 49
PEASGOOD NONSUCH 162	Pomme Water..... 49
Peasgood's Nonsuch 162	Pommewater (in Illinois)..... 49
Pennsylvania Vandevere..... 231	POMONA 164
Pepin de Kent 15	Pomona Britannica..... 3
Pepin Ribston 184	Pompey 233
Pipping Englishcher 57	PORTER 166
Perte d'Angleterre 21	Possaris Nalivia..... 6
PERRY REDSTREAK..... 162	Potter's Large..... 116
Perry Red Streak..... 162	Potter's Large Grey Seedling.. 116
PETER 163	Potter's Large Seedling..... 116
Peterson's Charlamoff 32	Pound 77
Petrovskoe 195	Pound Pippin..... 62
Petrowskoe 195	Pound Royal (synonym of Fall Pippin) 62
Philadelphia Pippin (synonym of Fall Pippin)..... 62	Pound Royal (synonym of Golden Pippin I.)..... 78
Philadelphia Pippin (synonym of White Spanish Reinette)..... 241	Pound Royal (synonym of Lowell) 128
Philadelphia Sweet 10	Pound Royal (synonym of Long Island Pearmain)..... 125
Phoenix 3	Pound's July..... 217
Phonix 3	POUND SWEET..... 167
Pickaway Rambo 89	Pound Sweet (synonym of Pumpkin Sweet)..... 171
Pie Apple (synonym of Holland Pippin) 101	Powers..... 168
Pie Apple (synonym of Sops of Wine) 203	President Napoleon..... 3
Pippin Kent 15	Pride of Genesee..... 73
Pittstown Pippin 79	PRIMATE 167
Plodowitka Cudakaja 170	Primiting 240
Plodowitka Cudakaja 170	Prince Bismark 19
PLUMB CIDER..... 164	

	PAGE.		PAGE.
<i>Prince of Wales</i>	21	<i>Ramsdell</i>	175
<i>Prince's Early Harvest</i>	51	<i>Ramsdell Red Sweet</i>	175
<i>Prince's Harvest</i>	51	<i>Ramsdell's Red</i>	175
<i>Prince's Large Pippin of N. Y.</i> ..	62	<i>Ramsdell's Red Pumpkin Sweet</i> ..	175
<i>Prince's Large Red and Green</i> <i>Sweeting</i>	177	<i>Ramsdell's Red Sweeting</i>	175
<i>Prince's Red and Green Sweet</i> ..	177	RAMSDELL SWEET.....	175
<i>Prince's Yellow Harvest</i>	51	<i>Ramsdell's Sweeting</i>	175
<i>Princess Louise</i>	127	<i>Randall's Red Winter</i>	175
<i>Princessin Apfel</i>	85	RASPBERRY	176
<i>Prolific Sweet</i>	170	<i>Ray Apple</i>	146
PROLIFIC SWEETING	169	RED AND GREEN SWEET.....	177
<i>Prussian</i>	38	<i>Red and Green Sweeting</i>	177
PUMPKIN RUSSSET	170	<i>Red Astracan</i>	178
PUMPKIN SWEET	171	RED ASTRACHAN.....	178
<i>Pumpkin Sweet</i> (synonym of <i>Cheeseboro</i>)	33	<i>Red Astrakhan</i>	178
<i>Pumpkin Sweet</i> (synonym of <i>Pumpkin Russet</i>).....	170	<i>Red Beitigheimer</i>	17
<i>Pumpkin Sweeting</i> (synonym of <i>Pumpkin Sweet</i>).....	172	<i>Red Bellflower</i> (synonym of Ohio <i>Nonpareil</i>)	148
<i>Pumpkin Sweeting</i> (synonym of <i>Pumpkin Russet</i>).....	170	<i>Red Bellflower</i> (synonym of <i>Long Red Pearmain</i>).....	125
<i>Pyrus Astracanicus</i>	239	<i>Red Bietigheimer</i>	19
<i>Queen</i> (synonym of <i>Summer</i> <i>Queen</i>)	212	<i>Red Bough</i>	177
<i>Queen</i> (synonym of <i>Williams</i>)..	242	<i>Red Cheek</i> (synonym of Rasp- <i>berry</i>)	177
<i>Queen Anne</i> (synonym of <i>Low-</i> <i>ell</i>)	128	<i>Red Cheek</i> (synonym of <i>Maiden</i> <i>Blush</i>)	130
<i>Queen Anne</i> (synonym of <i>Mother</i>)	143	<i>Red Cheek</i> (synonym of <i>Fall</i> <i>Orange</i>)	60
QUINCE (of <i>Cole</i>).....	174	<i>Red Gilliflower</i> (synonym of <i>Scollop Gilliflower</i>).....	197
QUINCE (of <i>Coxe</i>).....	174	<i>Red Gilliflower</i> (synonym of <i>Striped Gilliflower</i>)	207
<i>Quince Apple</i>	174	RED GRAVENSTEIN.....	180
<i>Rag Apple</i>	146	<i>Red Gravenstein</i> (synonym of <i>Banks</i>)	14
<i>Rambour</i>	213	<i>Red Hawthornden</i>	66
<i>Rambour Aigre</i>	213	RED HOOK.....	180
<i>Rambour Blanc</i>	213	<i>Red Joaneting</i>	142
<i>Rambour d'Amerique</i>	213	RED JUNE.....	181
<i>Rambour d'Ete</i>	213	<i>Red Junceating</i> (synonym of <i>Margaret</i>)	142
<i>Rambour Franc</i>	213	<i>Red Juneating</i> (synonym of <i>Red</i> <i>June</i>)	181
<i>Rambour Gros</i>	213	<i>Red Juneating</i> (synonym of <i>Early Strawberry</i>).....	55
<i>Rambour Lorraine</i>	89	<i>Red June, Carolina</i>	181
<i>Rambour Raze</i>	213	<i>Red June of South</i>	142
<i>Rambu</i>	213	<i>Red Juneting</i>	142
<i>Ramsdale's Sweeting</i>	175		

PAGE.	PAGE.
<i>Red Pearmain</i> (synonym of Kaighn)..... 113	<i>Reinette de Traver</i> 184
<i>Red Pearmain</i> (synonym of Long Red Pearmain)..... 125	<i>Reinette Grenade Anglaise</i> 184
<i>Red Phoenix</i> (synonym of Kaighn)..... 113	<i>Reinette Landsberger</i> 119
<i>Red Phoenix</i> (synonym of Long Red Pearmain)..... 125	<i>Reinette Liubski</i> 130
<i>Red Pippin</i> (synonym of Kaighn)..... 113	<i>Reinette Monstrueuse</i> 25
<i>Red Pippin</i> (synonym of Long Red Pearmain)..... 125	<i>Reinette Orange de Cox</i> 42
<i>Red Pumpkin Sweet</i> 175	<i>Reinette Quarrendon</i> 142
<i>Red Pumpkin Sweeting</i> 175	<i>Reinette Tendre</i> 241
<i>Red Reinette</i> 83	<i>Reinette von Montfort</i> 25
<i>Red Shropsarine</i> 203	<i>Renet Liubskui</i> 130
<i>Red Spitzenberg</i> (synonym of Kaighn)..... 113	REPKA..... 183
<i>Red Spitzenberg</i> (synonym of Long Red Pearmain)..... 125	<i>Remboure d'Ele</i> 213
<i>Red Spitzenberg</i> (synonym of Richard Graft)..... 186	<i>Roschestwenskoe</i> 19
<i>Red Spitzenburg</i> 113	<i>Revelstone</i> 248
<i>RED TRANSPARENT</i> 182	<i>Revelstone Pippin</i> 233
<i>Red Transparent</i> (synonym of Vineuse Rouge)..... 233	<i>Rhode Island Sweet</i> 172
<i>RED TYPE OF TWENTY OUNCE</i> ... 229	<i>Riabnouka</i> 39
<i>Red Vanderveere</i> 231	<i>Ribbed Gilliflower</i> 197
<i>RED WINE</i> 182	<i>Ribbed Pippin</i> 83
<i>Red Winter Pearmain</i> (synonym of Kaighn)..... 113	RIESTON..... 184
<i>Red Winter Pearmain</i> (synonym of Long Red Pearmain)..... 125	<i>Ribstone</i> 184
REED..... 183	<i>Ribstone Pippin</i> 184
<i>Reindell's Large</i> 175	<i>Ribstone Pepping</i> 184
<i>Reinette A Gobelet</i> 241	<i>Ribston Pippin</i> 184
<i>Reinette Belle de Boskoop</i> 25	<i>Richard</i> 186
<i>Reinette Blanche</i> 241	RICHARD GRAFT..... 186
<i>Reinette Blanche d'Espagne</i> (synonym of White Spanish Reinette)..... 241	<i>Ridge</i> (synonym of Ribston).... 184
<i>Reinette Blanche d'Espagne</i> (synonym of Fall Pippin).... 62	<i>Riefka</i> 183
<i>Reinette de Blenheim</i> 21	<i>Riff Apfel</i> 85
<i>Reinette d'Espagne</i> 241	<i>Rockhill's Russet</i> 184
<i>Reinette d'Holland</i> 101	<i>Rode Wyn Appel</i> 203
<i>Reinette d'Hollande</i> 101	ROLFE..... 187
<i>Reinette de Landsberg</i> 119	ROMAN STEM..... 188
	<i>Romenskoe</i> 189
	ROMNA..... 189
	<i>Romna</i> (synonym of Hiberna).... 97
	<i>Romnenskoe</i> 189
	RONK..... 190
	<i>Roschdestwenskoe</i> 19
	<i>Roschdestwenskoe</i> 19
	ROSE RED..... 190
	<i>Roshdestwenskoe</i> 19
	<i>Rother Astrakhan</i> 178
	<i>Rother Gravensteiner</i> 180
	<i>Rother Jacobs</i> 142
	<i>Rother Jacobs Apfel</i> 142
	<i>Rother Weinapfel</i> 182
	<i>Rotherwein Appel</i> 182

PAGE.	PAGE.
<i>Rough and Ready</i> 168	SCOLLOP GILLIFLOWER 197
<i>Round Catshead</i> 29	<i>Scollop Gilliflower</i> (synonym of
<i>Round Sweet</i> 172	<i>Striped Gilliflower</i>) 207
<i>Round Top</i> 56	<i>Scolloped Gilliflower</i> 197
<i>Rubets Vinogradnui</i> 244	<i>Scott</i> 168
<i>Rubez Vinogradnui</i> 244	SCOTT BEST 198
<i>Russam</i> (synonym of Kaighn) .. 113	<i>Seek-No-Further</i> (synonym of
<i>Russam</i> (synonym of Long Red	<i>Cooper</i>) 40
<i>Pearmain</i>) 125	<i>Seek-No-Further</i> (synonym of
<i>Russian Emperor</i> 3	<i>Green Seek-no-Further</i>)..... 88
<i>Russian Gravenstein</i> 6	<i>Seever's Red Streak</i> 230
<i>Rusty Core</i> 148	SENECA FAVORITE 198
	<i>Seneca Favorite</i> (synonym of
<i>Sabine</i> 85	<i>Quince</i> [of Cox]) 174
<i>Sabine of the Flemmings</i> 85	<i>Seneca Spice</i> 174
<i>Safstaholm</i> 191	<i>Shaker's Yellow</i> 53
SÄFSTAHDOLMS 191	<i>Shannon</i> 149
<i>Säfstaholmsäpple</i> 191	SHARP 190
<i>Säfstaholmsäpple</i> 191	<i>Sharpe's Early</i> 212
<i>Sailee Russet</i> 192	<i>Sharpe's Spice</i> 64
<i>Saille Sweet</i> 177	<i>Sheepnose</i> (synonym of <i>Egg</i>
<i>Sailly Autumn</i> 192	<i>Top</i>) 56
<i>Saint-Germain</i> 241	<i>Sheepnose</i> (synonym of Long
<i>St. Hilaire</i> 98	<i>Red Pearmain</i>) 125
<i>St. John Strawberry</i> 55	SHERMAN 200
<i>St. Lawrence</i> 192	<i>Sherman's Favorite</i> 200
<i>Saint-Lawrence</i> 193	<i>Sherman's Sweet</i> 200
<i>Saint Laurent</i> 193	<i>Sherwood's Favorite</i> 33
<i>St. Peter</i> 194	<i>Shiowasse Beauty</i> 201
<i>Sandy Glass</i> 195	SHIAWASSEE 69, 201
<i>Sanguineus</i> 65	<i>Shiawassee Beauty</i> 201
<i>Sapson</i> 203	<i>Shiawassie Beauty</i> 201
<i>Sapsoncine</i> 203	<i>Shoreditch White</i> 96
<i>Sassafras Sweet</i> 94	<i>Shropshirecine</i> 203
SAXTON 196	<i>Siberian August</i> 112
<i>Scalloped Gilliflower</i> 197	SINE-QUA-NON 202
<i>Scalloped Gillyflower</i> 197	<i>Skvosnoi krasnoi</i> 182
<i>Scarlet Pearmain</i> (synonym of	<i>Skvosnoi Krasnoi</i> 182
<i>Kaighn</i>) 113	<i>Skvosnoi Schotoi</i> 248
<i>Scarlet Pearmain</i> (synonym of	<i>Sleeper's Yellow</i> 53
Long Red Pearmain) 125	SLINGERLAND 202
SCARLET PIPPIN 196	<i>Slingerland's Fall Pippin</i> 202
<i>Schafnasé</i> 29	<i>Slingerland Pippin</i> 202
<i>Scharlottenthaler Golba</i> 22	<i>Small Admirable</i> 1
<i>Schoone van Baskoop</i> 25	<i>Small's Admirable</i> 1
<i>Schoone van Baskoop</i> 25	<i>Smithfield Spice</i> 49
SCHUYLER SWEET 197	<i>Smokehouse</i> (synonym of Van-
	<i>devere Pippin</i>) 231

PAGE.	PAGE.
<i>Smryna</i> 33	<i>Striped V'ondereve</i> 231
<i>Snow</i> 65	STROAT 208
<i>Solotoc Renet</i> 79	<i>Strode</i> 208
SOMERSET (N. Y.) 203	STRODE BIRMINGHAM 208
<i>Sops in Wine</i> 203	<i>Strode's</i> 208
SOPS OF WINE 203	<i>Strode's Birmingham</i> 208
<i>Sourbough</i> (synonym of Cham- plain) 30	<i>Strohmer</i> 85
SOUR BOUGH 204	<i>Stromling</i> 85
<i>Sour Bough</i> (synonym of Cham- plain) 30	STUMP 209
<i>Sour Bough</i> (synonym of Tart Bough) 220	<i>Stymer's</i> 210
<i>Sour Harvest</i> 168	STYMUS 210
<i>Speckled</i> 60	SUFFOLK BEAUTY 210
SPECTATOR 204	<i>Suisleffer</i> 219
<i>Spiced Oaxye</i> 231	<i>Summer Bellefleur</i> 211
<i>Staalclubs</i> 231	SUMMER BELLFLOWER 211
<i>Staalclubs</i> 230	<i>Summer Bellflower</i> (synonym of Autumn Bough) 10
<i>Stalcubs</i> 230	<i>Summer Hugloc</i> 92
STARKEY 205	SUMMER PEARMAIN 211
STARR 206	<i>Summer Pippin</i> (synonym of Champlain) 30
<i>Steklianka pesotchnaya</i> 195	<i>Summer Pippin</i> (synonym of Fall Pippin) 62
<i>Stewart's Nonpareil</i> 112	<i>Summer Pippin</i> (synonym of Holland Pippin) 101
STILLMAN 207	SUMMER QUEEN 212
STILLMAN EARLY 207	SUMMER RAMBO 213
<i>Stillman's Early</i> 207	<i>Summer Rambo</i> (synonym of Grosh) 89
<i>Stoke Tulip</i> 3	<i>Summer Rambour</i> 213
Straat 208	SUMMER REDSTREAK 214
<i>Strawberry</i> (synonym of Sops of Wine) 203	SUMMER ROSE 215
<i>Strawberry</i> (synonym of Che- nango) 33	<i>Summer Russet</i> 218
<i>Strawberry</i> (synonym of Late Strawberry) 120	SUMMER SPITZENBURG 215
<i>Strawberry</i> (synonym of Richard Graft) 186	<i>Summer Sweet</i> 216
<i>Strawberry</i> 207	<i>Summer Sweet</i> (synonym of Hightop Sweet) 98
<i>Striped Ashmore</i> 231	<i>Summer Sweeting</i> (synonym of Summer Sweet) 216
<i>Striped Bellflower</i> 207	<i>Summer Sweeting</i> (synonym of Hightop Sweet) 98
STRIPED FAMEUSE 68	<i>Summer Traveller</i> 142
STRIPED GILLFLOWER 207	<i>Superior White</i> 157
<i>Striped Harvest</i> 93	<i>Susy Clark</i> 181
<i>Striped Juncating</i> 112	<i>Svinsovka</i> 121
<i>Striped Pearmain</i> 125	<i>Swedish Borsdorf of Patten</i> 5
<i>Striped Quarrendon</i> 142	SWEET BOUGH 216
STRIPED RED JUNE 182	
<i>Striped Shropshire</i> 55	

	PAGE.		PAGE.
SWEET FALL PIPPIN	218	Tinmouth Sweet	223
Sweet Golden Pippin	12	Titovca	224
Sweet Harvest	217	TITOVKA	224
Sweet June	98	Titovka	224
Sweet Lyman's Pumpkin	172	Titus Apple	224
Sweet Pippin	100	Titus Riga	224
Sweet Rambo	89	Tom Harryman	85
SWEET RUSSET	218	Tompkins	49
Sweet Russet (synonym of		Tom Put	225
Pumpkin Russet)	170	TOM PUTT	225
Sweet Russet (synonym of		Transparente de Astracan.....	239
Cheesboro)	33	Transparente d'Ete	239
Sweet's Harvest	212	Transparent de Muscovie	239
Sweet Suvar	12	Transparente de Muscovie d'Ete.	239
Sweet Wine	64	Transparente de Saint-Leger	
Sweet's Harvest	212	(synonym of Vineuse Rouge).	233
Swinez	121	Transparente de Saint-Leger	
Swincowska	121	(synonym of Yellow Trans-	
SWITZER	218	parent)	248
SYLVESTER	220	Transparente de Zurich	239
		Transparente Jaune (synonym of	
Taffitai	239	Vineuse Rouge)	233
Tallow	128	Transparente jaune (synonym of	
Tallow Apple	128	Yellow Transparent)	248
Tallow Pippin	128	Transparente Rouge	233
Tars Thorn	45	Transparente Verte	233
TART BOUGH	220	Transparent Muscovie	239
Tart Bough (synonym of Early		Travers	184
Harvest)	51	Travers Apple	184
Tart Bough (synonym of Cham-		Travers Peppin	184
plain)	30	Travers Pippin	184
Teignmouth	223	Travers Reinette	184
Tennessee Early Red	55	Trenton Early	81
Tete d'Ange	29	Tsarskui Schip	45
Tete de Chat	29	TUPIS	226
Tetofsky	220	Tufts Baldwin	226
Tetofski (synonym of July)....	112	Tufts Seedling	226
Tetofski (synonym of Tetofsky).	220	Turn Off Lane	174
TETOFSKY	220	TWENTY OUNCE	227
Tetofsky (synonym of July)....	112	Twenty Ounce Pippin	227
THALER	222	TYRE BEAUTY	229
THOMPSON	222		
Thompson Seedling No. 24.....	222	Uncle Sam's Best	64
Thompson No. 24	222	Underdunk	30
Thompson's Seedling No. 29....	111	UTTER	229
Thompson's Seedling No. 38....	90	Utter's Large Red	230
TINMOUTH	223	Utter's Red	230

PAGE.	PAGE.
<i>Vandervee</i> (synonym of Vandevere) 230	<i>Watkins Early</i> 211
<i>Vandervee</i> (synonym of Vandevere Pippin) 231	<i>Watson's Vandevere</i> 231
<i>Vandervee Pippin</i> 231	<i>Watson's Vandevere</i> 231
<i>Vandervee</i> 230	<i>WEALTHY</i> 236
<i>VANDEVERE</i> 230	<i>Weinappel Rother</i> 182
<i>VANDEVERE PIPPIN</i> 231	<i>Weinappel Rother</i> 182
<i>Vandevere Yellow</i> 231	<i>Weiser Hawthornden</i> 96
<i>Van Dym's Pippin</i> 62	<i>Wells</i> 148
<i>Van Dync Apple</i> 77	<i>Westbrook</i> 60
<i>Van Dyn's Pippin</i> 62	<i>WESTERN BEAUTY</i> 239
<i>Van Kleek's Sweet</i> 100	<i>Western Beauty</i> (synonym of Grosh) 89
<i>Vargul</i> 6	<i>Western Beauty</i> (synonym of Ohio Nonpareil) 148
<i>Vermillon d'Ete</i> 178	<i>Wheeler's Kernel</i> 96
<i>Vermont</i> 30	<i>White Apple</i> (synonym of Hawthornden) 96
<i>Vermont Pippin</i> 223	<i>White Apple</i> (synonym of Parry White) 157
<i>Vermont Pumpkin Sweet</i> 172	<i>White Astracan</i> 239
<i>Vermont Sweet</i> 172	<i>WHITE ASTRACHAN</i> 239
<i>VICTORIA</i> 231	<i>White Borodovka</i> 15
<i>Victoria Sweet</i> 231	<i>White Graft of Wisconsin</i> 60
<i>Victoria Sweeting</i> 231	<i>White Hawthornden</i> 96
<i>VICTUALS AND DRINK</i> 233	<i>White Hawthornden</i> 96
<i>Vilkuil Mogul</i> 87	<i>White June</i> 118
<i>VINEUSE ROUGE</i> 233	<i>WHITE JUNEATING</i> 240
<i>Vinnoc Krasnoe Osennee</i> 182	<i>White Newell</i> 60
<i>Vinograd</i> 244	<i>White Seck-No-Further</i> 88
<i>Virginia June</i> 142	<i>WHITE SPANISH REINETTE</i> 241
<i>Virginia Sweet</i> 177	<i>White Spice</i> 49
<i>Voronesh No. 21</i> 247	<i>WILLIAMS</i> 242
<i>Wabash Bellflower</i> 125	<i>Williams Early</i> 242
<i>Walter Pease</i> 161	<i>Williams Early Red</i> 242
<i>Watworth</i> 30	<i>Williams Favorite</i> 242
<i>Warden's Pie Apple</i> 203	<i>Williams Favorite Red</i> 242
<i>Ward's Pippin</i> 21	<i>Williams Red</i> 242
<i>Warren Pennoek</i> 53	<i>Willis Sweet</i> 244
<i>Washington</i> (synonym of Sops of Wine) 203	<i>Willis Sweeting</i> 244
<i>Washington</i> (synonym of Washington Strawberry) 234	<i>Wilson's June</i> 181
<i>Washington</i> (synonym of Sweet Bough) 217	<i>Windover</i> 231
<i>Washington Co. Seedling</i> 234	<i>Wine</i> (synonym of Egg Top) 56
<i>Washington of Maine</i> 234	<i>Wine</i> (synonym of Fall Wine) 64
<i>WASHINGTON STRAWBERRY</i> 234	<i>Wine</i> (synonym of Richard Graft) 186
<i>WATER</i> 236	<i>Wine</i> (synonym of Twenty Ounce) 227
<i>Waterloo</i> 178	<i>Wine of Cole</i> 64

PAGE.	PAGE.		
WINE RUBETS	244	<i>Yellow June</i> (synonym of Kirk- bridge)	118
<i>Winter Chandler</i>	31	<i>Yellow Juneating</i>	51
<i>Winter Golden Sweet</i>	13	<i>Yellow May</i>	240
<i>Winter Pearmain</i> (synonym of Long Island Pearmain)	124	<i>Yellow Summer Pearmain</i>	166
<i>Winter Pearmain</i> (synonym of Long Red Pearmain)	125	<i>Yellow Sweeting</i>	81
<i>Winter Seek-No-Further</i>	88	YELLOW TRANSPARENT	247
WINTHROP GREENING	245	<i>Yellow Transparent</i> (synonym of Thaler)	222
WOLF RIVER	245	<i>Yellow Vanderveve</i>	231
<i>Wolf River</i> (synonym of Alex- der)	3	YOPP	249
<i>Wolman's Harvest</i>	215	<i>Yopp's Favorite</i>	249
<i>Woodstock</i> (synonym of Blen- heim)	21	YORK	249
<i>Woodstock</i> (synonym of Dyer)	49	<i>York and Lancaster</i>	193
<i>Woodstock Pippin</i>	21	<i>York Pippin</i> (synonym of Fall Pippin)	62
<i>Woolman's Early</i>	215	<i>York Pippin</i> (synonym of Golden Pippin 1)	78
<i>Woolman's Harvest</i>	215	<i>York Pippin</i> (synonym of White Spanish Reinette)	241
<i>Woolman's Striped Harvest</i>	215	<i>York Russet</i> (synonym of Cheeseboro)	33
<i>Woolverton</i>	127	<i>York Russet</i> (synonym of Pumpkin Russet)	170
WORKAROE	246	<i>York Russeting</i>	33
<i>Wunderapfel</i>	3	<i>Zarskischip</i>	45
<i>Yanker Apple</i>	172	<i>Zarski Schip</i>	45
<i>Yellow Bough</i>	217	<i>Zarski Zars</i>	45
YELLOW CALVILLE	247	<i>Zolotwi Renet</i>	79
<i>Yellow Flat</i>	118	<i>Zour Bough</i>	168
<i>Yellow Harvest</i>	51	<i>Zuzoff of Tuttle</i>	5
<i>Yellow June</i> (synonym of White Juneating)	240		

CRABAPPLES.

<i>Anther Crab</i>	269	CORAL	252
<i>Baccata fructa oblonga</i>	261	CURRANT	253
BAILEY CRIMSON	251	<i>Current Crab</i>	253
<i>Bailey's Crimson</i>	251		
<i>Briar Sweet</i>	251	DAKTMOUTH	253
BRIER	251		
<i>Brier's Sweet</i>	251	EXCELSIOR	254
<i>Brier Sweet</i> (synonym of Brier)	251		
<i>Brier Sweet</i> (synonym of Van Wyck)	268	FLORENCE	255
CHERRY	252	GIBB	256
		<i>Golden Beauty</i>	269

PAGE.	PAGE.
<i>Hislop</i> 257	QUAKER 263
HYSLOP 257	QUEEN CHOICE 264
LARGE RED SIBERIAN..... 258	<i>Queen's Choice</i> 264
LARGE YELLOW SIBERIAN..... 258	RED SIBERIAN 264
MARENGO 259	SEPTEMBER 264
<i>Marengo No. 1</i> 259	<i>Siberian Crab</i> (synonym of Red
<i>Marengo Winter Crab</i> 259	Siberian) 264
MARTHA 259	<i>Siberian Crab</i> (synonym of Yel-
MINNESOTA 260	low Siberian) 269
<i>Montreal</i> 261	SOULARD 265
MONTREAL BEAUTY 261	TRANSCENDENT 266
OBLONG 261	<i>Transcendant</i> 267
<i>Oblong-Fruited Siberian Crab.</i> 261	VAN WYCK 267
<i>Oblong Siberian Crab</i> 261	<i>Van Wyck</i> (synonym of Brier). 251
ORANGE 261	<i>Van Wyck Siberian</i> 268
	<i>Van Wyck Sweet</i> 268
PAUL IMPERIAL 262	WHITNEY 268
PICTA STRIATA 263	<i>Whitney No. 20</i> 268
<i>Pieta</i> 263	YELLOW SIBERIAN 269
<i>Pieta Striata</i> 263	
<i>Pomme Groseille</i> 253	

COMBINED INDEX TO VARIETIES.

Volumes 1 and 2.

(Accepted names appear wholly or in part in roman type; synonyms in italic.)

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Abe Lincoln</i>		178
<i>Aberdeen</i>	227	
<i>Accidental</i> ..	227	
<i>Acuba-leaf Reinette</i>	52	
<i>Acuba-leaved Reinette</i>	52	
ADIRONDACK		I
ADIRONDACK		I
<i>Æsopus Spitzenberg</i>	120	
<i>Æsopus Spitzenberg</i>	120	
<i>Æsopus Spitzenburg</i>	120	
<i>Æsopus Spitzsburgh</i>	120	
<i>Aiken</i>	41	
<i>Aikens Winter</i> (of Downing)	41	
<i>Aikin's Red</i>	41	
<i>Aiken</i>	41	
AKIN	41	
<i>Akin Red</i>	41	
<i>Akin Seedling</i>	41	
<i>Akin's Red</i>	41	
<i>Albemarle</i>	146	
<i>Albemarle Pippin</i>	18, 146	
<i>Albertin</i>		3
ALBION		2
ALEXANDER	17, 18, 20, 25	3
ALEXANDER GROUP	25	
<i>Alexander the First</i>		3
<i>Alexandre</i>		3
<i>Alleghany</i>	227	
ALLINGTON	42	
<i>Allington Pippin</i>	42	
ALLISON	43	
<i>Amindelig</i>	181	
AMASSIA	44	
<i>Amber Crab</i>		269

	VOL. I. PAGE.	VOL. II. PAGE.
<i>American</i>		110
<i>American Beauty</i> (synonym of <i>Sterling</i>).....	45, 319	
<i>American Beauty</i> (synonym of <i>Sheriff</i>).....	308	
AMERICAN BLUSH	45	
<i>American Blush</i> (synonym of <i>Hubbardston</i>).....	161	
<i>American Fall</i>		62
<i>American Fall Pippin</i>		241
<i>American Gloria Mundi</i>		77
<i>American Golden</i>		83
<i>American Golden Pippin</i>		83
<i>American Golden Russet</i>	89	
<i>American Golden Russet</i> (of <i>New England</i>).....	164	
<i>American Mammoth</i>		77
<i>American Mother</i>		143
<i>American Newtown Pippin</i>	146	
<i>American Nonpareille</i>	161	
<i>American Pearmain</i>		211
AMERICAN PIPPIN	45	
<i>American Red</i>		178
<i>American Red Juneating</i>		55
<i>American Seek-No-Further</i>	273	
<i>American Summer</i>		211
<i>American Summer Pearmain</i>		211
AMOS	46	
<i>Amos Jackson</i>	46	
AMSTERDAM		5
<i>Amsterdam Sweet</i>		5
ANDREWS	46	
<i>Andrews Winter</i>	46	
<i>Anglesea Pippin</i>		178
ANIS	26	
ANISIM		5
<i>Anisim</i> (of <i>Peterson</i>)		5
<i>Antonovka</i>		6
<i>Antonowka</i>		6
ANTONOVKA		6
<i>Antony</i>		6
<i>Api</i>	181	
<i>Api eller</i>	181	
<i>Api Fin</i>	181	
<i>Api Ordinaire</i>	181	
<i>Api Petit</i>	181	
<i>Api Rose</i>	181	
<i>Api Rouge</i>	181	
APOINT		6
APOINT GROUP	20, 25	
<i>Apoint</i> (synonym of <i>Apoint Orient</i>).....		7
<i>Aporta</i>		3

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Aporta Nalivia</i>		3
APORT ORIENT		7
<i>Aport Oriental</i>		7
<i>Apple of Commerce</i>	66	
<i>Apple of the Well</i>	363	
<i>Apy Rouge</i>	181	
<i>Arabka</i>		32
ARCTIC	24	7
<i>Arcad Krasivui</i>		14
<i>Arcad Krasivui</i>		14
<i>Arcad Krasivui</i>		14
ARKANSAS	24, 47	
<i>Arkansas Baptist</i>		140
ARKANSAS BEAUTY	49	
ARKANSAS BLACK	24, 49	
<i>Arkansas Black</i> (synonym of Arkansas)	47	
<i>Arkansas Black Twig</i>	47, 50	
<i>Arkansas</i>	47	
ARNOLD	50	
<i>Arnold's Beauty</i>	50	
<i>Aromatic Spike No. 351</i>		233
<i>Arsapple</i>	116	
ARTHUR	51	
<i>Astracan Blanch</i>		230
<i>Astracan d'Ete</i>		230
<i>Astracanischer Sommer</i>		230
<i>Astracan Rouge</i>		178
<i>Astrachan</i>		178
<i>Astrachan Red</i>		178
<i>Astrakhan Rouge</i>		178
<i>Astrachan White</i>		230
<i>Astrachan Rouge</i>		178
<i>Astravaskoc</i>		155
<i>Aubertin</i>		3
AUCUBA	52	
<i>Acubafolia</i>	52	
AUGUST		9
<i>August</i> (synonym of July)		112
<i>August Apple</i>		53
AUGUSTINE		10
<i>August Sweet</i>		216
<i>August Sweeting</i>		216
<i>Aunt Dorcas</i>	145	
<i>Aunt Ginnie</i>		75
<i>Aurora</i>		227
<i>Austin</i>	236	
<i>Autumnal Bough</i>		10
<i>Autumnal Swaar</i> (synonym of Autumn Swaar)		11

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Autumnal Swaar</i> (synonym of Autumn Sweet Swaar)....		12
<i>Autumnal Sweet</i>		12
<i>Autumnal Sweet Swaar</i>		12
AUTUMN BOUGH	15	10
<i>Autumn Bough</i> (synonym of Sweet Bough).....		216
<i>Autumn Pearmain</i> (synonym of Winter Pearmain).....	378	
<i>Autumn Pearmain</i> (synonym of Long Island Pearmain)..		124
<i>Autumn Pippin</i>		62
<i>Autumn Rose</i>		100
<i>Autumn Seeknofurther</i>		88
<i>Autumn Strawberry</i>		120
AUTUMN STREAKED		10
AUTUMN SWAAR		11, 12
<i>Autumn Swaar</i> (synonym of Autumn Sweet Swaar)....		12
<i>Autumn Sweet</i>		12
<i>Autumn Sweet Bough</i>		10
AUTUMN SWEET SWAAR		12
<i>Avery Sweet</i>		175
BABBITT	53	
<i>Babouskino</i>		83
<i>Babuscheno</i>		83
<i>Babuschkino</i>		83
<i>Babushkino</i>		83
<i>Baccata fructa oblonga</i>		261
<i>Bachelor</i>	88	
<i>Back Creek</i>	146	
<i>Baer</i>	158	
BAILEY CRIMSON (Crah)		251
<i>Bailey's Golden Sweet</i>	54	
BAILEY SPICE.....		13
<i>Bailey's Spice</i>		13
BAILEY SWEET	54	
<i>Bailey Sweet</i> (synonym of Sweet Winesap).....	333	
BAKER	55	
BAKER SWEET		13
<i>Baker's Sweet</i>		13
BALDWIN.....	12, 15, 17, 18, 19, 24, 32,	56
<i>Baldwin Rosenapfel</i>		56
<i>Baldwin's Rother Pippin</i>		56
<i>Ball Apple</i>		250
<i>Baltimore</i> (synonym of Roseau).....	292	
<i>Baltimore</i> (synonym of Gloria Mundi).....		77
<i>Baltimore</i> (synonym of Vandevere Pippin).....		231
<i>Baltimore Pippin</i> (synonym of Ben Davis).....	69	
<i>Baltimore Pippin</i> (synonym of Gloria Mundi).....		77
<i>Baltimore Red</i>	69	
<i>Baltimore Red Streak</i>	69	
<i>Banana</i>	377	

	VOL. I. PAGE.	VOL. II. PAGE.
BANANA SWEET	60, 377	
BANKS		14
<i>Banks Gravenstein</i>		14
<i>Banks Red Gravenstein</i>		14
BAPTIST	61	
BARBEL	61	
<i>Bard Apple</i>		49
<i>Baranowski</i>		151
<i>Baranowski</i>		151
<i>Barretts Spitzenburgh</i>	225	
BARRINGER	62	
BARRY	63	
<i>Batchellor</i>	88	
BATULLEN	64	
BAXTER	64	
<i>Baxter's Red</i>	64	
BEACH	66	
<i>Beard Burden</i>		49
<i>Beaute de Kent</i>		15
BEAUTIFUL ARCAD		14
<i>Beautiful Arcade</i>		14
<i>Beautiful Pippin</i>		184
<i>Beauty</i>	324	
<i>Beauty of America</i>	45, 319	
BEAUTY OF KENT		15
<i>Beauty of Queen</i>		3
<i>Beauty of the West</i> (synonym of Western Beauty)	239	
<i>Beauty of the West</i> (synonym of Grush)	89	
<i>Beauty Red</i>		40
<i>Bec de Lievre</i>	264	
<i>Beel Solotofskaja</i>		82
<i>Beitigheimer</i>		17
BELBOROLOOSKOE		15
<i>Belle Bonde</i>	75	
<i>Belle Bonne</i>	75	
<i>Belle d'Angers</i>		21
<i>Belle de Boskoop</i>		25
<i>Belle de Boskoop</i>		25
<i>Belle d'Orleans</i>		3
<i>Belle de Rome</i>	290	
<i>Belle Dubois</i>		77
BELLE ET BONNE	67	
<i>Belle Fille</i>	264	
<i>Belle Flavoise</i>	381	
<i>Belle-fleur</i>	381	
<i>Belle-fleur jaune</i>	381	
<i>Belle-Flower</i>	381	
<i>Belleflower Improved</i>	204	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Belle Josephine</i> (synonym of <i>Gloria Mundi</i>).....		77
<i>Belle Josephine</i> (synonym of <i>White Spanish Reinette</i>)....		241
<i>Belle of Boskoop</i>		25
<i>Bellerodovskoe</i>		15
<i>Belle Rose</i>		168
<i>Bell-Flower</i>	381	
<i>Bellflower</i>	381	
<i>Bellflower of the West</i>		148
<i>Bell's Early</i>		203
<i>Bell's Favorite</i>		203
<i>Bellyband</i>	67	
BELMONT	67	
<i>Belmont Late</i>	67	
<i>Belpre Russet</i>	293	
BEN DAVIS	17, 18, 19, 21, 24,	68
<i>Ben Ford</i>		45
<i>Benniger</i>		15
BENNINGER		15
<i>Bennington</i>		203
BENONI		16
BENTLEY	71	
<i>Bentley's Sweet</i>	71	
<i>Bergamot</i>		6
BERGEN	72	
<i>Berry</i>	227	
<i>Berry Apple</i>		39
<i>Bersford</i>	269	
BESS POOL.....	72	
<i>Best Pool</i>	72	
BETHEL	19, 24, 72,	321
BETHLEHEMITE	74	
<i>Bethlemite</i>	74	
<i>Bielborodovska</i>		15
BIETIGHEIMER		17
<i>Big Hill</i> (synonym of <i>Nickajack</i>).....	227	
<i>Big Hill</i> (synonym of <i>Pryor</i>).....	269	
<i>Big Rambo</i> (synonym of <i>Western Beauty</i>).....		239
<i>Big Rambo</i> (synonym of <i>Grosh</i>).....		89
<i>Big Romanite</i> (synonym of <i>Greyhouse</i>).....	153	
<i>Big Romanite</i> (synonym of <i>Pennock</i>).....	255	
<i>Big Sweet</i>		233
<i>Big Vandevere</i>		231
BILLY BOND.....	75	
BIRTH		18
<i>Bishop's Pippin of Nova Scotia</i>	381	
BISMARCK	25	19
<i>Bismark</i>		19
<i>Black American</i>	79	

COMBINED INDEX TO VARIETIES.

299

	VOL. I. PAGE.	VOL. II. PAGE.
BLACK ANNETTE.....	76	20
<i>Black Apple</i> (synonym of <i>Black Jersey</i>).....	79	
<i>Black Apple</i> (synonym of <i>Detroit Red</i>).....		46
<i>Black Baldwin</i>	60	
BLACK BEN DAVIS.....	76	
<i>Black Ben Davis</i> (synonym of <i>Gano</i>).....	135	
<i>Blackburn</i>	88	
<i>Black Detroit</i>		46
BLACK GILLFLOWER.....	12, 16, 17, 18, 32, 77	
<i>Black Jack</i>	153	
BLACK JERSEY.....	79	
BLACK LADY APPLE.....	182	
<i>Black Pennock</i>	153	
<i>Black Spitz</i>	77	
<i>Black Spitzenberg</i>	132	
<i>Black Spy</i>	189	
<i>Black Sweet</i>	219	
<i>Black Twig</i>	247	
<i>Black Vandervere</i>	153	
<i>Black Vandevere</i>	153	
<i>Blair</i>	208	
<i>Blanche</i>		241
<i>Blanche d'Espagne</i>		241
<i>Blanche Glacée d'Été</i>		239
BLenheim.....	20	
<i>Blenheim Orange</i>	21	
<i>Blenheim Pippin</i>	21	
<i>Blenheimsrenett</i>	21	
Bloom.....	69	
<i>Blooming Orange</i>		21
<i>Blue Baldwin</i>	185	
BLUE PEARMAIN.....	17, 18, 19, 24, 80	
BLUE PEARMAIN GROUP.....	24	
<i>Bismen Catrill</i>		85
BLUSHED CHAMBLE.....		22
<i>Blushed June</i>		181
<i>Blush June</i>		181
<i>Rootman's Seedling</i>	350	
<i>Bogdanoff</i> (synonym of <i>Bogdanoff Glass</i>).....	81	
<i>Bogdanoff</i> (synonym of <i>Grandmother</i>).....		83
BODDANOFF GLASS.....	81	
<i>Bogdanoff Steklhanka</i>		83
<i>Bohannon</i>		118
BOIKEN.....	82	
<i>Boiken Apfel</i>	82	
<i>Bonford</i>	269	
BONUM.....		23
BOROVINKA.....		24

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Borovinka Angluskaja</i>		24
<i>Borovitsky</i> (synonym of Oldenburg).....	25	151
<i>Borovitsky</i> (synonym of Borovinka).....		24
<i>Borowicki</i>		151
<i>Borowitski</i>		151
<i>Borowitsky</i>		151
BORSIDORF	83	
<i>Borsdorfer</i> (synonym of Borsdorf).....	84	
<i>Borsdorfer</i> (synonym of Anisim).....		5
BOSKOOP		25
BOSTON Russet	85	
<i>Boston Russet</i> (synonym of Roxbury).....	293	
BOTTLE GREENING	24, 85	
BOUCKEN	86	
<i>Bough</i>		216
<i>Bough, Early Sweet</i>		216
<i>Bough Sweet</i>		26, 216
<i>Bow Apple</i>		216
<i>Bowers Apple</i>	329	
BOYS DELIGHT	86	
<i>Bracken</i>		240
<i>Bracy's Seek-No-Further</i>		88
<i>Brandywine</i>	213	
<i>Bread and Cheese</i>	273	
BRESKOVKA		26
<i>Brier Sweet</i>		251
BRIER (Crab)		251
<i>Brier Sweet</i> (synonym of Brier).....		251
<i>Brier Sweet</i> (synonym of Van Wyck).....		268
BRILLIANT		(v)
BRISTOL	86	
<i>Bristol</i> (synonym of Red Canada).....	276	
<i>Broad River</i>	289	
<i>Brooke Pippin</i>	146	
<i>Brown' Golden Sweet</i>	343	
BROWNLEES	86	
<i>Brownlees Russet</i>	86	
<i>Brownlees Seedling Russet</i>	86	
BROWN SWEET	87	
BUCKINGHAM	18, 88	
<i>Buckley</i>		33
<i>Buckram</i>		97
<i>Buchanan</i>		149
<i>Buler</i>	174	
BULLOCK	15, 89	
<i>Bullock</i> (synonym of Hunt Russet).....	164	
<i>Bullock's Pippin</i> (synonym of Bullock).....	89	
<i>Bullock's Pippin</i> (synonym of Ewalt).....	124	

COMBINED INDEX TO VARIETIES.

301

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Bullripe</i>		40
BUNKER HILL		27
<i>Burlington</i>	225	
<i>Burlington Greening</i>	282	
<i>Burlington Spitzenberg</i>	225	
BUTTER		27
<i>Butter Pippin</i>		78
<i>Byers</i>	88	
<i>Byers Red</i>	88	
<i>Cabane du Chien</i>		98
<i>Cabashate</i>		227
CABASHEA (<i>Winter</i>)	91, 176	
CABASHEA		28
<i>Cabashea</i> (synonym of <i>Twenty Ounce Pippin</i>)	349	
<i>Cabashie</i>		28
<i>Calkin's Pippin</i>		30
<i>Calville Butter</i>	56	
<i>Calville de Gravenstein</i>		85
<i>Calville Grafensteiner</i>	85	
<i>Calville Krasmut</i>		22
<i>Cambour des Lorrains</i>		213
<i>Camoisis du roi d'Espagne</i>		241
<i>Camoise Blanche</i>		241
<i>Camoisee Blanche</i>		241
CAMPFIELD	91	
<i>Camucsar</i>		241
<i>Camuezas</i>		241
<i>Camuzar</i>		241
CANADA BALDWIN	25, 92	69
<i>Canada Peach</i>		159
<i>Canada Pippin</i> (synonym of <i>Canada Reinette</i>)	93	
<i>Canada Pippin</i> (synonym of <i>White Pippin</i>)	368	
<i>Canada Red</i> (synonym of <i>Red Canada</i>)	276	
<i>Canada Red</i> (synonym of <i>Roseau</i>)	292	
<i>Canada Redstreak</i>	276	
CANADA REINETTE	32, 93	
<i>Canada Reinette</i> (synonym of <i>Cheeseboro</i>)		33
<i>Canadian Reinette</i>	93	
<i>Canadisk Reinnet</i>	93	
<i>Caufield</i>	91	
CANNON <i>Pearmain</i>	95	
<i>Capp Mammoth</i>	206	
<i>Cardinale</i>	181	
CARLOUGH	95	
<i>Carnation Apple</i>	96	
<i>Carolina</i> (synonym of <i>Nickajack</i>)	227	
<i>Carolina</i> (synonym of <i>White Juneating</i>)		240
<i>Carolina June</i>		181

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Carolina Red</i>		181
<i>Carolina Red June</i>		181
<i>Carolina Red Streak</i>	69	
<i>Carolina Spice</i>	227	
<i>Caroline</i> (synonym of Nickajack)	227	
<i>Caroline</i> (synonym of White Juneating)		240
CARPENTIN	96	
<i>Carpentin Reinette</i>	96	
<i>Carthouse</i>	138	
CATHEAD		29
<i>Cathead</i> (synonym of Cheeseboro)		33
<i>Cathead</i> (synonym of Fall Pippin)		62
<i>Cathead Greening</i>		29
<i>Catshad</i>		29
<i>Catshad Greening</i>		29
<i>Cattell Apple</i>		148
<i>Caux</i>	114	
<i>Cayuga Red Streak</i>		227
CAYWOOD	96	
CELENTIA		29
<i>Chaltram Pippin</i>	227	
<i>Champion</i>	90	
<i>Champion Red</i>	90	
CHAMPLAIN		30
CHANDLER		31
<i>Chandler's Red</i>		31
CHARLAMOFF		32
<i>Charlamoski</i>		32
<i>Charlamovskoe</i>		32
<i>Charlamowski</i>		151
<i>Charlamowski d'Automne</i>		151
<i>Charlamowskircher Nalleoid</i>		151
<i>Charlamowskoe</i>		32
<i>Charlamowsky</i>		151
<i>Charles Apple</i>	199	
<i>Charlottenthaler</i>		222
<i>Charlottenthaler Apple</i>		222
<i>Charlottenthaler Golba</i>		222
<i>Charmant Blanc</i>		213
<i>Chase</i>	166	
<i>Chatham Pippin</i>	227	
<i>Cheat</i> (synonym of Domine)	109	
<i>Cheat</i> (synonym of Wells)	363	
<i>Cheatan Pippin</i>	227	
<i>Cheataw</i>	227	
<i>Cheesborough</i>		33
CHEESEBORO		33
<i>Cheeseboro's Russet</i>		33

COMBINED INDEX TO VARIETIES.

303

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Cheeseborough</i>		33
<i>Cheeseborough Russet</i>		33
CHENANGO		33
<i>Chenango Strawberry</i>		33
CHERRY (Crab)		252
<i>Chimney Apple</i>		65
<i>Christ Birth</i>		18
<i>Christ Birth Apple</i>		18
<i>Christmas</i>		18
<i>Christmas Apple</i>	181	
<i>Christ's Golden Reinette</i>	114	
<i>Choice Kentuck</i>	311	
<i>Cider</i>	311	
<i>Cider Apple</i>	311	
<i>Cinnamon</i>		6
CLAPPER FLAT		34
CLARKE		35
<i>Clarke Beauty</i>		35
CLANTON	66	
<i>Cling Tight</i>	109	
<i>Clothes-yard Apple</i>	145	
CLYDE		36
<i>Clyde Beauty</i>		36
<i>Coalbrook</i>	225	
<i>Cobbett's Fall</i> (synonym of Fall Pippin)	62	
<i>Cobbett's Fall</i> (synonym of White Spanish Reinette)	241	
<i>Cobbett's Fall Pippin</i> (synonym of Fall Pippin)	62	
<i>Cobbett's Fall Pippin</i> (synonym of White Spanish Reinette)	241	
COLLING	117	
<i>Codlin, Keswick</i>	117	
<i>Coc's Spice</i>	49	
COFFELT	97	
<i>Coffel Beauty</i>	97	
<i>Coggeswell</i>	98	
COGSWELL	98	
<i>Cogswell Pearmain</i>	98	
<i>Coleman</i>		227
<i>Cole's Quince</i>		174
COLLAMER		36
<i>Collamer Twenty Ounce</i>		36
COLLINS	99	
<i>Collins' Red</i>	99	
COLTON		37
<i>Colton Early</i>		37
COLVERT		38
<i>Combermere Apple</i>	206	
<i>Compound</i>	329	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Compte Orloff</i>		233
<i>Comte Woronzoff</i>		3
<i>Concombre Ancien</i> (synonym of Fall Pippin).....		62
<i>Concombre Ancien</i> (synonym of White Spanish Reinette)		241
<i>Conford</i>	269	
<i>Conic June</i>		118
<i>Connecticut Seek-No-Further</i>	364	
CONSTANTINE		39
<i>Coon</i>	107	
<i>Coon Red</i>	107	
COOPER	16	40
COOPER MARKET	109	
<i>Cooper's Red</i> (synonym of Cooper Market)	101	
<i>Cooper's Red</i> (synonym of Etowah).....	123	
<i>Cooper's Redling</i>	101	
<i>Copmanthorpe Crab</i>	114	
<i>Copp's Mammoth</i>		77
<i>Corail</i>		3
CORAL (Crab)		252
CORNELL		40
<i>Cornell Fancy</i>		41
<i>Cornell's Favorite</i>		41
<i>Cornell's Savewell</i>	299	
CORNER		41
CORNISH GILLIFLOWER	138	
<i>Cornish Gilliflower</i> (synonym of Scollop Gilliflower)....		197
<i>Corse's St. Lawrence</i>		193
<i>Cos Orange</i>		42
<i>Coss Champion</i>	99	
<i>Costard</i>		29
<i>Costard Ray</i>		29
<i>Count Orloff</i>		233
<i>Cowstard</i>		29
COX ORANGE		42
<i>Cox's Orange Pippin</i>		42
<i>Cox's Pomona</i>		165
CRANBERRY PIPPIN	300	43
<i>Crandall Seedling</i>		107
<i>Crane's Pippin</i>	244	
CREAM		44
<i>Crimson Beauty</i>		196
<i>Crimson Pippin</i> (synonym of Scarlet Pippin).....		196
<i>Crimson Pippin</i> (synonym of Detroit Red).....		46
<i>Crimson Scarlet Pippin</i>		196
CROTTIS	102	
CROW EGG		44
CROWNS	102	
<i>Cuir, De</i>	264	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Cumming's Rambo</i>		89
CURRENT (Crab)	3	253
Current (Crab)		253
Curtis Greening	331	
Cut Wine		244
Czarskui Schip		45
CZAR THORN		45
Daklonaga	227	
Danc de Menage	200	
DANVERS Sweet	103	
Danvers Winter	103	
Danvers Winter Sweet	15, 103	
Dark Baldwin	60	
DARTMOUTH (Crab)		253
DEACON JONES	104	
DEADERICK		45
Dean's Codlin		165
De Bretagne	93	
De Cuir	334	
De Glace d'Ete		239
De Glace Hative		230
Deil's	201	
De La Madeleine Rouge		55
Delaware	273	
Delaware Red Winter	189	
Delaware Winter	189	
De Lorraine		213
Demary		73
DEMOCRAT	105	
De Muscovie d'Isle		239
De Noige		105
De Notre Dame		213
De Rambourg		213
De Rambour		213
De Ratcau		62
De Ratteau		241
Der Carpentin	96	
De Revel (synonym of Yellow Transparent)		248
De Revel (synonym of Vineuse Rouge)		233
Derrick and Ann		180
Derrick's Graft		180
Der Schwere Apfel	320	
De Seigneur d'Automne		29
D'Espagne (synonym of Fall Pippin)		62
D'Espagne (synonym of White Spanish Reinette)		211
Detrding's Early		178
Detroit (synonym of Ortley)	244	
Detroit (synonym of Detroit Red)		46

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Detroit Black</i>		46
DETROIT RED	12	46
<i>Detroit of the West</i>	244	
<i>D'Eve</i>		141
<i>De Vin du Conn</i>		227
<i>Dewitt Apple</i>	107	
<i>De Witt</i>	107	
<i>Dickenson</i>	106	
DICKINSON	106	
<i>Die Haarlemer Reinette</i>	93	
<i>Diels Sommerkönig</i>		85
<i>Die Weiberreinette</i>	93	
DISHAROON	107	
<i>Dix-huit Onces</i>		227
DOCTOR	17, 107	
<i>Doctor Dewitt</i>	107	
DOCTOR WALKER	108	
<i>Dodge's Black</i>	79	
<i>Dodge's Early Red</i>		203
DOMINE	17, 109	
<i>Domine</i> (synonym of Wells?)	363	
<i>Dominie</i>	109	
<i>Doppelte Casselar Reinette</i>	114	
<i>D'Or d'Angleterre</i>	141	
DOUBLE ROSE	110	
<i>Douse</i>		95
<i>Downing's Winter Maiden Blush</i>	152	
<i>Dows</i>		95
<i>Dowse</i>		95
<i>Dredge's Fame</i>		21
DU BOIS	111	
<i>Duchess</i>		151
<i>Duchesse d'Oldenbourg</i>		151
<i>Duchess No. 3</i>		158
<i>Duchess of Oldenburg</i>		48, 151
<i>Duchess of Oldenburgh</i>		151
<i>Ducks Bill</i>	378	
DUDLEY		48
<i>Dudley Winter</i>		48
<i>Duitsch Mignonne</i>	114	
DUKE OF DEVONSHIRE	111	
<i>Duke of Wellington</i>	112	
<i>Du Marechal</i>		65
DUMELOW	112	
<i>Dumelow's Crab</i>	112	
<i>Dumelow's Pippin</i>	112	
<i>Dumelow's Seedling</i>	112	
<i>Dumpling</i>		208

	VOL. I. PAGE.	VOL. II. PAGE.
DUNCAN	113	
<i>Durable Trois ans</i>	116	
<i>Dutchess</i> ..		151
<i>Dutchess of Oldenberg</i>		151
<i>Dutch Greening</i>	253	
DUTCH MIGNONNE	32, 114	
<i>Dutch Mignonne</i> (synonym of Blenheim)		21
<i>Dutch Minion</i>	114	
<i>Duverson's June</i>		141
DUZENBURY	115	
DYER		49
<i>Early Baldwin</i>		168
<i>Early Bough</i>		216
EARLY CHANDLER	16	
<i>Early Colton</i>		37
<i>Early Congress</i>		85
<i>Early French Reinette</i> (synonym of Early Harvest)		51
<i>Early French Reinette</i> (synonym of Sweet Bough)		216
<i>Early Golden Sweet</i>		81
EARLY HARVEST	15, 17, 18	50
<i>Early Jack</i>		166
<i>Early Jenetting</i>		240
EARLY JOE		52
<i>Early July Pippin</i>		51
<i>Early June</i> (synonym of Early Harvest)		51
<i>Early June</i> (synonym of Egg Top)		56
<i>Early June of South</i>		141
<i>Early Margaret</i>		141
<i>Early May</i>		240
EARLY PENNOCK		53
<i>Early Red</i>		141
<i>Early Red Juneating</i>		141
<i>Early Red Margaret</i>		141
<i>Early Red Pippin</i>		93
<i>Early Redstreak</i>		93
<i>Early Red Streak</i>		93
EARLY RIPE		54
EARLY STRAWBERRY	15, 17	54
<i>Early Striped Juneating</i>		141
<i>Early Summer Pearmain</i>		211
<i>Early Sweet</i>		98
<i>Early Sweet Bough</i>		216
<i>Early Sweetheart</i>		216
<i>Early Tart Harvest</i>		168
<i>Early Washington</i>		203
<i>Edelborsdorfer</i>	84	
<i>Edgar County Red Streak</i>	357	
<i>Edgar Red Streak</i>	357	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Edgerly Sweet</i>	54	
EDWARDS	115	
<i>Edwards</i> (synonym of Nickajack)	227	
<i>Edwards Favorite</i>	115	
<i>Edward Shantee</i>	227	
<i>Egg Jap</i>		44
EGG TOP		56
<i>Eighteen Ounce</i>		227
<i>Eighteen Ounce Apple</i>		227
EISER	116	
<i>Eiser Rouge</i>	116	
ELGIN PIPPIN		56
<i>Elgin Pippin</i> (synonym of White Spanish Reinette)		241
ELLSWORTH	117	
<i>Empereur Alexandre I</i>		3
<i>Empereur Alexandre de Russie</i>		3
<i>Empereur de Russie</i>		3
<i>Emperor Alexander</i>		3
<i>Englar Orange Appel</i>		154
<i>Englische Granat-Reinette</i>		184
<i>Englischer Pepping</i>		57
<i>English Beauty of Pennsylvania</i>	109	
<i>English Borovinka</i>		24
ENGLISH CODLING		117
<i>English Golden</i>	143	
<i>English Golden Pippin</i>	141	
<i>English Golden Russet</i>	143	
<i>English Jannetting</i>		230
<i>English King</i>		3
ENGLISH PEARMAIN	16	
<i>English Pearmain</i> (synonym of Long Red Pearmain)		125
ENGLISH PIPPIN		57
<i>English Pippin</i> (synonym of Longfield)		123
<i>English Pippin</i> (synonym of Golden Reinette)	142	
<i>English Rambo</i> (synonym of Grosh)		89
<i>English Rambo</i> (synonym of Domine)	109	
<i>English Rambo</i> (synonym of Wells)	363	
<i>English Red Streak</i> (synonym of Domine)	109	
<i>English Red Streak</i> (synonym of Redstreak)	278	
<i>English Red Streak</i> (synonym of Wells)	363	
<i>English Redstreak</i>	278	
ENGLISH RUSSET	118	
<i>English Russet</i> (synonym of Long Island Russet I)	194	
<i>English Spitzenberg</i>	225	
<i>English Spitzenberg</i>	225	
<i>English Sweet</i>		58, 175
<i>English Sweeting</i>		175
<i>English Wandevere</i>	312	

COMBINED INDEX TO VARIETIES.

309

	VOL. I. PAGE.	VOL. II. PAGE.
<i>English Winter Red Streak</i> (synonym of Domine)	109	
<i>English Winter Red Streak</i> (synonym of Wells)	363	
<i>Episcopal</i>		62
<i>Episcopale</i>		241
<i>Eppes Sweet</i>	103	
<i>Epses Sweet</i>	103	
<i>Equinately</i>	88	
<i>Ernst's Apple</i>		149
<i>Ernst's Pippin</i>		149
<i>Esopus</i>	120	
<i>Esopus Spitzenberg</i>	121	
<i>Isopus Spitzenberg</i>	121	
<i>Esopus Spitzenberg</i> (New)	172	
<i>Esopus Spitzenburg</i>	12, 15, 16, 17, 18, 19, 24, 120	
<i>Esopus Spitzenburgh</i>	121	
<i>Essex Pippin</i>		184
ETOWAH	123	
<i>Etowah</i> (synonym of Cooper Market)	101	
ETRIS	123	
<i>Eve</i>		56
<i>Eve Apple</i> (synonym of Egg Top)		56
<i>Eve Apple</i> (synonym of Margaret)		141
<i>Evebearing</i>		108
EVENING PEAR	123	
EWALD	124	
EXCELSIOR (Crab)		254
<i>Faldtvalder</i>	125	
FALIX	125	
FALLAWATER	18, 32, 125	
<i>Fall Bough</i>		10
<i>Fall de Waldes</i>	125	
<i>Fallentvalder</i>	125	
<i>Fall Genetling</i>		59
<i>Fall Gennetling</i>		59
FALL GREENING		58
<i>Fall Green Sweet</i>		233
FALL HARVEY		58
<i>Fall Jenetling</i>		59
FALL JENNETTING		59
<i>Fall Jennetting</i>		59
FALL ORANGE		12, 60
FALL PIPPIN	12, 15, 16, 17, 18, 24	61
FALL PIPPIN GROUP	24	
<i>Fall Pippin</i> (synonym of Fall Harvey)		58
<i>Fall Pippin</i> (synonym of Holland Pippin)		101
<i>Fall Pippin</i> (synonym of White Spanish Reinette)		241
<i>Fall Queen</i> (synonym of Buckingham)	88	
<i>Fall Queen</i> (synonym of Haas)		91
VOL. II.—25		

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Fall Queen of Kentucky</i>	88	
<i>Fall Romanite</i>	273	
<i>Fall Strawberry</i>		120
<i>Fall Stripe</i>		196
<i>Fall Swaar</i>		11
<i>Fall Swaar of the West</i>		11
<i>Fall Vanderzere</i>		231
FALL WINE		63
<i>Fall Winesap</i>	89	
FAMEUSE.....	15, 16, 17, 18, 19, 23,	25
<i>Fameuse</i> (synonym of <i>Utter</i>)		230
<i>Fameuse Baldwin</i>		98
FAMEUSE, GREEN		69
FAMEUSE GROUP	20, 25	68
FAMEUSE NOIRE		69
FAMEUSE SURE		69
FAMILY	127	
FANNY		69
FARRIS	128	
<i>Faust's Rome Beauty</i>	290	
<i>Fay's Russet</i>	164	
<i>Felch</i>	56	
<i>Femme de Menage</i>	206	
FERDINAND	129	
FERRIS	130	
<i>Feuilles D'Aucuba</i>	52	
<i>Filliken</i>	153	
<i>Fin d'Automne</i>		3
<i>Fine Winter</i>	373	
<i>Fink</i>	336	
<i>Fink's Seedling</i>	336	
FISHKILL	32	70
<i>Fishkill Beauty</i>		70
<i>Five-Quartered Gilliflower</i>		197
<i>Flanders Pippin</i>	206	
<i>Flat</i>		34
<i>Flat Spitzenburg</i>	345	
<i>Flint Russet</i>		170
FLORENCE	130	
FLORENCE (Crab)		255
FLORY	131	
<i>Flory Bellflower</i>	131	
<i>Flower</i> (of <i>Genesee</i>)		73
<i>Flushing</i> (synonym of <i>Flushing Spitzenburg</i>)	132	
<i>Flushing</i> (synonym of <i>Newtown Spitzenburg</i>)	225	
<i>Flushing Seek-No-Further</i>		88
<i>Flushing Spitzenburg</i>	132	
FLUSHING <i>Spitzenburg</i>	132	

COMBINED INDEX TO VARIETIES.

311

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Flushing Spitzenburg</i>	132	
<i>Flushing Spitzenburg</i> (synonym of Roseau)	292	
<i>Flushing Spitzenburgh</i>	132	
FORD		71
FOREST	133	
<i>Forever Pippin</i>		33
<i>Formosa</i>		184
<i>Formosa Pippin</i>		184
<i>Fornwalder</i>	125	
<i>Forsythe's Seedling</i>	227	
<i>Fourth of July</i>		112
<i>Fowler</i>	311	
<i>Fraise</i>		55
FRAKER	133	
<i>Fraker's Seedling</i>	133	
FRANCHOT		71
<i>Frank</i>		33
<i>Frankfort Queen</i>	88	
<i>Frank Rambour</i>		213
FRENCH PIPPIN	24, 134	
<i>French Pippin</i> (synonym of Holland Pippin)		101
<i>French Pippin</i> (synonym of Roman Stem)		188
<i>French Pippin</i> (synonym of Newark Pippin)	223	
<i>French Rambo</i>		89
<i>French Reinette</i>		215
<i>French Russet</i>	261	
<i>French Spitzenburgh</i>		215
<i>French Spitzenburg of Vermont</i>	292	
<i>Fuller</i>	311	
FULLERTON SWEET		71
<i>Fullhauser</i>	60	
<i>Gait</i>	67	
GANO	21, 135	
GARDEN ROYAL		72
<i>Gardener's Apple</i>		143
GARDNER SWEET PEARMAIN		73
<i>Gale</i>	67	
<i>Gay's Romanite</i>	255	
<i>Gelber Bellefleur</i>	381	
<i>Gelber Englischer Schönblühender</i>	381	
<i>Gelee d'Ete</i>		239
<i>General Chandler</i>		31
GENESEE FLOWER		73
<i>Genet</i>	271	
<i>Genetion</i>	271	
<i>Geneva Pearmain</i>		30
GENEVA PIPPIN	136	
<i>Geniton</i>	271	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Gennetin</i>	271	
<i>Genneting</i>	271	
<i>Gennetting</i> (synonym of Ralls)	271	
<i>Gennetting</i> (synonym of White Juneating)		240
<i>Germaine</i>	378	
<i>German Calville</i>		6
<i>German Calville 324</i>		6
<i>German Green</i>	93	
<i>German Spitzenberg</i>	153	
<i>Georgia June</i>		181
GIBB (Crab)		256
<i>Gibbon's Smokehouse</i> (synonym of Vandevere Pippin)....		231
<i>Gibbons Smokehouse</i> (synonym of Smokehouse)	312	
GIDEON		74
GIDEON SWEET	24, 137	
<i>Gideon White</i>		74
<i>Gillet's Seedling</i>	290	
GILLIFLOWER	16	
<i>Gillflower</i>	77, 138	
GILPIN	17, 34, 138	
<i>Ginet</i>	271	
<i>Ginetting</i>		240
GINNIE		75
GIVENS	140	
<i>Glacé de Zelande</i>		239
<i>Glacé d'Ete</i>	239	
GLADSTONE	25	75
<i>Glazewood</i>		77
<i>Glazewood Gloria Mundi</i>		77
GLENLOCH	140	
GLORIA MUNDI		76
<i>Glory of York</i>		184
<i>Gloucester Pippin</i>		21
<i>Golden Apple</i>		83
<i>Golden Beauty</i>		269
<i>Golden Gray</i>	328	
GOLDEN MEDAL	141	
GOLDEN PIPPIN	16, 141	
GOLDEN PIPPIN (I)		78
GOLDEN PIPPIN (II)		79
<i>Golden Pippin</i> (synonym of Fall Pippin)		62
<i>Golden Pippin</i> (synonym of Golding)		83
<i>Golden Pippin</i> (synonym of Belmont)	67	
<i>Golden Pippin</i> (synonym of Ortley)	244	
GOLDEN PIPPINS		78
GOLDEN RED	142	
GOLDEN REINETTE	142	79
<i>Golden Rennet</i>	142	

COMBINED INDEX TO VARIETIES.

313

	VOL. I. PAGE.	VOL. II. PAGE.
GOLDEN RUSSET	17, 18, 143, 295	
<i>Golden Russet</i> (synonym of Bullock)	89	
<i>Golden Russet</i> (synonym of Perry Russet)	257	
<i>Golden Russet, American</i>	89	
<i>Golden Russet</i> (not of New York)	164	
<i>Golden Russet</i> of Massachusetts.....	164	
<i>Golden Russet</i> of New England.....	164	
<i>Golden Russet</i> of New York	143	
<i>Golden Russet</i> of Western New York	143	
<i>Golden Spice</i>		49
GOLDEN SWEET		81
<i>Golden Sweet</i> (synonym of Northern Sweet)		147
<i>Golden Sweetening</i>		81
GOLDEN WHITE		82
GOLDING		82
<i>Gold Medal</i>	141	
<i>Goldreinette von Blenheim</i>		21
<i>Good Peasant</i> (synonym of Longfield)		123
<i>Good Peasant</i> (synonym of Anisim)		5
<i>Gov. Seaward's</i>		227
<i>Gowden</i>	227	
<i>Gowdie</i>	227	
<i>Grafen-Apfel</i>		85
<i>Grafensteiner</i>		85
<i>Graham's Red Warrior</i>	227	
<i>Grenat-Reinette</i>		184
<i>Grand Alexander</i>		3
<i>Grand Alexandre</i>		3
<i>Grand Duc Constantin</i>		39
<i>Grand Duke Constantine</i>	25	39
GRANDMOTHER		83
<i>Grand Mother</i>		83
<i>Grandmother's Apple</i>	145	
<i>Grand Sultan</i> (synonym of Vineuse Rouge)		233
<i>Grand Sultan</i> (synonym of Yellow Transparent)		248
GRANITE BEAUTY	145	
<i>Grantham</i>		108
GRAVENSTEIN	12, 15, 17, 18	84
<i>Gravensteiner</i>		85
<i>Gravenstein Rouge</i>		180
<i>Gravenstine</i>		85
<i>Grave Slige</i>		85
<i>Grave Slige</i>		85
<i>Gray Apple</i> (synonym of McAfee)	196	
<i>Gray Apple</i> (synonym of Pomme Grise)	264	
<i>Gray Baldwin</i>	60	
<i>Grayhouse</i>	153	
<i>Gray Romanite</i> (synonym of Greyhouse)	153	
<i>Gray Romanite</i> (synonym of Rambo)	273	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Gray's Keeper</i>	196	
<i>Greasy Pippin</i> (synonym of Ortley)	244	
<i>Greasy Pippin</i> (synonym of Lowell)		128
GREAT BARBE	145	
GREAT MOGUL		87
<i>Great Pearmain</i>	378	
<i>Green Bellflower</i>	244	
<i>Greening</i>	145, 282	
<i>Green Mountain Pippin</i> (synonym of Fallawater).....	125	
<i>Green Mountain Pippin</i> (synonym of Virginia Greening)...	352	
GREEN NEWTOWN	17, 18, 24, 145, 149	
<i>Green Newtown Pippin</i>	146	
<i>Green Newton Pippin</i> (synonym of Rhode Island Green- ing)	282	
GREEN PIPPIN	16	
GREEN SEEK-NO-FURTHER		88
GREEN SWEET	150	
<i>Green Sweet</i> (synonym of Repka Malenka)	282	
<i>Green Sweet</i> (synonym of Victuals and Drink)		233
GREEN SWEETING	151	
<i>Green Sweeting</i> (synonym of Green Sweet)	151	
<i>Green Transparent</i>		233
GREENVILLE	152	
<i>Green Winter Pearmain</i>	378	
<i>Green Winter Pippin</i>	146	
GREYHOUSE	153	
GRIMES	18, 19, 23, 153	
<i>Grimes Golden</i>	154	
<i>Grimes Golden Pippin</i>	154	
<i>Grindstone</i>	45	
<i>Grise</i>	204	
<i>Gros-Alexandre</i>		3
<i>Gros Api Rouge</i>	181	
GROSH		80
<i>Grosh</i> (synonym of Summer Rambo).....		213
<i>Grosh's Mammoth</i>		80
<i>Groszkoe Selenka Grüner</i>		233
<i>Gros Pomier</i>		91
<i>Gros Pommier</i>		91
<i>Gros-Rambour d'Ete</i>		213
<i>Gros Rambour d'Hiver</i>	206	
<i>Grosse Casselar Reinette</i>	114	
<i>Grosse Reinette d'Angleterre</i>	93	
<i>Grosser Mogul</i>		87
<i>Grosse-Schatnase</i>		20
GRUNDY		90
<i>Gul. Bellefleur</i>	381	
HAAS		91

	VOL. I. PAGE.	VOL. II. PAGE.
HAGLOE		92
<i>Hagloe Crab</i>		92
<i>Hampers American</i>		178
<i>Hampshire Greening</i>	282	
<i>Hang-On</i>	339	
<i>Hard Red</i>	153	
<i>Hardwick</i>	326	
HARGROVE	155	
<i>Harmony</i>		53
<i>Harrigan</i>	208	
<i>Harvest</i> (synonym of Early Harvest)		51
<i>Harvest</i> (synonym of Primate)		168
<i>Harvest Apple</i>		215
HARVEST REDSTREAK		93
<i>Harvey</i>		58
HASKELL		93
<i>Haskell Sweet</i>		94
<i>Hass</i>		91
<i>Haus Mütterchen</i>	206	
<i>Haverstraw Pippin</i>		30
<i>Hawkins Pippin</i>		73
HAWLEY	15, 24	94
<i>Hawley</i> (synonym of Hawthornden)		96
<i>Hawthornden</i>		90
HAWTHORNDEN		96
<i>Hays</i>	373	
<i>Hays Apple</i>	373	
<i>Hays Wine</i>	373	
<i>Hays Winter</i>	373	
<i>Hays Winter Wine</i>	373	
HAYWOOD	155	
HAZEN	156	
<i>Heaster</i>	158	
<i>Heicke's Summer Queen</i>		53
<i>Heister</i>	158	
HELEN	183	
<i>Hempstead</i>	321	
<i>Hendrick</i>	333	
<i>Hendrick Sweet</i>	333	
HENNIKER	156	
<i>Henrick</i>	333	
<i>Henrick Sweet</i>	333	
<i>Henry Sweet</i>	333	
<i>Henshaw</i>	88	
<i>Herbstbreitling</i>		213
<i>Herbst Streifling</i>		10
<i>Herbst Streifling</i>		10
HEREFORDSHIRE	157	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Herefordshire Beefing</i>	157	
<i>Herefordshire Pearmain</i>	378	
<i>Herefordshire Redstreak</i>	278	
<i>Herr's June</i>		141
<i>Herefordshire Pearmain</i>	378	
HIBERNAL	26	96
HICKS		97
HIESTER	158	
HIGHLAND BEAUTY	159, 183	
<i>Highland Pippin</i>		168
HIGHTOP SWEET		97
<i>High Top Sweet</i>		98
<i>High Top Sweet</i> (synonym of Amsterdam)		5
<i>High Top Sweetening</i>		98
HILAIRE		69, 98
HILTON		99
<i>Himbeerapfel Lieslander</i>		129
<i>Himbeerapfel Lirlander</i>		129
<i>Hishop</i>		257
HOADLEY		99
<i>Hogan</i>	109	
HOG ISLAND SWEET		100
<i>Hogpen</i>		60
<i>Holden</i>		69
<i>Holden Pippin</i>		60
HOLLAND PIPPIN		101
<i>Holland Pippin</i> (synonym of Fall Pippin)		62
<i>Holland Pippin</i> (synonym of Domine)	24, 109	
<i>Holland Pippin</i> (synonym of Holland Winter)	159	
<i>Holland's Red Winter</i>	375	
HOLLAND WINTER	159	
<i>Hollow Cored Pippin</i>	244	
<i>Hollow Core Pippin</i>	244	
<i>Hollow Crown Pearmain</i> (synonym of Wine)	373	
<i>Hollow Crown Pearmain</i> (synonym of Long Island Pear- main)		124
HOLMES	161	
HOLMES SWEET	161	
<i>Hominy</i>		203
<i>Honey Greening</i>	16, 151	
<i>Honey Sweet</i>	380	
HOOKE		193
<i>Hoop</i>	153	
<i>Hoopes</i>	153	
<i>Hoopes Pearmain</i>	153	
<i>Hoops</i>	153	
<i>Hopsy</i>	153	
<i>Hopson</i>	153	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Horning</i>		203
<i>Horse</i>		91
<i>Hoss</i>		91
<i>House</i> (synonym of Greyhouse).....	153	
<i>House</i> (synonym of Fall Wine).....		64
<i>Howard</i>	227	
HOWARD BEST		103
<i>Howard Russet</i>		33
<i>Howard's Best</i>		103
<i>Howard's Best Russian</i>		103
<i>Howard's Sweet</i>	54	
<i>Howe Apple</i>		245
<i>Howe</i>		64
<i>Howen</i>		60
<i>Hubbard</i>	227	
HUBBARDSTON	17, 18,	161
<i>Hubbardston Nonsuch</i> (synonym of Hubbardston).....		161
<i>Hubbardston Nonsuch</i> (synonym of Sutton).....		324
<i>Hudson Red Streak</i>		125
HUNTER PIPPIN		104
HUNT RUSSET		104
<i>Hunt's Fine Green Pippin</i>	146	
<i>Hunt's Green Newtown Pippin</i>	149	
HUNTSMAN	18,	105
<i>Huntsman's Ear rite</i>		105
<i>Hurlbut</i>		105
HURLBUT		104
<i>Hurlbut</i> (synonym of Ramsdell Sweet).....		175
<i>Hurlbut Stripe</i>		105
<i>Hurlbut Sweet</i>		175
<i>Hurlbut</i>		105
HYDE KING		166
<i>Hyde's King (of the West)</i>		166
HYSLOP (Crab)		257
<i>Illinois Greening</i>	331	
<i>Imperial Vandevere</i>		231
<i>Imperial White</i>		157
<i>Imperatrice Eugenie</i>		21
<i>Imperatrice Josephine</i>		77
<i>Indian</i>	106	
<i>Indiana Jannetling</i>	271	
<i>Indiana Vandevere</i>		231
<i>Indian Queen</i>		53
<i>Ingraham</i>	107	
INGRAM	24,	167
<i>Ingram Seedling</i>		167
<i>Inman</i>	244	
INMAN		106

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Isham Sweet</i>		106
<i>Isle of Wight Orange</i> (synonym of Marigold).....	202	
<i>Isle of Wight Orange</i> (synonym of Orange Pippin)....		154
<i>Isle of Wight Pippin</i> (synonym of Marigold).....	202	
<i>Isle of Wight Pippin</i> (synonym of Orange Pippin).....		154
JACK		106
<i>Jack Apple</i>		106
JACKSON	168	
<i>Jackson</i> (synonym of Amos).....	46	
<i>Jackson</i> (synonym of Chenango).....		33
<i>Jackson Red</i>	227	
<i>Jackson Seedling</i>	168	
<i>Jackson Winesap</i>	311	
<i>Jacks Red</i>	135	
<i>Jacobs</i>	169	
JACOBS SWEET	169	
<i>Jacobs Winter Sweet</i>	169	
<i>James River</i> (synonym of Limbertwig).....	193	
<i>James River</i> (synonym of Willow).....	370	
<i>Janet</i>	271	
<i>Janetting</i>	271	
<i>Januarea</i>	93	
JARVIS		107
<i>Jefferies</i>		108
JEFFERIS		108
JEFFERSON COUNTY		109
<i>Jefferson Pippin</i>	271	
<i>Jeffries</i>		108
<i>Jellyflower</i>		197
<i>Jenlon</i>	271	
<i>Jenneting</i>		240
<i>Jennett</i>	271	
<i>Jennette</i>	271	
<i>Jennetting</i>		109
<i>Jennetting</i> (synonym of White Juneating).....		240
<i>Jennings</i>		166
<i>Jenniton</i>	271	
<i>Jersey Black</i>	79	
<i>Jersey Greening</i> (synonym of Ortley).....	244	
<i>Jersey Greening</i> (synonym of Rhode Island Greening)....	282	
JERSEY SWEET	12	110
<i>Jersey Sweeting</i>		110
JEWETT Red	19, 170	
<i>Jewett's Fine Red</i>	170	
<i>Jewett's Red</i>	170	
<i>Joaneting</i>		240
<i>Joe Berry</i>	225	
<i>Joe Précoce</i>		52

COMBINED INDEX TO VARIETIES.

319

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Johnathan</i>	172	
<i>John May</i>	161	
<i>Johnson</i>	278	
<i>Johnson's Fine Winter</i>	385	
<i>Jolly Gentleman</i>		3
JONATHAN	17, 18, 24, 32, 172	
JONATHAN BULER	174	
<i>Jonathan of Buler</i>	174	
<i>Jonathan of the North</i>		5
<i>Jones Early Harvest</i>		181
<i>Jones Pippin</i>		60
<i>Jones Seedling</i>	43	
<i>Josephine</i> (synonym of <i>Gloria Mundi</i>).....		77
<i>Josephine</i> (synonym of <i>White Spanish Reinette</i>)		241
<i>Josie Moore</i>	219	
JUBSON		111
JULY		112
<i>July Apple</i>		168
<i>July Early Pippin</i>		51
<i>July, Fourth of</i>		112
<i>July Pippin</i>		51
<i>June</i> (synonym of <i>Margaret</i>).....		141
<i>June</i> (synonym of <i>Red June</i>).....		181
<i>Juneating</i>		240
<i>Juneating</i>		109
<i>Juneating</i> (synonym of <i>White Juneating</i>).....		240
<i>Juneting</i>		240
<i>Juniata</i>		234
KAIGHN	183	113
<i>Kaighn's Spitzenburg</i>		113
<i>Kaighn's Spitzenberg</i>		113
<i>Kaighn's Spitzenbergh</i>		113
<i>Kaighn's Spitzenburg</i> (synonym of <i>Kaighn</i>).....		113
<i>Kaighn's Spitzenburg</i> (synonym of <i>Long Red Pearmain</i>)..		125
<i>Kaighn's Spitzenburgh</i>		113
<i>Kaighn's Spitzenburg</i>		113
<i>Kaiser Alexander</i>		3
<i>Kajabazcha</i>		114
KALKIDON		114
<i>Kalkidonskoe</i>		114
<i>Kalkidouskoe</i>		114
<i>Kalkidovskoe</i>		114
<i>Kaleil jeltui</i>		247
<i>Kaleil scholtti</i>		247
<i>Kanada Reinette</i>	94	
<i>Kanada-renett</i>	93	
<i>Kansas</i>	175	
KANSAS GREENING	175	

	VOL. I. PAGE.	VOL. II. PAGE.
KANSAS KEEPER	175	
<i>Karaboff</i>		114
KARABOVKA		114
<i>Karabovska</i>		115
<i>Kelley's Sweet</i>	219	
<i>Kelly White</i>	67	
<i>Kempster's Pippin</i>		21
<i>Kennebec Seedling</i>		245
<i>Kent Beauty</i>		15
<i>Kent Fillbasket</i>		115
<i>Kentish Filbasket</i>		115
KENTISH FILLBASKET		115
<i>Kentish Pippin</i>		15
KENT PIPPIN	10	
<i>Kentucky Bellflower</i>		125
<i>Kentucky Gullflower</i>		125
<i>Kentucky Pippin</i>	60	
<i>Kentucky Queen</i>	88	
<i>Kentucky Red Streak</i>	357	
<i>Kentucky Streak</i>	60	
KESWICK		116
<i>Keswick Codlin</i>		117
<i>Keswick Codling</i>		117
<i>Kettageska</i>	178	
<i>Keystone</i>	153	
<i>Khalkidonskoe</i>		114
<i>Kinderhook Pippin</i>		77
<i>King</i> (synonym of Buckingham).....	88	
<i>King</i> (synonym of Tompkins King).....	176, 345	
<i>King</i> (synonym of Twenty Ounce Pippin).....	349	
<i>King Apple</i>	345	
<i>King George the Third</i>	84	
<i>King of Tompkins County</i>	345	
<i>King Philip</i>	172	
<i>Kingsbury Russet</i> (synonym of Cheeseboro).....		33
<i>Kingsbury Russet</i> (synonym of Pumpkin Russet).....		170
KINNARD	18, 176	
<i>Kinnaird's Choice</i>	176	
<i>Kinnaird's Favorite</i>	176	
<i>Kinnard</i>	176	
<i>Kinnard's Choice</i>	176	
KIRKBRIDGE		118
<i>Kirkbridge White</i>		118
<i>Kirk's Golden Reinette</i>	142	
KIRKLAND	177	
KITTAGESKFE	18, 178	
<i>Kleine Grane Reinette</i>	96	
<i>Kleiner Api</i>	181	

COMBINED INDEX TO VARIETIES.

321

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Knight's Red June</i>		181
<i>Krallen Apfel</i>		3
<i>Kountz</i>	225	
<i>La Belle Fameuse</i>		65
LACKEE	179	
<i>Lacquer</i>	179	
<i>Ladies Favorite of Tennessee</i>	88	
<i>Ladies Sweet</i> (synonym of Lady Sweet).....	184	
<i>Ladies Sweet</i> (synonym of Sweet Winesap).....	333	
<i>Ladies Sweeting</i>	184	
LADY	15, 17, 32, 180	
<i>Lady Apple</i>	181	
<i>Lady Blush</i>		139
<i>Lady de Grey's</i>		115
LADY FINGER	183	118
<i>Lady Finger</i> (synonym of Kaighn).....	183	113
<i>Lady Finger</i> (synonym of Long Red Pearmain).....		125
<i>Lady Finger Pippin</i>	183	
LADY GROUP	182	
<i>Lady Henniker</i>	156	
<i>Lady Pippin</i>	66	
<i>Lady's Apple</i>		242
LADY SEEDLINGS	183	
<i>Lady Suffolk</i>		126
<i>Lady Sutherland</i>		126
<i>Lady's Sweet</i>	184	
<i>Lady's Sweeting</i>	184	
LADY SWEET	184	
<i>Lady Sweet</i> (synonym of Sweet Winesap).....	333	
<i>Lady Sweeting</i>	333	
<i>Lady Washington</i> (synonym of Yellow Bellflower).....	381	
<i>Lady Washington</i> (synonym of Cooper).....		40
<i>La Fameuse</i>		65
<i>Lambertwig</i>	193	
<i>Lammas</i>		142
<i>Lancaster Queen</i>		212
LONDON	185	
LANDSBERG		119
<i>Landsberger Reinette</i>		119
<i>Landsburg</i>		119
<i>Landsburger Reinette</i>		119
<i>Lane Albert</i>	268	
<i>Lane's Prince Albert</i>	268	
<i>Langerfeldskoe</i>		123
<i>Langford</i>	186	
LANKFORD	18, 186	
<i>Lankford's Seedling</i>	186	
<i>Lansberger Reinette</i>		119

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Lansinburg</i>	188	
LANSINGBURG	187	
<i>Lansingburgh</i>	188	
<i>Lansingburg Pippin</i>	188	
<i>Laquier</i>	179	
<i>Large Bough</i>		216
<i>Large Early</i>		51
<i>Large Early Bough</i>		216
<i>Large Early Harvest</i>		51
<i>Large Early Yellow Bough</i>		216
<i>Large Fall</i>		241
<i>Large Fall Pippin</i>		241
<i>Large Golden Pippin</i> (synonym of Champlain)		30
<i>Large Golden Pippin</i> (synonym of Golden Pippin)		78
LARGE LADY APPLE	182	
<i>Large Newtown Pippin</i>	146	
<i>Large Rambo</i> (synonym of Rambo)	273	
<i>Large Rambo</i> (synonym of Grosh)		89
<i>Large Red and Green Sweetling</i>		177
LARGE RED SIBERIAN (Crab)		258
<i>Large Red Sweetling</i>		177
<i>Large Romanite</i>	255	
<i>Large Striped Pearmain</i>	196	
<i>Large Striped Winter Pearmain</i>	196	
<i>Large Summer Rambo</i>		89
<i>Large Sweet Bough</i>		216
<i>Large Vanderveer</i>		231
<i>Large White Juncating</i>		51
<i>Large Winter Red</i>	373	
<i>Large Yellow Bough</i>		216
<i>Large Yellow Newton Pippin</i>	146	
<i>Large Yellow Newtown Pippin</i>	146	
LARGE YELLOW SIBERIAN (Crab)		258
<i>La Rue</i>	64	
<i>Late Baldwin</i>	56	
<i>Late Bough</i>		10
<i>Late Chandler</i>		31
LATE DUCHESS	25	
<i>Late Golden Sweet</i>		13
LATE STRAWBERRY		120
LATHAM		121
LA VICTOIRE	188	69
<i>La Victoria Seedling</i>	188	
LAWVER	189	
LEAD		121
<i>Lead Apple</i>		121
<i>Leanham</i>	227	
<i>Leather Apple of Turic</i>	264	

COMBINED INDEX TO VARIETIES.

323

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Lecker</i>	179	
<i>Leder</i>	264	
LEEDS BEAUTY		106
LEE SWEET	191	
LEIGH Greening	134, 192	
<i>Lexington Queen</i>	88	
<i>Liceland Raspberry</i>		129
<i>Lille Api</i>	181	
LILLY OF KENT	193	
<i>Lily of Kent</i>	193	
<i>Lima</i>		227
<i>Limber Twig</i>	193	
LIMBERTWIG	193	
LIMBERTWIG (small or red)	193	
LIMBERTWIG (large or green)	194	
<i>Lincoln</i>		245
LINCOLN PIPPIN		122
<i>Lincoln Pippin</i> (synonym of Yellow Bellflower)	381	
<i>Lincoln Pippin</i> (synonym of Winthrop Greening)		245
<i>Lincolnshire Pippin</i>		96
LINPENWALD		122
<i>Linnæus Pippin</i>	381	
<i>Linnæus Pippin</i>	381	
<i>Lippincott</i>		215
<i>Litsey</i>	168	
<i>Little Pearmain</i>	89	
<i>Little Red Romanite</i> (synonym of Gilpin)	138	
<i>Little Red Romanite</i> (synonym of Romanite)	289	
<i>Little Ropka</i>	282	
<i>Little Seedling</i>	282	
<i>Liveland Raspberry</i>		129
<i>Livesley's Imperial</i>		126
<i>Livland Raspberry</i>		129
<i>Livre</i>	266	
<i>Loeb's Favorite</i>	361	
<i>Lodge's Early</i>		215
<i>Logan's Northern Pippin</i>	213	
<i>Long Bois</i>	181	
LONGFIELD	26	122
<i>Longfield's Apple</i>		123
<i>Long Island</i>		60
<i>Long Island Graft</i>		60
LONG ISLAND PEARMAN		124
LONG ISLAND PIPPIN	16	
LONG ISLAND RUSSET (I)	17, 194	
LONG ISLAND RUSSET (II)	195	
<i>Long Island Seek-No-Further</i>	130	
<i>Long John</i> (synonym of Kaighn)		113

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Long John</i> (synonym of Long Red Pearmain).....		125
<i>Long Pearmain</i> (synonym of Kaighn).....		113
<i>Long Pearmain</i> (synonym of Long Red Pearmain).....		125
<i>Long Pippin</i>	183	
LONG RED PEARMAIN		125
LONG STEM OF PENNSYLVANIA	196	
LONG SIEM		125
<i>Long Stem Sweet</i>		13
LONGWORTH		125
<i>Longworth Red Winter</i>		125
<i>Lopside</i>	153	
<i>Lop-sided Pearmain</i>	153	
<i>Lop-side Pearmain</i>	153	
<i>Lord Kingston</i>		96
<i>Lord Nelson</i>		21
LORD SUFFIELD		126
<i>Lothringer Rambour</i>		89
<i>Lothringer Rambour d'Etc.</i>		213
LOU		126
<i>Louis XIII</i>		55
LOUISE	20, 25	69, 126
<i>Louise, Princess</i>		127
LOWELL	16, 24	128
<i>Lowell Pippin</i>		128
LOWLAND RASPBERRY		129
LUBSK QUEEN		120
<i>Lubsk Reinette</i>		130
<i>Lucius Apfel</i>		21
<i>Lyman's Large Yellow</i>		171
<i>Lyman's Pumpkin Sweet</i>		171
LYSCOM		130
MABIE	24	131
<i>McAdow's June</i>		112
McAFEE	196	
<i>McAfee Red</i>	196	
<i>McAfee's Nonesuch</i>	196	
<i>McAfee's Nonesuch</i>	196	
<i>McAfee's Red</i>	196	
<i>McAfee</i>	197	
<i>McAfee's Nonesuch</i>	197	
Mc CARTY		132, 173
<i>McClellan</i>		135
<i>McClouds Family</i>	127	
MACDONOUGH		132
McINTOSH	17, 18, 20, 25	69, 132
<i>McIntosh Red</i>		133
<i>Mackie's Clyde Beauty</i>		36
<i>Mackinlay</i>	197	

COMBINED INDEX TO VARIETIES.

325

	VOL. I. PAGE.	VOL. II. PAGE.
McKINLEY	197	
McKINNEY	198	
McLELLAN		131
<i>McLellan</i>		135
<i>McLouds Family</i>	127	
<i>McMahan</i>		136
<i>McMahan White</i>		136
McMAHON		136
<i>McMahon White</i>		136
<i>Macomber</i>		187
<i>Magdalene</i>		142
MAGENTA	198	
<i>Magnum Bonum</i>		23
MAGOG		138
<i>Magog Red Streak</i>		138
MAIDEN BLUSH	17, 18	139
MAIDEN FAVORITE		141
<i>Maiden's Apple</i>		141
<i>Maiden's Blush</i> (synonym of Hawthornden).....		96
MAKELFIELD	312	
MALA CARLE	199	
<i>Malcarle</i>	199	
<i>Mal Carle</i>	199	
<i>Male Carle</i>	199	
MALINDA	199	
<i>Mailinokskoe</i>		177
<i>Mamma Bean</i>	67	
<i>Mamma Bean</i>	67	
<i>Mammoth</i> (synonym of Gloria Mundi).....		77
<i>Mammoth</i> (synonym of Golden Pippin).....		78
<i>Mammoth Black Twig</i> (synonym of Arkansas).....	47	
<i>Mammoth Black Twig</i> (synonym of Arkansas Black)....	50	
<i>Mammoth Black Twig</i> (synonym of Paragon).....	247	
<i>Mammoth Pippin</i>		77
<i>Mammoth Rambo</i>		89
MANCHESTER	200	
<i>Manks Codling</i>	379	
MANN	201	
<i>Maralandica</i>		51
MARUNGO (Crab)		259
<i>Marungo No. 1</i>		259
<i>Marungo Winter Crab</i>		259
MARGARET		141
<i>Margaret Early</i>		142
<i>Margaret Early Red</i>		142
<i>Margaretha Apfel</i>		142
<i>Marget Apple</i>		142

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Marguerite</i>		142
<i>Marietta Russet</i>	293	
<i>Marietta Seek-No-Further</i>	364	
MARIGOLD	202	
<i>Marigold</i> (synonym of Orange Pippin)		154
<i>Marigold Creed's</i>		154
<i>Marigold Pippin</i> (synonym of Marigold)	202	
<i>Marigold Pippin</i> (synonym of Orange Pippin)		154
<i>Marle Carle</i>	199	
<i>Maroquin</i>	264	
<i>Marrow-bone</i>		225
MARTHA (Crab)	3	259
<i>Martin</i>		135
<i>Marygold</i> (synonym of Marigold)	202	
<i>Marygold</i> (synonym of Orange Pippin)		154
MASON ORANGE	204	
<i>Mason's Improved</i>	204	
<i>Mason's Orange</i>	204	
<i>Massachusetts Golden Russett</i>	164	
MASTEN	204	
<i>Masten's Seedling</i>	204	
<i>Matchless</i>	225	
<i>Matthews Stripe</i>		131
<i>Maudlin</i>		142
<i>May</i>	153	
<i>May Apple</i>	153	
<i>May Seek-No-Farther</i>	153	
<i>May Seek-No-Further</i>	153	
<i>Marchem Sweet</i>		146
<i>Mela Carla</i>	199	
<i>Mela de Carlo</i>	199	
<i>Mela di Carlo</i>	199	
<i>Mela Januero</i>	94	
<i>Melinda</i>	199	
MELON	15, 24, 204	
<i>Melon</i> (synonym of Gloria Mundi)		77
<i>Melon Apple</i>	204	
<i>Melon de Norton</i>	204	
<i>Melon, Norton</i>	204	
<i>Melting Pippin</i>	244	
<i>Menage</i>	206	
MÉNAGÈRE	206	
<i>Menagerie</i>	206	
<i>Mere de Menage</i>	206	
<i>Merit</i>	88	
MERRILL	207	
<i>Merrill's</i>	207	

COMBINED INDEX TO VARIETIES.

327

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Merrill's Apple</i>	207	
<i>Metzgerapfel</i>	381	
<i>Metzger's Calvill</i>	381	
<i>Michel Miller</i>	158	
<i>Michigan Beauty</i>		201
MIDDLE	207	
MILAM	208	
MILDEN	209	
<i>Milding</i>	209	
<i>Millcreek</i> (synonym of Smokehouse).....	312	
<i>Millcreek</i> (synonym of Vandevere Pippin).....		231
<i>Millcreek Vandevere</i>	312	
<i>Millcreek Vandevere</i>		231
MILLER		142
<i>Miller</i> (synonym of Hiester).....	158	
<i>Miller Seedling</i>		142
MILLIGEN		142
MILWAUKEE	25, 211	
MINISTER	212	
<i>Minister Apple</i>	212	
MINKLER	18, 213	
MINNESOTA (Crab)		260
MISSING LINK	214	
<i>Mississippi</i>		77
<i>Missouri</i>	215	
<i>Missouri Janet</i>	271	
<i>Missouri Keeper</i>	215	
<i>Missouri Orange</i>	215	
MISSOURI PIPPIN	18, 215	
<i>Missouri Pippin</i> (synonym of Nickajack).....	227	
<i>Missouri Red</i>	227	
<i>Missouri Superior</i>	107	
<i>Mr. Gladstone</i>		75
<i>Mittle</i>	207	
<i>Mabbs</i>	227	
MONMOUTH	210	
<i>Monmouth Pippin</i>	217	
MONROE SWEET	24	
<i>Montgomery Sweet</i>		10
<i>Monstreuse Pippin</i>		77
<i>Monstrous Pippin</i>		77
<i>Monstrous Rambo</i>		89
<i>Montreal</i> (synonym of Montreal Beauty).....		260
<i>Montreal</i> (synonym of St. Lawrence).....		193
MONTREAL BEAUTY (Crab)		261
<i>Montreal Peach</i>		159
MOON	218	
<i>Moore's Late Sweet</i>	219	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Moore's Shanty</i>	219	
<i>Moore's Sweet</i>	219	
<i>Moore's Sweeting</i>	219	
MOORE SWEET	219	
<i>Moor's Sweeting</i>	219	
<i>Moose</i>		145
<i>Morgan's Favorite</i>		227
<i>Morris Red</i>	324	
MOSHER		143
<i>Mosher Sweet</i>		143
MOTHER	15, 24	143
<i>Mother Apple</i>		143
<i>Mother of America</i>		143
<i>Mountaineer</i>		145
<i>Mountain Flora</i>		77
<i>Mountain Pippin</i> (synonym of Fallawater)	125	
<i>Mountain Pippin</i> (synonym of Newtown Pippin)	146	
MOUNTAIN SWEET		145
MOUSE		145
MOYER	220	
<i>Moyer Prize</i>	220	
<i>Mudhole</i>		125
<i>Mumper Vandevere</i>	213	
MUNSON		146
<i>Munson Sweet</i>		146
<i>Munson Sweeting</i>		146
<i>Musgrove</i>		89
<i>Musgrove's Cooper</i> (synonym of Grosh)		89
<i>Musgrove's Cooper</i> (synonym of Western Beauty)		239
<i>Mushroom</i>		24
<i>Musk Spice</i>		64
<i>Mygatt's Bergamot</i>		49
<i>Myer's Nonpareil</i>		148
<i>Naliwi Jabloky</i>		239
<i>Naylor Rambo</i>		89
<i>Neige</i>		65
<i>Neige-Framboise de Gielen</i>		65
<i>Neisley's Winter</i>	255	
<i>Neisley's Winter Penick</i>	255	
NELSON	221	
<i>Nelson Sweet</i>	221	
<i>Ne Plus Ultra</i>	88	
NERO	222	
<i>Neustadt's gelber Pepping</i>	146	
<i>Never Fail</i>	271	
<i>Neverfail</i>	271	
NEWARK PIPPIN	223	
<i>Newark Sweeting</i>	91	

COMBINED INDEX TO VARIETIES.

329

	VOL. I. PAGE.	VOL. II. PAGE.
<i>New Brunswick</i>		151
<i>Newby</i>	107	
<i>New England Golden Russet</i>	164	
<i>New England Russet</i>	164	
<i>New England Seeknofurther</i>	364	
<i>New Greening</i>	331	
<i>V. J. Red Streak</i>		53
NEWMAN	24, 224	
<i>Newman Seedling</i>	224	
<i>New Missouri</i>	197	
<i>New Rhode Island Greening</i>	331	
<i>Newton's Pippin</i>	146	
<i>Newton Spitzenberg</i>	225	
<i>Newton Spitzenburgh</i>	225	
<i>Newton Yellow Pippin</i>	146	
<i>Newtown Greening</i>		83
NEWTOWN PIPPIN	11, 15, 16, 19, 146	
<i>Newtown Spitzenberg</i>	225	
<i>Newtown Spitzenbergh</i>	225	
NEWTOWN SPITZENBURG	15, 17, 225	
<i>Newtown Spitzenburgh</i>	225	
NEW WATER	226	
<i>New York Bellflower</i>		60
<i>New York Gloria Mundi</i>		77
<i>New York Greening</i>	146	
<i>New York Greening</i> (synonym of Golding)		83
<i>New York Pippin</i> (synonym of Ben Davis)	69	
<i>New York Pippin</i> (synonym of Newtown Pippin)	146	
<i>Niack Pippin</i> (synonym of Sweet Bough)		217
NICKAJACK	18, 227	
<i>Nickejack</i>	227	
<i>Nodhead</i>	170	
<i>Nonesuch</i> (synonym of Red Canada)	276	
<i>Nonesuch</i> (synonym of Hubbardston)	161	
<i>Nonpareil</i>		148
<i>Nonpareille</i>		184
<i>Nonpareille de Hubbardston</i>	161	
<i>Nonsuch</i> (synonym of Hubbardston)	161	
<i>Nonsuch</i> (synonym of McAfee)	197	
<i>Nonsuch</i> (synonym of Red Canada)	276	
<i>Normanton Wonder</i>	112	
<i>North American Best</i>		168
<i>Northampton</i>		21
<i>North Carolina</i>	227	
<i>Northern Golden Sweet</i>		147
<i>Northern Golden Sweeting</i>		147
NORTHERN SPY	12, 15, 17, 18, 21, 24, 37, 229	
NORTHERN SPY GROUP	24	

	VOL. I. PAGE.	VOL. II. PAGE.
NORTHERN SWEET		147
<i>Northern Sweeting</i> (synonym of Munson)		146
<i>Northern Sweeting</i> (synonym of Northern Sweet)		147
<i>North Star</i>		48
NORTHWESTERN Greening	24, 233	
<i>North West Greening</i>	233	
<i>Northwick Pippin</i>		21
<i>Norton's Melon</i>	204	
<i>Norton Watermelon</i>	204	
No. 3 M		121
No. 4 M		155
No. 6 M		83
No. 9 M		24
No. 11 M		189
No. 14 M		5
No. 18 M		5
No. 21 M		115
No. 22 M		22
No. 23 M		7
No. 24 M		195
No. 26 M		6
No. 54 M		87
No. 56 M		123
No. 57 M		123
No. 80 M		195
No. 94 M		114
No. 105 M		32
No. 134 M		224
No. 139 M		183
No. 140 M		45
No. 147 M		222
No. 152 M		27
No. 161 M		18
No. 161		123
No. 205		115
No. 206		45
No. 210		244
No. 224		6
No. 230		224
No. 236		6
No. 245		24
No. 252		7
No. 262		32
No. 288		177
No. 333		182
No. 340		129
No. 343		182
No. 351		170

COMBINED INDEX TO VARIETIES.

331

	VOL. I. PAGE.	VOL. II. PAGE.
No. 372		195
No. 410	282	
No. 418	282	
No. 442		247
No. 444		130
No. 453		14
No. 457		39
No. 467	61	
No. 469		83
No. 472		155
No. 477		18
No. 540		114
No. 587		57
No. 590		189
No. 964		10
No. 978		82
No. 979		82
No. 981		82
No. 12 Orel		7
No. 21 Veronesh		247
No. 51 Vor.		79
No. 84 Vor.		83
Nyack		30
Nyack Pippin		30
OAKLAND	234	
Oakland County Seek-No-Further	234	
Oats		51
OBLONG (Crab)		261
Oblong-Fruited Siberian Crab		261
Oblong Siberian Crab		261
OCCIDENT	24, 235	
Oel	236	
OEL AUSTIN	24, 236	
OLDENSBURG		147
Ohio Beauty (synonym of Western Beauty)		230
Ohio Beauty (synonym of Grosh)		89
Ohio Favorite	244	
OHIO NONPAREIL		148
Ohio Nonpareil (synonym of Gravenstein)		85
Ohio Nonpareil		148
OHIO PIPPIN		149
Ohio Wine (synonym of Fall Wine)		64
Ohio Wine (synonym of Kaighn)		113
OKABENA		150
Okabena		150
OLDENBURG	17, 18, 25, 26	
Oldenburg, Duchess of		151
OLDENBURG GROUP	25	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Oldenburgh</i>		151
<i>Old English Pearmain</i>	378	
<i>Old Golden Pippin</i>	141	
<i>Old Hawthorndean</i>		96
<i>Old Nonsuch</i>	276	
<i>Old Pearmain</i>	378	
OLIVE	237	
OLIVER	238	
<i>Oliver's Red</i>	238	
OLYMPIA	24, 60, 239	
<i>Olympia Baldwin</i>	239	
<i>Omensk</i>		189
ONTARIO	21, 24, 240	
OPALESCENT	242	
<i>Oporto</i>		7
ORANGE		153
ORANGE (Crab)		261
<i>Orange</i> (synonym of Fall Orange)		60
<i>Orange</i> (synonym of Lowell)		128
<i>Orange</i> (synonym of Orange Pippin)		154
<i>Orange Blenheim</i>		21
<i>Orange de Cox</i>		42
ORANGE OF NEW JERSEY		153
ORANGE OF PENNSYLVANIA		153
ORANGE PIPPIN		154
<i>Orange Pippin</i> (synonym of Marigold)	202	
<i>Orange Pippin</i> (synonym of Blenheim)		21
<i>Orange Russet</i>		154
ORANGE SWEET		154
<i>Orange Sweet</i> (synonym of Munson)		146
<i>Orange Sweet</i> (synonym of Golden Sweet)		81
<i>Orange Sweeting</i> (synonym of Golden Sweet)		81
<i>Orange Sweeting</i> (synonym of Orange Sweet)		154
<i>Orleans</i>	161	
<i>Orloff</i>		233
<i>Orlovskoe</i>		233
<i>Orlovskoe</i>		233
ORNAMENT	243	
<i>Ornement de Table</i>	243	
<i>Ornement de Table</i>	243	
<i>Orsimoi</i>		97
ORTLEY	244	
<i>Ortley Apple</i>	244	
<i>Ortley Pippin</i>	244	
<i>Osgood's Favorite</i>		131
<i>Osimoë</i>		97
<i>Oskaloosa</i>		106
OSTRAKOFF		155

COMBINED INDEX TO VARIETIES.

333

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Ostrakoff Glass</i>		155
<i>Ostrekoft</i>		155
<i>Ostrekoft's Glass</i>		155
<i>Ostrekovskaya Steklianika</i>		155
<i>Ostrekovskaja Steklianika</i>		155
<i>Ostrokoft</i>		155
<i>Ostrokoft's Glass</i>		155
<i>Ox Apple</i>		77
<i>Ox-Eye</i> (synonym of Buckingham)	88	
<i>Ox Eye</i> (synonym of Newtown Spitzenburg)	225	
<i>Oxeye</i> (synonym of Vandevere)		230
<i>Oxheart</i> (synonym of Cheeseboro)		33
<i>Oxheart</i> (synonym of Twenty Ounce Pippin)	349	
<i>Owen's Golden Beauty</i>		240
<i>Ozark</i>	135	
<i>Ozark Pippin</i>		45
PALOUSE		156
PALMER	246	
<i>Palmer Greening</i>	361	
<i>Palmer of N. Z.</i>	246	
<i>Paper</i>		30
<i>Paper-skin</i>		30
<i>Paradies Apfel</i>		85
<i>Paradise Winter</i>	380	
<i>Paradise Winter Sweet</i>	380	
PARAGON	24, 246	
<i>Paragon</i> (synonym of Arkansas)	47	
<i>Pariser Rambour Reinette</i>	94	
<i>Park</i> (synonym of McAfee)	197	
<i>Park</i> (synonym of Park Spice)	248	
<i>Park</i> (synonym of Long Red Pearmain)		123
<i>Poik Apple</i>	248	
<i>Parks Keeper</i>	197	
PARK SPICE	248	
PARLIN	248	
<i>Parlin's Beauty</i>	248	
<i>Parmain d'Angleterre</i>	378	
<i>Parmain d'Hiver</i>	378	
<i>Parmain-Pepping</i>	378	
PARRY WHITE		157
PARSON	249	
<i>Parson's Sweet</i>	249	
<i>Paternoster Apple</i>	114	
<i>Paterson's Sweet</i>	54	
PATTEN		158
<i>Patten's Duchess No. 3.</i>		158
<i>Patten Greening</i>		158
<i>Patterson's Sweet</i>	54	

	VOL. I. PAGE.	VOL. II. PAGE.
PAUL IMPERIAL (Crab).....		262
PAWPAW	250	
<i>Pawpaw Seedling</i>	250	
PAYNE	250	
<i>Payne Late Keeper</i>	250	
<i>Payne's Keeper</i>	250	
<i>Payton</i>	135	
PEACH	251	
PEACH (Montreal)		159
<i>Peach Apple of Montreal</i>		159
<i>Peach of Kentucky</i>	251	
<i>Peach of Montreal</i>		159
PEACH POND		160
<i>Peach Pond Sweet</i>		160
<i>Peach Pound Sweet</i>		160
<i>Pear Apple</i>		174
<i>Pear Lot</i>		244
PEARMAN	252	
<i>Pearmain</i> (synonym of Winter Pearmain).....	378	
<i>Pearmain</i> (synonym of Long Red Pearmain).....		125
<i>Pearmain d'Hiver</i>	378	
<i>Pearmain Herefordshire</i>	378	
PEARSALL		161
<i>Pearsall's Sweet</i>		161
<i>Pear-Tree Lot</i>		244
PEASE		161
<i>Pease, Waller</i>		161
PEASGOOD NONSUCH		162
<i>Peasgood's Nonesuch</i>		162
<i>Peau</i>	264	
<i>Peck</i>	253	
<i>Pecker</i>	56	
PECK Pleasant	12, 24, 32, 253	
<i>Peck's Pleasant</i>	253	
<i>Peewaukee</i>	258	
<i>Pelican</i>	255	
<i>Penick</i>	255	
<i>Pennick</i>	255	
PENNOCK	17, 255	
<i>Pennock's Red Winter</i>	255	
<i>Pennsylvania Cider</i>	311	
<i>Pennsylvania Red-Streak</i>	373	
<i>Pennsylvania Vandevere</i>		231
<i>Pipin de Kent</i>		15
<i>Pepin de New-York</i>	69	
<i>Pepin d'Or</i>	141	
<i>Pepin Parmain d'Angleterre</i>	378	
<i>Pepin Parmain d'Hiver</i>	378	

COMBINED INDEX TO VARIETIES.

335

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Pepin Ribston</i>		184
<i>Pepping Englischer</i>		57
<i>Peremeses</i>	379	
<i>Perle d'Angleterre</i>		21
<i>Permaine</i>	379	
<i>Permain</i>	379	
PERRY REDSTREAK		162
<i>Perry Red Streak</i>		162
PEERY RUSSET	256	
PETER	25	163
<i>Petersburgh Pippin</i>	146	
<i>Peterson's Charlamoff</i>		32
<i>Petit Api</i>	181	
<i>Petit Api Rose</i>	181	
<i>Petit Api Rouge</i>	181	
<i>Petit Apis</i>	181	
<i>Petite Reine, & Grise</i>	90	
<i>Petrozskoe</i>		195
<i>Petrozskoe</i>		195
PEWAWKEE	25, 258	
<i>Pfeiffer</i>	202	
<i>Pfeiffer</i>	262	
<i>Pfund</i>	206	
<i>Philadelphia Pippin</i> (synonym of Fall Pippin)		62
<i>Philadelphia Pippin</i> (synonym of White Spanish Reinette)		241
<i>Philadelphia Sweet</i>		10
<i>Phillip Rick</i>	172	
<i>Phoenix</i> (synonym of Pennock)	255	
<i>Phoenix</i> (synonym of Rome)	290	
<i>Phanix</i>		3
<i>Phönix</i>		3
<i>Picard</i>	260	
<i>Picard's Reserve</i>	260	
<i>Pickard</i>	260	
PICKARD RESERVE	260	
<i>Pickard's Reserve</i>	260	
<i>Pickaway Rambo</i>		84
PICTA STRIATA (Crab)		263
<i>Pie Apple</i> (synonym of Holland Pippin)		101
<i>Pie Apple</i> (synonym of Sops of Wine)		203
<i>Pieta</i>		263
<i>Pieta Striata</i>		263
PIFER	262	
<i>Pilliken</i>	153	
<i>Pim's Beauty of the West</i>	125	
<i>Pineapple</i>	257	
<i>Pineapple Russet</i>	257	
<i>Pine's Beauty of the West</i>	125	

	VOL. I. PAGE.	VOL. II. PAGE.
PINE STUMP	263	
<i>Piper</i>	262	
PIPPIN	263	
<i>Pippin</i> (synonym of Newtown Pippin).....	146	
<i>Pippin, Bullock</i>	89	
<i>Pippin Kent</i>		15
<i>Pittstown Pippin</i>		79
<i>Pitzer Hill</i>	269	
<i>Platarchium</i>	379	
<i>Plodowitka Caudkaja</i>		170
<i>Plodowitka Cuadkaja</i>		170
PLUMB CIDER		164
<i>Plum Cider</i>		164
<i>Pointed Pipka</i>		32
<i>Polecat</i>		212
<i>Polhemus</i>	219	
<i>Pomme Astrachan</i>		239
<i>Pomme d'Api</i>	181	
<i>Pomme d'Api Rouge</i>	181	
<i>Pomme d'Apis</i>	181	
<i>Pomme d'Astrachan</i>		239
<i>Pomme de Blenheim</i>		21
<i>Pomme de Caen</i>	94	
<i>Pomme de Charles</i>	199	
<i>Pomme de Cuir</i>	269	
<i>Pomme d'Ete of Canada</i>		51
<i>Pomme de Fameuse</i>		65
<i>Pomme de Fer</i>	292	
<i>Pomme de Hawthornden</i>		96
<i>Pomme de Laak</i>	114	
<i>Pomme de Neige</i>		65
<i>Pomme de Notre-Dame</i>		213
<i>Pomme d'Orange</i>		154
<i>Pomme de Transylvania</i>	64	
<i>Pomme Fameuse</i>		65
<i>Pomme Finale</i>	199	
<i>Pomme Graefenstein</i>		85
<i>Pomme Grece</i>	264	
<i>Pomme Gris</i>	264	
POMME GRISE	264	
<i>Pomme Grise d'Or</i>	328	
<i>Pomme Groseille</i>		253
<i>Pomme Peche</i>		159
<i>Pomme Rose</i>	181	
<i>Pommeroy</i>	184	
<i>Pomme Royal</i>		49
<i>Pomme Royale</i>		49
<i>Pomme Roye</i> (synonym of Pennock).....	255	

COMBINED INDEX TO VARIETIES.

337

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Pomme Royale</i> (synonym of Dyer).....		49
<i>Pomme Water</i>		49
<i>Pommeswater</i> in Illinois.....		49
POMONA		164
<i>Pomona Britannica</i>		3
<i>Pomone d'Apis</i>	181	
<i>Pompey</i>		233
<i>Poplar Bluff</i>	311	
<i>Popular Bluff</i>	311	
PORTER	12, 15	166
<i>Portugal</i>	94	
<i>Possaris Nalicia</i>		6
<i>Potter's Large</i>		116
<i>Potter's Large Grey Seedling</i>		116
<i>Potter's Large Seedling</i>		116
<i>Poughkeepsie Russet</i> (synonym of English Russet)	118	
<i>Poughkeepsie Russet</i> (synonym of Perry Russet)	257	
<i>Pound</i> (synonym of Fallawater)	125	
<i>Pound</i> (synonym of Nickajack)	227	
<i>Pound</i> (synonym of Gloria Mundi)		77
<i>Pound Pippin</i>		62
<i>Pound Royal</i> (synonym of Fall Pippin)		62
<i>Pound Royal</i> (synonym of Golden Pippin)		78
<i>Pound Royal</i> (synonym of Lowell)		128
<i>Pound Royal</i> (synonym of Long Red Pearmain)		125
<i>Pound's July</i>		217
POUND SWEET	266	167
<i>Pound Sweet</i> (synonym of Moore Sweet)	219	
<i>Pound Sweet</i> (synonym of Pumpkin Sweet)		171
<i>Powers</i>		168
<i>Prager Reinette Franche de Grandville</i>	264	
<i>Pratt</i>	266	
PRATT SWEET	266	
<i>President Napoleon</i>		3
<i>Pride of Genesee</i>		73
<i>Pride of Hudson</i>	62	
<i>Pride of Texas</i>	337	
<i>Pride of the Hudson</i>	62	
PRIESTLY	267	
<i>Priestley</i>	267	
<i>Priestley's American</i>	267	
PRIMATE		167
<i>Primiting</i>		240
PRINCE ALBERT	268	
<i>Prince Albert (Lane)</i>	268	
<i>Prince Bismark</i>		19
<i>Princesse Noble</i>	142	
<i>Prince of Wales</i>		21

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Prince's Early Harvest</i>		51
<i>Prince's Harvest</i>		51
<i>Prince's Large Pippin of New York</i>		62
<i>Prince's Large Red and Green Sweeting</i>		177
<i>Prince's Red and Green Sweet</i>		177
<i>Prince's Yellow Harvest</i>		51
<i>Princess Louise</i>		127
<i>Princessin-Apfel</i>		85
<i>Prior's Late Red</i>	269	
<i>Prior's Red</i>	269	
PROLIFIC BEAUTY	16	
<i>Prolific Beauty</i> (synonym of Pennock)	255	
<i>Prolific Beauty</i> (synonym of Blue Pearmain)	80	
<i>Prolific Sweet</i>		170
PROLIFIC SWEETING		169
<i>Prussian</i>		38
PRYOR	269	
<i>Pryor Red</i>	269	
<i>Pryor's Pearmain</i>	269	
<i>Pryor's Red</i>	269	
PUMPKIN RUSETT		170
PUMPKIN SWEET	16, 17, 18	171
<i>Pumpkin Sweet</i> (synonym of Cheeseboro)		33
<i>Pumpkin Sweet</i> (synonym of Pumpkin Russet)		170
<i>Pumpkin Sweeting</i> (synonym of Pumpkin Sweet)		172
<i>Pumpkin Sweeting</i> (synonym of Pumpkin Russet)		170
<i>Putman's Russet</i>	293	
<i>Putnam Russet</i>	293	
<i>Putnam's Sawetwell</i>	299	
<i>Pyrus Astracanicus</i>		230
QUAKER (Crab)		263
<i>Quaker</i> (synonym of Streaked Pippin)	321	
<i>Queen</i> (synonym of Buckingham)	88	
<i>Queen</i> (synonym of Summer Queen)		212
<i>Queen</i> (synonym of Williams)		242
<i>Queen Anne</i> (synonym of Lowell)		128
<i>Queen Anne</i> (synonym of Mother)		143
QUEEN CHOKE (Crab)		264
<i>Queen of Haywood</i>	155	
<i>Queens</i>	84	
<i>Queen's Choice</i>		263
QUINCE (of Cole)		174
QUINCE (of Cox)	17	174
<i>Quince Apple</i>		174
<i>Rag Apple</i>		146
RALLS	18, 24, 270	
RALLS GROUP	24	
<i>Ralls Genet</i>	271	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Ralls Janet</i>	271	
RAMBO	17, 18, 32, 273	
<i>Rambouillet</i>	273	
<i>Ramboulette</i>	273	
<i>Ramboulrette</i>	109	
<i>Rambour</i>		213
<i>Rambour Barre</i>	379	
<i>Rambour Blanc</i>		213
<i>Rambour d'Amerique</i>		213
<i>Rambour d'Ete</i>		213
<i>Rambour Franc</i>		213
<i>Rambourg Aigre</i>		213
<i>Rambour Gros</i>		213
<i>Rambour Lorraine</i>		89
<i>Rambour Raye</i>		213
<i>Rambu</i>		213
<i>Ramsdale's Sweeting</i>		175
<i>Ramsdell</i>		175
<i>Ramsdell Red Sweet</i>		175
<i>Ramsdell's Red</i>		175
<i>Ramsdell's Red Sweeting</i>		175
RAMSDALL SWEET		175
<i>Ramsdell's Red Pumpkin Sweet</i>		175
<i>Randall's Red Winter</i>		175
RASPBERRY		176
<i>Raule Janet</i>	271	
<i>Raule's Genet</i>	271	
<i>Raule's Janet</i>	271	
<i>Raule's Janette</i>	271	
<i>Raule's Jannetting</i>	271	
<i>Raule's Jennetting</i>	271	
<i>Raul's Gennetting</i>	271	
<i>Razole's Genet</i>	271	
<i>Razole's Janet</i>	271	
<i>Razole's Jennet</i>	271	
<i>Razole's Jennetting</i>	271	
<i>Razole's Jennette</i>	271	
<i>Rox Apple</i>		146
<i>Reagan</i>	135	
RED AND GREEN SWEET		177
<i>Red and Green Sweeting</i>		177
<i>Red Astracan</i>		178
RED ASTRACHAN	15, 17, 18, 25, 26	178
<i>Red Astokhan</i>		178
<i>Red Baldwin Pippin</i>	56	
<i>Red Beilgheimer</i>		17
<i>Red Bellflower</i> (synonym of Ohio Nonpareil)		148
<i>Red Bellflower</i> (synonym of Long Red Pearmain)		125

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Red Ben Davis</i>	135	
<i>Red Bietigheimer</i>		19
<i>Red Bough</i>		177
RED CANADA 15, 24, 37,	275	
<i>Red Canada</i> of Ontario	292	
<i>Red Cathead</i>	267	
<i>Red Cheek</i> (synonym of Monmouth)	217	
<i>Red Cheek</i> (synonym of Raspberry)		177
<i>Red Cheek</i> (synonym of Maiden Blush)		139
<i>Red Cheek</i> (synonym of Fall Orange)		60
<i>Red Checked Pippin</i>	217	
<i>Red Cheek Pippin</i>	217	
<i>Red Codlin</i>	132	
<i>Red Doctor</i>	107	
<i>Red Eiser</i>	116	
<i>Red Everlasting</i>	153	
<i>Red Gilliflower</i> (synonym of Black Gilliflower)	77	
<i>Red Gilliflower</i> (synonym of Scollop Gilliflower)		197
<i>Red Gilliflower</i> (synonym of Striped Gilliflower)		207
<i>Red Gloria Mundi</i>	88	
RED GRAVENSTEIN		180
<i>Red Gravenstein</i> (synonym of Banks)		14
<i>Red Hawthornden</i>		96
<i>Red Hazel</i>	227	
RED HOOK		180
<i>Red Horse</i>	88	
<i>Red Joaneling</i>		142
RED JUNE		181
<i>Red Juneating</i> (synonym of Margaret)		142
<i>Red Juneating</i> (synonym of Red June)		181
<i>Red Juneating</i> (synonym of Early Strawberry)		55
<i>Red June, Carolina</i>		181
<i>Red June of South</i>		142
<i>Red Juneling</i>		142
<i>Red Lady Finger</i>	183	
<i>Red Limbertwig</i>	193	
<i>Redling</i>	101	
<i>Red Neverfail</i>	271	
<i>Red Ox</i>	255	
<i>Red Pearmain</i> (synonym of Kaighn)		113
<i>Red Pearmain</i> (synonym of Long Red Pearmain)		125
<i>Red Pennock</i>	255	
<i>Red Phoenix</i> (synonym of Kaighn)		113
<i>Red Phoenix</i> (synonym of Long Red Pearmain)		125
<i>Red Pippin</i> (synonym of Ben Davis)	69	
<i>Red Pippin</i> (synonym of Nickajack)	227	
<i>Red Pippin</i> (synonym of Streaked Pippin)	321	
<i>Red Pippin</i> (synonym of Kaighn)		113

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Red Pippin</i> (synonym of Long Red Pearmain)		125
<i>Red Pound</i>	64	
<i>Red Pumpkin Sweet</i>		175
<i>Red Pumpkin Sweeting</i>		175
<i>Red Reinette</i>		83
<i>Red Rock</i>	188	
<i>Red Romanite</i>	289	
<i>Red Romanite of Ohio</i>	138	
RED RUSSET	278	
<i>Red Russet</i> (synonym of Pryor)	269	
<i>Red Shropshire</i>		203
RED SIBERIAN (Crab)		264
<i>Red Spitzenberg</i> (synonym of Kaighn)		113
<i>Red Spitzenberg</i> (synonym of Long Red Pearmain)		125
<i>Red Spitzenberg</i> (synonym of Richard Graft)		186
<i>Red Spitzenburg</i>		113
REDSTREAK	278	
<i>Red Streak</i>	278	
<i>Red Sweet Pippin</i>	219	
<i>Red Sweet Winsap</i>	333	
RED TRANSPARENT		182
<i>Red Transparent</i> (synonym of Vineuse Rouge)		233
RED TYPE OF TWENTY OUNCE		229
RED TYPE OF WESTFIELD <i>Seck-No-Further</i>	366	
<i>Red Vanderveere</i>		231
<i>Red Vanderveere</i>	312	
<i>Red Warrior</i>	247	
RED WINE		182
<i>Red Winter</i>	276	
RED WINTER PEARMAN	183	
<i>Red Winter Pearmain</i>	279	
<i>Red Winter Pearmain</i> (synonym of Westfield <i>Seck-No-Further</i>)	364	
<i>Red Winter Pearmain</i> (synonym of Kaighn)		113
<i>Red Winter Pearmain</i> (synonym of Long Red Pearmain) ..		125
RED WINTER PIPPIN	183	
<i>Red Winter Sweet</i>	219	
RIED		183
<i>Reindell's Large</i>		175
REINETTE	279	
<i>Reinette a feuille d'Acuba</i>	52	
<i>Reinette A Gobelet</i>		241
<i>Reinette Belle de Boskoop</i>		25
<i>Reinette Blanche</i>		241
<i>Reinette Blanche d'Espagne</i> (synonym of White Spanish Reinette)		241
<i>Reinette Blanche d'Espagne</i> (synonym of Fall Pippin)		62
<i>Reinette Canada</i>	94	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Reinette Carpentin</i>	96	
<i>Reinette de Blenheim</i>		21
<i>Reinette de Caen</i>	94	
<i>Reinette de Canada</i>	94	
<i>Reinette de Canada d'Cotes</i>	94	
<i>Reinette de Caux</i>	114	
<i>Reinette de Darnetal</i>	264	
<i>Reinette d'Espagne</i>		241
<i>Reinette d'Holland</i>		161
<i>Reinette de Landsberg</i>		119
<i>Reinette de New York</i>	146	
<i>Reinette d'Or</i>	114	
<i>Reinette de Traver</i>		184
<i>Reinette Doree</i>	114	
<i>Reinette du Canada</i>	94	
<i>Reinette du Canada d'Cortes</i>	94	
<i>Reinette du Canada Blanche</i>	94	
<i>Reinette Grenade Anglaise</i>		184
<i>Reinette Grise</i>	264	
<i>Reinette Grise Brownles</i>	86	
<i>Reinette Grise de Darnetal</i>	264	
<i>Reinette Grise Double</i>	265	
<i>Reinette Grise de Grandville</i>	265	
<i>Reinette Grise d'Ilizer</i>	265	
<i>Reinette Grise Extra</i>	265	
<i>Reinette Grise Franchaise</i>	264	
<i>Reinette Grosse de Angleterre</i>	94	
<i>Reinette Landsberger</i>		119
<i>Reinette Liubski</i>		130
<i>Reinette Monstrueuse</i>		25
<i>Reinette Monstreuse de Canada</i>	94	
<i>Reinette Orange de Cox</i>		42
<i>Reinette Pepin</i>	280	
REINETTE PIPPIN	280	
<i>Reinette Quarrendon</i>		142
<i>Reinette Tendre</i>		241
<i>Reinette toute Grise</i>	265	
<i>Reinette von Canada</i>	94	
<i>Reinette von Montfort</i>		25
<i>Reipka Melenkaya</i>	282	
<i>Renet Liubskui</i>		130
REPKA		183
REPKA MALENKA	281	
<i>Remboure d'Ete</i>		213
<i>Reschestwenskoe</i>		19
<i>Revelstone</i>		248
<i>Revelstone Pippin</i>		233
<i>Rhode Island</i>	282	

COMBINED INDEX TO VARIETIES.

343

	VOL. I. PAGE.	VOL. II. PAGE.
RHODE ISLAND Greening	12, 15, 16, 17, 18, 19, 24, 282	
RHODE ISLAND Greening Group.....	24	
<i>Rhode Island Russet</i>	257	
<i>Rhode Island Seek-No-Further</i>	130	
<i>Rhode Island Sweet</i>		172
<i>Riabinouka</i>		39
<i>Ribbed Gilliflower</i>		197
<i>Ribbed Pippin</i>		83
RIBSTON	17	184
<i>Ribstone</i>		184
<i>Ribstone Pippin</i>		184
<i>Ribston Pepping</i>		184
<i>Ribston Pippin</i>		184
<i>Richard</i>		186
RICHARD GRAFT		186
<i>Richardson's Red</i>	66	
<i>Richfield Nonsuch</i>	276	
<i>Rickman's Red</i>	227	
RIDGE	287	
<i>Ridge</i> (synonym of Ribston).....		184
<i>Ridge Pippin</i>	287	
<i>Riepka</i>		183
<i>Ripp Apfel</i>		85
<i>Roa You</i>	184	
<i>Robinson</i>	316	
<i>Rock</i>	289	
<i>Rock Apple</i>	188	
<i>Rockhill's Russet</i>		184
ROCK PIPPIN	288	
<i>Rock Romain</i>	271	
<i>Rock Rimmon</i>	271	
<i>Rode Wyn Appel</i>		203
ROLFE		187
<i>Rolland</i>	67	
<i>Roman Beauty</i>	290	
ROMANITE	289	
<i>Romanite</i> (synonym of Gilpin).....	138	
<i>Romanite</i> (synonym of Greyhouse).....	153	
<i>Romanite</i> (synonym of Pennock).....	255	
<i>Romanite</i> (synonym of Rambo).....	273	
<i>Romanite</i> of the South.....	289	
<i>Romanite</i> of the West.....	138	
<i>Romanite, South</i>	289	
<i>Roman Knight</i>	255	
ROMAN STEM	17, 18	188
ROME	32, 290	
<i>Rome Beauty</i>	290	
<i>Romenskoe</i>		189

	VOL. I. PAGE.	VOL. II. PAGE.
ROMNA		189
<i>Romna</i> (synonym of <i>Hibernal</i>).....		97
<i>Romnenskoe</i>		189
RONK		190
<i>Roschdestvenskoe</i>		19
<i>Roschdestvenskoe</i>		19
ROSEAU	292	
ROSE-COLORED LADY APPLE.....	182	
ROSE RED		190
<i>Rose Sweet</i>	333	
<i>Roschdestvenskoe</i>		19
<i>Rosseau</i>	292	
<i>Ross Greening</i>	352	
<i>Rother Astrakhan</i>		178
<i>Rother Eiser</i>	116	
<i>Rother Eiser Apfel</i>	116	
<i>Rother Gracensteiner</i>		180
<i>Rother Jacobs</i>		142
<i>Rother Jacobs Apfel</i>		142
<i>Rother Weinapfel</i>		182
<i>Rotherwein Apfel</i>		182
<i>Rouge de Pryor</i>	269	
<i>Rouge Rayee</i> (synonym of <i>Eiser</i>).....	116	
<i>Rouge Rayee</i> (synonym of <i>Redstreak</i>).....	278	
<i>Rough and Ready</i>		168
<i>Round Catshead</i>		29
<i>Round Secret</i>		172
<i>Round Top</i>		56
<i>Rox</i>	294	
ROXBURY	15, 16, 17, 18, 23,	293
<i>Roxbury Russet</i>	293	
<i>Roxbury Russetting</i>	293	
<i>Rox Russet</i>	294	
<i>Royal Janelle</i>	271	
<i>Royal Red of Kentucky</i>	375	
<i>Rubets Vinogradnui</i>		244
<i>Rubetz Vinogradnui</i>		244
<i>Rubicon</i>	250	
<i>Ruckman</i>	227	
<i>Ruckman's Red</i>	227	
<i>Russam</i> (synonym of <i>Kaighn</i>).....		113
<i>Russam</i> (synonym of <i>Long Red Pearmain</i>).....		125
<i>Russet</i>	294	
RUSSET BALDWIN	59	
<i>Russet, Boston or Roxbury</i>	294	
<i>Russet, Golden</i>	143	
<i>Russet Pearmain</i>	164	
<i>Russet Seek-No-Further</i>	364	

COMBINED INDEX TO VARIETIES.

345

	VOL. I. PAGE.	VOL. II. PAGE.
RUSSIAN BALDWIN	205	
<i>Russian Emperor</i>		3
<i>Russian Gauenstein</i>		6
<i>Rusty Core</i>		148
RUBINE	206	
<i>Sabine</i>		85
<i>Safstaholm</i>		191
SAFSTAHOLMS		191
<i>Säfstaholmsäple</i>		191
SAILEE Russet		192
<i>Saile Sweet</i>		177
SAILLY AUTUMN		192
<i>Saint-Germaine</i>		241
<i>St. Helena Russet</i>	94	
<i>St. Hilare</i>		98
<i>St. John's Strawberry</i>		55
ST. LAWRENCE	17, 18	192
<i>Saint-Laurence</i>		193
<i>Saint Laurent</i>		193
ST. PETER		194
SALISBURY	297	
<i>Salisbury Pippin</i>	297	
<i>Salisbury Winter</i>	297	
SANDY	24, 207	
SANDY CLASS		195
<i>Sanguineus</i>		65
<i>Sapsin</i>		203
<i>Sapsoneine</i>		203
<i>Sassaparilla Sweet</i>		94
SAWELL	209	
SAXTON		196
<i>Scalloped Gilliflower</i>		197
<i>Scalloped Gilliflower</i>		197
SCARLET BEAUTY	24	
SCARLET CRANBERRY	209	
<i>Scarlet Pearmain</i> (synonym of Kaighn).....		113
<i>Scarlet Pearmain</i> (synonym of Long Red Pearmain).....		125
SCARLET PIPPIN	25	196
<i>Scharmaré</i>		29
<i>Scharlottenthaler Golba</i>		222
SCHODACK	300	
<i>Schöner von Boskoop</i>		25
<i>Schoonacker</i>	301	
<i>Schoone van Boskoop</i>		25
SCHOONMAKER	301	
SCHUYLER SWEET		197
<i>Schwere Apfel</i>	326	
SCOLLOP GILLIFLOWER		197

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Scollop Gilliflower</i> (synonym of Striped Gilliflower).....		207
<i>Scolloped Gilliflower</i>		197
SCOTT	301	
<i>Scott</i> (synonym of Baker).....	55	
<i>Scott</i> (synonym of Primate).....		168
SCOTT BEST		198
<i>Scott's Red Winter</i>	302	
<i>Scott's Winter</i>	302	
SCRIBNER	303	
<i>Scribner's Spitzenberg</i>	303	
<i>Scribner's Spitzenburgh</i>	303	
<i>Scudamore's Crab</i>	278	
<i>Scudamous Crab</i>	278	
<i>Seek-No-Farther</i>	273	
<i>Seeknofurther</i>	364	
<i>Seek-No-Further</i> (synonym of Rambo).....	273	
<i>Seek-No-Further</i>	303	
<i>Seek-No-Further</i> of Del., N. J., and Penn.....	273	
<i>Seek-No-Further</i> (synonym of Cooper).....		40
<i>Seek-No-Further</i> (synonym of Green Seek-No-Further).....		86
<i>Seever's Red Streak</i>		230
<i>Senator</i>	238	
SENECA FAVORITE		198
<i>Seneca Favorite</i> (synonym of Quince [of Coxel]).....		174
<i>Seneca Spice</i>		174
SEPTEMBER (Crab)		264
SHACKLEFORD	304	
<i>Shackleford's Best</i>	304	
<i>Shaker's Yellow</i>		53
SHANNON (I)	305	
SHANNON (II)	306	
<i>Shannon</i> (synonym of Ohio Pippin).....		149
<i>Shannon Pippin</i>	305	
SHARP		199
<i>Sharpe's Early</i>		212
<i>Sharpe's Spice</i>		64
SHEDDAN	306	
<i>Sheepnose</i> (synonym of Bullock).....	89	
<i>Sheepnose</i> (synonym of Lady Finger).....	183	
<i>Sheepnose</i>	307	
<i>Sheepnose</i> (synonym of Egg Top).....		56
<i>Sheepnose</i> (synonym of Long Red Pearmain).....		125
<i>Sheep Shire</i>	131	
<i>Sheep's Nose</i>	89	
<i>Sheep's Snout</i>	89	
SHERRIFF	308	
SHERMAN		200
<i>Sherman's Favorite</i>		200

COMBINED INDEX TO VARIETIES.

347

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Sherman's Sweet</i>		200
<i>Sherwood's Favorite</i>		33
<i>Shiawasse Beauty</i>		201
SHIAWASSEE	25	69, 201
<i>Shiawassce Beauty</i>		201
<i>Shiawassie Beauty</i>		201
<i>Shippens Russet</i>	294	
SHIRLEY	308	
<i>Shoreditch White</i>		95
<i>Shropshirevine</i>		203
<i>Siberian August</i>		112
<i>Siberian Crab</i>	17	
<i>Siberian Crab</i> (synonym of Red Siberian)		264
<i>Siberian Crab</i> (synonym of Yellow Siberian)		264
SINE-QUA-NON		202
SKANK	309	
<i>Skanka</i>	81	
<i>Skanka Bogdanoff</i>	81	
<i>Skank</i>	321	
<i>Skeosnoi Krasnoi</i>		182
<i>Skeosnoi Krasnoi</i>		182
<i>Skeosnoi Schotol</i>		248
<i>Sleeper's Yellow</i>		53
SLEIGHT	310	
<i>Sleight's Lady Apple</i>	310	
<i>Slight's Lady Apple</i>	310	
SLINGERLAND		202
<i>Slingerland's Fall Pippin</i>		202
<i>Slingerland's Pippin</i>		202
<i>Small Admirable</i>		1
<i>Small's Admirable</i>		1
SMITH CIDER	18, 310	
<i>Smithfield Spice</i>		49
<i>Smith's</i>	311	
<i>Smith's Cider</i>	311	
SMOKEHOUSE	15, 312	
<i>Smoke House</i>	312	
<i>Smokehouse</i> (synonym of Vandevere Pippin)		231
<i>Smyrna</i>		33
<i>Snorter</i>	197	
<i>Snow</i>		65
<i>Sol Carter</i>	88	
<i>Solotoc Renet</i>		79
SOMERSET (N. Y.)		203
<i>Sondergleichen von Hubbardston</i>	161	
<i>Sops in Wine</i>		203
SOPS OF WINE		203
SOULARD (Crab)		295

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Sourbough</i> (synonym of Champlain)		30
SOUB BOUGH		204
<i>Sour Bough</i> (synonym of Champlain)		30
<i>Sour Bough</i> (synonym of Tart Bough)		220
<i>Sour Harvest</i>		168
<i>Southern Romanite</i>	289	
<i>Speckled</i>		65
SPECTATOR		204
<i>Spiced Oxeye</i>		231
<i>Spiced Ox Eye</i>	225	
SPICE SWEET	12	
<i>Spitzenberg</i>	121	
<i>Spitzenberg Newtown</i>	225	
<i>Spitzenburg</i>	314	
<i>Spitzenburgh</i> (synonym of Esopus <i>Spitzenburg</i>)	121	
<i>Spitzenburgh</i> (synonym of Newtown <i>Spitzenburg</i>)	225	
<i>Spitzenburgh Newtown</i>	225	
SPRING PIPPIN	314	
<i>Springport</i>	314	
<i>Springport Pippin</i>	314	
<i>Spy</i>	229, 315	
<i>Staalclubs</i> (synonym of Newtown <i>Spitzenburg</i>)	225	
<i>Staalclubs</i> (synonym of Vandevere Pippin)		231
<i>Staalclubs</i>		230
<i>Stalcubs</i>		230
STANARD	315	
<i>Stanard's Seedling</i>	315	
<i>Stannard</i>	315	
<i>Stannard's Seedling</i>	315	
STARKE	316	
STARKEY		205
<i>Starke Apple</i>	316	
STAR LADY APPLE	182	
STARR		206
<i>Stayman</i>	318	
<i>Stayman's Winesap</i>	318	
STAYMAN WINESAP	318	
<i>Steele's Red</i>	324	
<i>Steele's Red Winter</i> (synonym of Baldwin)	56	
<i>Steele's Red Winter</i> (synonym of Red Canada)	276	
<i>Steele's Red Winter</i> (synonym of Sutton)	324	
<i>Steel's Red</i>	276	
<i>Stekly</i>	158	
<i>Steklianka Bogdanoff</i>	81	
<i>Steklianka pesotchnaya</i>		195
STERLING	319	
<i>Sterling Beauty</i>	45, 319	
<i>Stettin Pippin</i>	114	

COMBINED INDEX TO VARIETIES.

349

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Stevenson Pippin</i>	197	
<i>Stewart's Nonpareil</i>		112
<i>Stillman</i>		207
STILLMAN EARLY		207
<i>Stillman's Early</i>		207
<i>Stine</i>	197	
<i>Stoke Tulip</i>		3
STONE	24, 320	
<i>Stone</i> (synonym of American Pippin)	45	
<i>Stor Casseler Reinnet</i>	114	
<i>Stor Kasselrenett</i>	114	
<i>Storr's Wine</i>	197	
STOWE	321	
<i>Stowe's Winter</i>	321	
<i>Straat</i>		208
<i>Strawberry</i> (synonym of Sops of Wine)		203
<i>Strawberry</i> (synonym of Chenango)		33
<i>Strawberry</i> (synonym of Late Strawberry)		120
<i>Strawberry</i> (synonym of Richard Graft)		186
<i>Stracherry</i>		207
STREAKED PIPPIN	321	
<i>Striped Ashmore</i>		231
<i>Striped Bellflower</i>		207
STRIPE FAMIFUSE		68
STRIPE GILLIFLOWER	16	207
<i>Striped Harvest</i>		93
<i>Striped Juneating</i>		142
<i>Striped Pearmain</i> (synonym of McAfee)	197	
<i>Striped Pearmain</i> (synonym of Long Red Pearmain)		125
<i>Striped Quarrendon</i>		142
<i>Striped Rambo</i>	273	
<i>Striped Red June</i>		182
<i>Striped Rhode Island Greening</i> (synonym of Domine)	109	
<i>Striped Rhode Island Greening</i> (synonym of Wells)	363	
<i>Striped Shropshire</i>		55
STRIPE SWEETING	16	
<i>Striped Sweet Pippin</i>	197	
<i>Striped Vanderveere</i>		231
<i>Striped Winter Pearmain</i>	197	
STROAT		208
<i>Strode</i>		208
STRODE BIRMINGHAM		208
<i>Strode's</i>		208
<i>Strode's Birmingham</i>		208
<i>Strohmer</i>		85
<i>Stromling</i>		85
<i>Stuart</i>	323	
STUART GOLDEN	323	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Stuart's Golden</i>	323	
STUMP		200
<i>Stump</i> (synonym of <i>Stuart Golden</i>)	323	
<i>Stymer's</i>		210
STYMUS		210
SUFFOLK BEAUTY		210
<i>Sugar Barbel</i>	61	
<i>Suislepper</i>		219
<i>Summer Bellefleur</i>		211
SUMMER BELLFLOWER		211
<i>Summer Bellflower</i> (synonym of <i>Autumn Bough</i>)		10
<i>Summer Hagloe</i>		92
<i>Summerour</i>	227	
SUMMER PEARMAIN	15	211
SUMMER PIPPIN	11	
<i>Summer Pippin</i> (synonym of <i>Champlain</i>)		30
<i>Summer Pippin</i> (synonym of <i>Fall Pippin</i>)		62
<i>Summer Pippin</i> (synonym of <i>Holland Pippin</i>)		101
SUMMER QUEEN	16	212
SUMMER RAMEO		213
<i>Summer Rambo</i> (synonym of <i>Grosh</i>)		89
<i>Summer Rambour</i>		213
SUMMER REDSTREAK		214
SUMMER ROSE	15	215
<i>Summer Russet</i>		218
SUMMER SPITZENBURG		215
<i>Summer Spitzenburgh</i>		215
SUMMER SWEET		216
<i>Summer Sweet</i> (synonym of <i>Hightop Sweet</i>)		68
<i>Summer Sweetling</i> (synonym of <i>Summer Sweet</i>)		216
<i>Summer Sweetling</i> (synonym of <i>Hightop Sweet</i>)		68
<i>Summer Traveller</i>		142
<i>Superior White</i>		157
<i>Sussex Scarlet Pearmain</i>	379	
<i>Susy Clark</i>		181
SUTTON	32, 324	
<i>Sutton Beauty</i> (synonym of <i>Dumelow</i>)	112	
<i>Sutton Beauty</i> (synonym of <i>Sutton</i>)	324	
<i>Swinetka</i>		121
SWAAR	15, 16, 17, 18, 326	
<i>Swaar Apple</i>	326	
<i>Swazyie Pomme Gris</i>	328	
<i>Swazyie</i>	328	
<i>Swazyie Pomme Grise</i>	328	
SWAZIE	328	
<i>Swazie Pomme Grise</i>	328	
<i>Swazie's Pomme Gris</i>	328	
<i>Swazy</i>	328	

COMBINED INDEX TO VARIETIES.

351

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Swazy Pomme Gris</i>	328	
<i>Swedish Borsdorf of Patten</i>		5
SWEET AND SOUR	329	
SWEET BOUGH	12, 15,	216
SWEET FALL PIPPIN		218
<i>Sweet Golden Pippin</i>		12
SWEET GREENING	151, 331	
<i>Sweet Harvest</i>		217
<i>Sweet June</i>		98
SWEET KING	333	
<i>Sweet Lyman's Pumpkin</i>		172
<i>Sweet Maiden's Blush</i>	91	
<i>Sweet Pearmain</i>	333	
<i>Sweet Pippin</i> (synonym of Moore Sweet).....	219	
<i>Sweet Pippin</i> (synonym of Hog Island Sweet).....		100
<i>Sweet Rambo</i>		89
<i>Sweet Rhode Island Greening</i>	331	
SWEET RUSSET	333	
SWEET RUSSET		218
<i>Sweet Russet</i> (synonym of Pumpkin Russet).....		170
<i>Sweet Russet</i> (synonym of Cheeseboro).....		33
<i>Sweet's Harvest</i>		212
<i>Sweet Sugar</i>		12
<i>Sweet Wine</i>		64
SWEET WINESAP	333	
<i>Sweet Wine Sap</i>	333	
SWENKER	335	
<i>Swett's Harvest</i>		212
<i>Swinec</i>		121
<i>Swinzovsha</i>		121
SWITZER		218
<i>Sylvan Russet</i>	294	
SYLVESTER		220
<i>Taffalai</i>		239
<i>Tallman's Sweet</i>	343	
<i>Tallman Sweet</i>	343	
<i>Tallman Sweeting</i>	343	
<i>Tallow</i>		128
<i>Tallow Apple</i>		128
<i>Tallow Pippin</i>		128
<i>Talman's Sweet</i>	343	
<i>Talman Sweet</i>	343	
<i>Talman's Sweeting</i>	343	
<i>Talman Sweeting</i>	343	
<i>Tars Thorn</i>		45
<i>Tart Bough</i> (synonym of Early Harvest).....		51
<i>Tart Bough</i> (synonym of Champlain).....		30
TART BOUGH		220

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Teignmouth</i>		223
<i>Teller</i>	206	
<i>Tennessee Early Red</i>		55
<i>Tenon Hills</i>	67	
<i>Terry's Redstreak</i>	273	
<i>Tete d'Ange</i>		29
<i>Tete de Chat</i>		29
<i>Tetoffsky</i>		220
<i>Tetofski</i> (synonym of <i>July</i>)		112
<i>Tetofski</i> (synonym of <i>Tetofsky</i>)		220
TETOFSKY	25, 26	220
<i>Tetofsky</i> (synonym of <i>July</i>)		112
<i>Tewkesbury Blush</i>	336	
<i>Tewkesbury Winter Blush</i>	336	
<i>Tewksberry Winter Blush</i>	336	
Tewksbury	336	
<i>Tewksbury Blush</i>	336	
<i>Tewksbury Winter Blush</i>	336	
<i>Texan Red</i>	375	
TEXAS	337	
THALER		222
<i>Thomas</i>	208	
THOMPSON		222
<i>Thompson No. 24</i>		222
<i>Thompson's Seedling No. 24</i>		222
<i>Thompson's Seedling No. 29</i>		111
<i>Thompson's Seedling No. 38</i>		90
<i>Timothy</i>	339	
<i>Timothy Titus Sort</i>	339	
TINMOUTH		223
<i>Tinmouth Sweet</i>		223
<i>Titacca</i>		224
TITOVKA		224
<i>Titovka</i>		224
TITUS	339	
<i>Titus</i> (synonym of <i>Titus Pippin</i>)	339	
<i>Titus Apple</i>		224
TITUS PIPPIN	338	
<i>Titus Riga</i>		224
TOBIAS	340	
<i>Tobias Apple</i>	340	
TOBIAS BLACK	341	
TOBIAS PIPPIN	342	
<i>Tolman</i>	343	
TOLMAN Sweet	16, 17, 18, 343	
<i>Tolman's Sweetling</i>	343	
<i>Toma Red</i>	345	
<i>Tom Harryman</i>		85

COMBINED INDEX TO VARIETIES.

353

	VOL. I PAGE.	VOL. II PAGE.
<i>Tommy Red</i>	345	
<i>Tompkins</i>		49
<i>Tompkins County King</i>	345	
TOMPKINS KING	17, 18, 19, 345	
<i>Tom Put</i>		225
TOM PUT		245
<i>Tom's Red</i>	345	
<i>Tom Woodward Pippin</i>	244	
<i>Transcendant (Crab)</i>		267
TRANSCENDENT (Crab)		266
<i>Transparente de Astracan</i>		239
<i>Transparente d'Ete</i>		239
<i>Transparent de Muscovie</i>		239
<i>Transparente de Muscovie d'Ete</i>		239
<i>Transparente de Saint-Leger (syn. of Vineuse Rouge)</i>		233
<i>Transparente de Saint-Leger (syn. of Yellow Transparent)</i>		248
<i>Transparente de Zurich</i>		230
<i>Transparente Jaune (synonym of Vineuse Rouge)</i>		233
<i>Transparente Jaune (synonym of Yellow Transparent)</i>		248
<i>Transparente Rouge</i>		233
<i>Transparente Verte</i>		233
<i>Transparent Muscovie</i>		230
<i>Travers</i>		184
<i>Travers Apple</i>		184
<i>Travers Pippin</i>		184
<i>Travers Pippin</i>		184
<i>Travers Renette</i>		184
<i>Trenham</i>	227	
<i>Trenham</i>	227	
<i>Trenton Early</i>		81
<i>True Spitzenburgh</i>	121	
<i>Trumpington</i>	273	
<i>Tsarskui Schip</i>		45
TUFTS	24	226
<i>Tufts Baldwin</i>		226
<i>Tufts Seedling</i>		226
<i>Tulpahocken</i>	125	
<i>Tulpehocken</i>	125	
<i>Turn Off Lane</i>		174
TWENTY OUNCE	17, 18, 193	227
TWENTY OUNCE PIPPIN	349	
<i>Twenty Ounce Pippin (synonym of Twenty Ounce)</i>		227
<i>Twitty's Paragon</i>	247	
TYRE BRUITY		229
<i>Ulsie Seedling</i>	172	
<i>Uncle Sam's Best</i>		64
<i>Underdunk</i>		30
UTTER		229

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Utter's Large Red</i>		230
<i>Utter's Red</i>		230
<i>Valandingham</i>	197	
<i>Vandevere</i> (synonym of Newtown Spitzenburg)	225	
<i>Vandevere</i> (synonym of Smokehouse)	312	
<i>Vandevere</i> of New York	225	
<i>Vandevere</i> (synonym of Vandevere)		230
<i>Vandevere</i> (synonym of Vandevere Pippin)		231
<i>Vandevere Pippin</i>		231
<i>Vandever</i>		230
VANDEVERE	18	230
<i>Vandevere</i> (synonym of Newtown Spitzenburg)	225	
<i>Vandevere, English</i>	312	
<i>Vandevere</i> of New York	225	
VANDEVERE PIPPIN		231
<i>Vandevere Yellow</i>		231
<i>Van Dym's Pippin</i>		62
<i>Van Dyne</i>	244	
<i>Van Dyne</i>	244	
<i>Van Dyne Apple</i>		77
<i>Van Dyn's Pippin</i>		62
VANHOY	351	
<i>Van Hoy</i>	351	
<i>Van Hoy No Core</i>	351	
<i>Van Kleeck's Sweet</i>		100
<i>Van Fleet</i>	161	
VAN WYCK (Crab)		267
<i>Van Wyck</i> (synonym of Brier)		251
<i>Van Wyck Siberian</i>		268
<i>Van Wyck Sweet</i>		268
<i>Vargul</i>		6
<i>Varick</i>	105	
<i>Vermillon d'Ete</i>		178
<i>Vermont</i>		30
<i>Vermont Pippin</i>		223
<i>Vermont Pumpkin Sweet</i>		172
<i>Vermont Sweet</i>		172
<i>Verte de l'Île de Rhodes</i>	283	
<i>Verte de Rhode Island</i>	283	
<i>Vickers</i>	186	
<i>Victoire, La.</i>	188	
VICTORIA	24	231
<i>Victoria Pippin</i>	69	
<i>Victoria Red</i>	69	
<i>Victoria Sweet</i>		231
<i>Victoria Sweetling</i>		231
VICTUALS AND DRINK		233
<i>Vilikui Mogul</i>		87

COMBINED INDEX TO VARIETIES.

355

	VOL. I. PAGE.	VOL. II. PAGE.
VINEUSE ROUGE		233
<i>Vinnoe Krasnoe Oskunee</i>		182
<i>Vinograd</i>		244
VIRGINIA GREENING	352	
<i>Virginia June</i>		142
<i>Virginia Pippin</i> (synonym of Newtown Pippin)	146	
<i>Virginia Pippin</i> (synonym of Virginia Greening)	352	
<i>Virginia Sweet</i>		177
<i>Peromesh No. 21</i>		247
WABASH	353	
<i>Wabash Bellflower</i> (synonym of Long Red Pearmain) ...		125
<i>Wabash Bellflower</i> (synonym of Wabash)	353	
WABASH RED	353	
<i>Wabash Red Winter</i>	353	
WAGNER	21, 24, 354	
<i>Wahr Reinette</i>	94	
<i>Walsh</i>	227	
WALBRIDGE	356	
<i>Walker</i>	358	
WALKER BEAUTY	358	
<i>Walker's Winter</i>	358	
<i>Walker Yellow</i>	358	
<i>Wall</i>	227	
WALLACE HOWARD	359	
<i>Wallbridge</i>	357	
<i>Walter Pease</i>		161
<i>Waltz Apple</i>	253	
<i>Wakworth</i>		30
<i>Wander</i>	227	
WANDERING SPY	360	
<i>Warden's Pie Apple</i>		203
<i>Ward's Pippin</i>		21
<i>Warren Pennock</i>		53
<i>Warren Pippin</i> (synonym of Ortley)	244	
<i>Warren Pippin</i> (synonym of Yellow Bellflower)	381	
<i>Washington</i> (synonym of Sops of Wine)		203
<i>Washington</i> (synonym of Washington Strawberry)		234
<i>Washington</i> (synonym of Sweet Bough)		217
<i>Washington County Seedling</i>		214
<i>Washington of Maine</i>		234
WASHINGTON ROYAL	361	
WASHINGTON STRAWBERRY		234
WATER		236
<i>Waterloo</i>		178
<i>Watermelon</i>	204	
<i>Watkins Early</i>		211
<i>Watson's Vandercere</i>		231
<i>Watson's Vandercere</i>		231

	VOL. I. PAGE.	VOL. II. PAGE.
Watts Apple	253	
WATWOOD	362	
Waxen	67	
Waxen Apple	67	
WEALTHY	17, 19, 25	236
WEALTHY GROUP	25	
Weinappel Rother		182
Weinappel Rother		182
Weiser Hachthornden		96
Weisser Metzgerapfel	381	
Well Apple (synonym of Domine)	109	
Well Apple (synonym of Titus Pippin)	339	
Wellington	112	
Wellington's Reinette	112	
WELLS	363	
Wells (synonym of Domine)	109	
Wells (synonym of Ohio Nonpareil)		148
Wells Apple	363	
Wesse Antillische W'interreinette	94	
Westbrook		60
Westchester Seek-No-Further	130	
Western Baldwin (synonym of Babbitt)	53	
Western Baldwin (synonym of Pawpaw)	259	
WESTERN BEAUTY		239
Western Beauty (synonym of Hyde King)	166	
Western Beauty (synonym of Grosh)		89
Western Beauty (synonym of Ohio Nonpareil)		148
Westfield	364	
Westfield Seek-No-Further	364	
WESTFIELD Seek-No-Further	12, 16, 17, 18, 32, 364	
WESTFIELD Seek-No-Further; Red type	366	
Wheeters Kernel		96
White	67	
White Apple (synonym of Hawthornden)		96
White Apple (synonym of Parry White)		157
White Astrucan		239
WHITE ASTRUCAN	25	239
White Bellefleur	244	
White Bellflower	244	
White Borodocka		15
White Crow	197	
White Detroit	244	
White Graft of Wisconsin		60
White Hawthornden		96
White Hawthornden		96
White June		118
WHITF JUNEATING		240
White Newell		60

COMBINED INDEX TO VARIETIES.

357

	VOL. I. PAGE.	VOL. II. PAGE.
WHITE PEARMAIN	367	
WHITE PIPPIN	24, 368	
<i>White Pippin</i> (synonym of Canada Reinette)	94	
<i>White Pippin</i> (synonym of Ortley)	244	
<i>White Robinson</i>	380	
WHITE SEEK-NO-FURTHER	15	
<i>White Seek-No-Further</i> (synonym of Green Seek-No-Further)		88
WHITE SPANISH REINETTE		241
<i>White Spice</i>		49
<i>White Winter Pearmain</i>	367	
<i>White Zurel</i>	387	
WHITNEY (Crab)		268
<i>Whitney No. 20</i>		268
<i>Willcox's Winter</i>	302	
<i>Wild Apple</i>	384	
WILLIAMS		242
<i>Williams Early</i>		242
<i>Williams Early Red</i>		242
<i>Williams Favorite</i>		242
<i>Williams Favorite Red</i>		242
<i>Williamson</i>	109	
<i>Williams Red</i>		242
WILKES SWEET		244
<i>Wilkes Sweeting</i>		244
WILLOW	18, 370	
<i>Willow Leaf</i>	370	
<i>Willow Leaf Pippin</i>	244	
<i>Willow Twig</i>	370	
WILLSBORO	372	
<i>Willsons June</i>		181
<i>Windover</i>		231
WINDSOR	372	
<i>Windsor Chief</i>	372	
WINE	15, 373	
<i>Wine</i> (synonym of Jonathan)	172	
<i>Wine</i> (synonym of Newtown Spitzenburg)	225	
<i>Wine</i> (synonym of Egg Top)		56
<i>Wine</i> (synonym of Fall Wine)		64
<i>Wine</i> (synonym of Richard Graft)		186
<i>Wine</i> (synonym of Twenty Ounce)		227
<i>Wine Apple</i>	373	
<i>Wine of Cole</i>		64
WINE RUBES		244
WINESAP	17, 24, 374	
WINESAP GROUP	24	
<i>Winesap</i> (synonym of Jonathan)	172	
<i>Winesap</i> (synonym of Roseau)	292	

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Wine Sop</i>	375	
<i>Wine Sweet</i>	380	
WINTER BANANA	377	
<i>Winter Belle Bonne</i>	67	
<i>Winter Blush</i> (synonym of Fallawater)	125	
<i>Winter Blush</i> (synonym of Rock Pippin of Eastern New York)	288	
<i>Winter Chandler</i>		31
<i>Winter Genneting</i>	271	
<i>Winter Golden Sweet</i>		13
WINTER HOG ISLAND SWEET	378	
<i>Winter Horse</i>	227	
<i>Winter Jannetting</i>	271	
<i>Winter King</i>	345	
<i>Winter Nonsuch</i>	276	
<i>Winter Peach</i>	251	
WINTER PEARMAN	378	
<i>Winter Pearmain</i> (synonym of McAfee)	197	
<i>Winter Pearmain</i> (synonym of Milam)	208	
<i>Winter Pearmain</i> (synonym of Long Island Pearmain) ...		124
<i>Winter Pearmain</i> (synonym of Long Red Pearmain)		125
<i>Winter Pippin</i>	197	
<i>Winter Pippin of Geneva</i>	136	
<i>Winter Queen</i>	88	
<i>Winter Quening</i>	88	
<i>Winter Rose</i>	227	
<i>Winter Russet</i> (synonym of English Russet)	118	
<i>Winter Russet</i> (synonym of Perry Russet)	257	
WINTER ST. LAWRENCE	379	
<i>Winter Seek-No-Further</i>		88
<i>Winter Sweet Paradise</i>	380	
<i>Winter Wine</i>	373	
WINTHROP GREENING		245
WISMER	380	
<i>Wisner's Dessert</i>	380	
<i>Wisner's Dessert</i>	380	
WOLF RIVER	20, 25	245
<i>Wolf River</i> (synonym of Alexander)		3
<i>Wolman's Harvest</i>		215
<i>Wonder</i>	227	
<i>Woodman's Song</i>	244	
<i>Woodpecker</i>	56	
<i>Woodstock</i> (synonym of Blenheim)		21
<i>Woodstock</i> (synonym of Dyer)		49
<i>Woodstock Pippin</i>		21
<i>Woodward's Pippin</i>	244	
<i>Woolman's Early</i>		215
<i>Woolman's Harvest</i>		215

COMBINED INDEX TO VARIETIES.

359

	VOL. I. PAGE.	VOL. II. PAGE.
<i>Woolman's Long</i>	244	
<i>Woolman's Long Pippin</i>	244	
<i>Woolman's Striped Harvest</i>		215
<i>Woolnary Long</i>	244	
<i>Woolventon</i>		127
WORKABOE		240
<i>World's Wonder</i>	227	
<i>Wunderappel</i>		3
<i>Wyandotte</i>	196	
<i>Yankee Apple</i>		172
<i>Yeats</i>	310	
<i>Yellow Bellefleur</i>	381	
<i>Yellow Belleflower</i>	381	
YELLOW BELLFLOWER	15, 16, 17, 18, 19, 23,	381
YELLOW BELLFLOWER GROUP		383
<i>Yellow Bough</i>		217
YELLOW CALVILLE		247
<i>Yellow Flat</i>		118
YELLOW FOREST		384
<i>Yellow German Reinette</i>	142	
YELLOW HARVEST	12	
<i>Yellow Harvest</i> (synonym of Early Harvest)		51
<i>Yellow Junett</i>	271	
<i>Yellow June</i> (synonym of White Juneating)		240
<i>Yellow June</i> (synonym of Kirkbridge)		118
<i>Yellow Juneating</i>		51
<i>Yellow May</i>		240
<i>Yellow Newton's Pippin</i>	140	
YELLOW NEWTOWN	17, 18, 24, 145, 150,	385
<i>Yellow Newtown</i> (synonym of Canada Reinette)	94	
<i>Yellow Newtown Pippin</i>	146	
<i>Yellow Pippin</i> (synonym of Newark Pippin)	223	
<i>Yellow Pippin</i> (synonym of Ortley)	444	
YELLOW SIBERIAN (Crab)		260
<i>Yellow Summer Pearmain</i>		160
<i>Yellow Sweeting</i>		81
YELLOW TRANSPARENT	17, 25	247
<i>Yellow Transparent</i> (synonym of Thaler)		222
<i>Yellow Vanderwe</i>		231
YOPP	18	249
<i>Yopp's Favorite</i> (Synonym of Yopp)		249
<i>Yopp's Favorite</i> (synonym of Yellow Newtown)	146	
YORK		249
<i>York and Lancaster</i>		193
YORK IMPERIAL	18, 19, 32,	385
YORK PIPPIN	24	
<i>York Pippin</i> (synonym of Fall Pippin)		62
<i>York Pippin</i> (synonym of Golden Pippin I)		78

	VOL. I. PAGE.	VOL. II. PAGE.
<i>York Pippin</i> (synonym of White Spanish Reinette)		241
<i>York Russet</i> (synonym of Cheeseboro)		33
<i>York Russet</i> (synonym of Pumpkin Russet)		170
<i>York Russeting</i>		33
<i>Zarskischip</i>		45
<i>Zarski Schip</i>		45
<i>Zarski Zars</i>		45
<i>Zeeke</i>	197	
<i>Zolotoi Renet</i>		79
<i>Zour Bough</i>		168
ZURBEL	387	
<i>Zusoff</i> of Tuttle		5

CPSIA information can be obtained at www.ICGtesting.com
Printed in the USA
LVOW080018060912

297478LV00004B/57/A





WWW.KESSINGER.NET



Cover design: GBRCreations.com